



Mark Scheme (Results)

Summer 2025

Pearson Edexcel GCE
In Further Mathematics (8FM0)
Paper 24 Further Statistics 2

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2025

Question Paper Log Number P76374A

Publications Code 8FM0_24_2506_MS

All the material in this publication is copyright

© Pearson Education Ltd 2025

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 40.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol $\checkmark$ will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - $\square$ The second mark is dependent on gaining the first mark
4. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 5. Where a candidate has made multiple responses and indicates which response they wish to submit, examiners should mark this response.
If there are several attempts at a question which have not been crossed out, examiners should mark the final answer which is the answer that is the most complete.
 6. Ignore wrong working or incorrect statements following a correct answer.
 7. Mark schemes will firstly show the solution judged to be the most common response expected from candidates. Where appropriate, alternatives answers are provided in the notes. If examiners are not sure if an answer is acceptable, they will check the mark scheme to see if an alternative answer is given for the method used.

Question	Scheme	Marks	AOs
2(a)	0	B1	1.1b
		(1)	
(b)	$E(X) [= \frac{-3+2}{2}] = -\frac{1}{2}$	B1	1.1b
		(1)	
(c)	$\text{Var}(X) [= \frac{(2-(-3))^2}{12}] = \frac{25}{12}$	B1	1.1b
		(1)	
(d)	Sketches straight line with positive gradient	M1	2.1
	Labels $(-3, 0)$ and ends graph at $(2, 1)$	A1	1.1b
	Labels $(0, 0.6)$	A1	1.1b
		(3)	
(e)	$X > 1.4$ or $X < -1.4$	M1	1.1b
	$P(X > 1.4) + P(X < -1.4) = \frac{2-1.4}{2-(-3)} + \frac{-1.4-(-3)}{2-(-3)} \text{ or } \frac{0.6}{5} + \frac{1.6}{5} \text{ oe}$	M1	3.1a
	$\frac{11}{25}$	A1	1.1b
		(3)	
(9 marks)			
Notes:			
(a)	B1:	cao	
(b)	B1:	o.e.	
(c)	B1:	o.e. (allow awrt 2.08)	
(d)	M1:	Sketching straight line with positive gradient.	
	A1:	Correct label and domain	
	A1:	Correct label on vertical axis	
(e)	M1:	Either correct region	
	M1:	Attempt to find probability using both regions	
	A1:	o.e. e.g. 0.44	
NB		Stating $P(-1.4 < X < 1.4) = 0.56$ can score M1A1A0	

Question	Scheme		Marks	AOs
3(a)	$S_{xy} = \sum xy - \frac{(\sum x \sum y)}{20} = 2056.63 - \frac{(19.8 \times 20) \times (5.2 \times 20)}{20} = -2.57^*$		B1cso*	1.1b
			(1)	
(b)	$r = \frac{-2.57}{\sqrt{15.78 \times 0.94}}, = -0.667291... = \text{awrt } \underline{\underline{-0.667}}$		M1, A1	1.1b 1.1b
			(2)	
(c)	$H_0 : \rho = 0 \quad H_1 : \rho < 0$		B1	2.5
	5% one-tail cv for r is: $(\pm)0.3783$		M1	1.1b
	Significant result so there is evidence of negative correlation (between BMI and speed)		A1	2.2b
			(3)	
(d)	$b = \frac{'-2.57'}{15.78} [= -0.16286...]$		M1	3.3
	$a = 5.2 - 'b'(19.8)$		M1	1.1b
	$v = 8.42 - 0.16x$		A1	1.1b
			(3)	
(e)	$RSS = S_{yy} \times (1 - r^2)$ or $S_{yy} - \frac{(S_{xy})^2}{S_{xx}}, = 0.521... = \text{awrt } \underline{\underline{0.52}}$		M1, A1	1.1b 1.1b
			(2)	
(f)	$v_A = 8.42 - 0.16(20.9) + 0.18$ $v_B = 8.42 - 0.16(19.7) - 0.22$		M1	3.4
	$v_A = \text{awrt } 5.2 \text{ or awrt } 5.3 \quad v_B = \text{awrt } 5.0$		A1	1.1b
	$v_A > v_B$ therefore A took the shorter amount of time.		A1	2.2a
			(3)	
(14 marks)				
Notes:				
(a)	B1*:	complete calculation (oe) for S_{xy} leading to given answer. Allow ... $-\frac{396 \times 104}{20}$		
(b)	M1:	use of formula for r with all given values		
	A1:	awrt -0.667		
(c)	B1:	for both hypotheses correct in terms of ρ		
	M1:	for use of tables to find the cv of 0.3783 (Condone 0.378)		
	A1:	for a correct conclusion that mentions negative correlation. (B0M1A1 possible)		
NB		A comparison is not required but if seen it must be correct for the A1		
(d)	M1:	setting up linear model by finding gradient		
	M1:	attempting v -intercept of linear model		
	A1:	correct model with $b = \text{awrt } -0.16$ and $a = \text{awrt } 8.42$ (NB 8.43 is A0 here)		
(e)	M1:	attempt at either correct expression		
	A1:	awrt 0.52		
(f)	M1:	using the linear model to find the speed of either runner. Must <u>see use</u> of residuals attempted but condone wrong sign.		
	A1:	both speeds found correctly. Can allow 2 nd A1 for slightly incorrect speeds M1A0A1		
	A1:	deducing that A took the shortest time (dep on M1 and consistent with speeds found)		

Question	Scheme	Marks	AOs
4(a)	$F(2) \rightarrow a(2^4) - \frac{1}{8}(2^3) + b(2^2)$	M1	3.4
	$16a + 4b - 1$	A1	1.1b
		(2)	
(b)	$f(t) = \frac{d}{dt} \left(at^4 - \frac{1}{8}t^3 + bt^2 \right) = 4at^3 - \frac{3}{8}t^2 + 2bt$	M1	1.1b
	$f(t) = \begin{cases} 4at^3 - \frac{3}{8}t^2 + 2bt & 0 \leq t \leq 4 \\ 0 & \text{otherwise} \end{cases}$	A1	1.1b
		(2)	
(c)	$F(4) = 1 \rightarrow 256a - 8 + 16b = 1$	M1	3.1b
	Solving simultaneously $256a + 16b = 9$ $16a + 4b = \frac{27}{16}$	M1	1.1b
	$a = \frac{3}{256}$ $b = \frac{3}{8}$	A1	1.1b
	$\frac{d}{dt}(f(t)) = 12at^2 - \frac{3}{4}t + 2b$	M1	3.1b
	$\frac{9}{64}t^2 - \frac{3}{4}t + \frac{3}{4} = 0$ or e.g. $3t^2 - 16t + 16 = 0$	M1	2.1
	$t = \frac{4}{3}$ or $t = 4$	M1	1.1b
	$[t = 4 \text{ is a minimum and } f(0) = 0 \text{ so}] \text{ mode of } T \text{ is } \frac{4}{3}$	A1	3.2b
		(7)	
(11 marks)			
Notes:			
(a)	M1:	Use of $F(2)$ but e.g. $F(4) - F(2)$ is M0	
	A1:	cao	
(b)	M1:	attempt to differentiate $F(t)$ with at least one term correct	
	A1:	both lines correct with correct limits	
(c)	M1:	use of $F(4) = 1$	
	M1:	solving simultaneously. Method to eliminate one variable or implied by A1	
	A1:	both values correct	
	M1:	differentiating $f(t)$ ft their a and b values. Allow with letters a and b condone 1 slip	
	M1:	(dep on 3 rd M1) setting equal to 0	
	M1:	solving quadratic. Method must be seen or implied by the correct answer.	
	A1:	selecting $\frac{4}{3}$ as only solution	

