



# Mark Scheme (Results)

Summer 2025

Pearson Edexcel International Advanced Level  
In Further Pure Mathematics F2 (WFM02)  
Paper 01

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2025

Question Paper Log Number P76392A

Publications Code WFM02\_01\_2506\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2025

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## General Instructions for Marking

The total number of marks for the paper is 75.

Edexcel Mathematics mark schemes use the following types of marks:

### 'M' marks

These are marks given for a correct method or an attempt at a correct method. In Mechanics they are usually awarded for the application of some mechanical principle to produce an equation, e.g. resolving in a particular direction; taking moments about a point; applying a suvat equation; applying the conservation of momentum principle; etc.

The following criteria are usually applied to the equation.

To earn the M mark, the equation

- (i) should have the correct number of terms
- (ii) each term needs to be dimensionally correct

For example, in a moments equation, every term must be a 'force x distance' term or 'mass x distance', if we allow them to cancel 'g' s.

For a resolution, all terms that need to be resolved (multiplied by sin or cos) must be resolved to earn the M mark.

'M' marks are sometimes dependent (DM) on previous M marks having been earned, e.g. when two simultaneous equations have been set up by, for example, resolving in two directions and there is then an M mark for solving the equations to find a particular quantity – this M mark is often dependent on the two previous M marks having been earned.

### 'A' marks

These are dependent accuracy (or sometimes answer) marks and can only be awarded if the previous M mark has been earned. e.g. M0 A1 is impossible.

### 'B' marks

These are independent accuracy marks where there is no method (e.g. often given for a comment or for a graph).

A and B marks may be f.t. – follow through – marks.

## General Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes:

- bod means benefit of doubt
- ft means follow through
  - o the symbol  $\checkmark$  will be used for correct ft
- cao means correct answer only
- cso means correct solution only, i.e. there must be no errors in this part of the question to obtain this mark
- isw means ignore subsequent working

- awrt means answers which round to
- SC means special case
- oe means or equivalent (and appropriate)
- dep means dependent
- indep means independent
- dp means decimal places
- sf means significant figures
- \* means the answer is printed on the question paper
- $\square$  means the second mark is dependent on gaining the first mark

All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

If a candidate makes more than one attempt at any question:

- If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
- If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.

Ignore wrong working or incorrect statements following a correct answer.

## General Principles for Further Pure Mathematics Marking

*(NB specific mark schemes may sometimes override these general principles)*

### Method mark for solving 3 term quadratic:

- Factorisation
  - o  $(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$
  - o  $(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$
- Formula
  - o Attempt to use the correct formula (with values for  $a$ ,  $b$  and  $c$ ).
- Completing the square
  - o Solving  $x^2 + bx + c = 0$  :  $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$ ,  $q \neq 0$ , leading to  $x = \dots$

### Method marks for differentiation and integration:

- Differentiation
  - o Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )
- Integration
  - o Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

### Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first. Normal marking procedure is as follows:

- Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.
- Where the formula is not quoted, the method mark can be gained by implication from correct working with values but may be lost if there is any mistake in the working.

### Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

### Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.

**June 2025**  
**WFM02 Further Pure Mathematics F2**  
**Mark Scheme**

| Question Number | Scheme                                                                                                                                                         | Notes                                                                                                                                                                                                 | Marks |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <b>1(a)</b>     | $r = 4\sqrt{2}$                                                                                                                                                | Correct $r$<br>Accept $\pm 4\sqrt{2}$ or $\pm\sqrt{32}$ or other processed equivalents.                                                                                                               | B1    |
|                 | $\theta = -\pi + \arctan\left(\frac{2\sqrt{6}}{2\sqrt{2}}\right) = -\pi + \frac{\pi}{3} = -\frac{2\pi}{3}$                                                     | Complete method to find an argument in the correct quadrant for their $r$                                                                                                                             | M1    |
|                 | $\Rightarrow -2\sqrt{2} - (2\sqrt{6})i = 4\sqrt{2}e^{-\frac{2\pi}{3}i}$                                                                                        | Correct answer in the form specified. Accept with $\sqrt{32}$                                                                                                                                         | A1    |
| <b>(3)</b>      |                                                                                                                                                                |                                                                                                                                                                                                       |       |
| <b>(b)</b>      | $z^5 = -2\sqrt{2} - (2\sqrt{6})i$                                                                                                                              |                                                                                                                                                                                                       |       |
|                 | $z = (4\sqrt{2})^{\frac{1}{5}} e^{-\frac{2\pi + 2k\pi}{5}i}$                                                                                                   | Attempts De Moivre correctly with either all roots indicated or simplified $r$ with one argument. See additional notes.                                                                               | M1    |
|                 | $\sqrt{2}e^{-\frac{2\pi}{15}i}, \sqrt{2}e^{\frac{4\pi}{15}i}, \sqrt{2}e^{\frac{2\pi}{3}i},$<br>$\sqrt{2}e^{-\frac{8\pi}{15}i}, \sqrt{2}e^{-\frac{14\pi}{15}i}$ | Three correct roots provided coefficients of $\pi$ for $\theta$ are single fractions, allow for $0 < \theta < 2\pi$<br>Allow $(4\sqrt{2})^{\frac{1}{5}}$ or $\sqrt{2}$ and correct roots in trig form | A1    |
|                 |                                                                                                                                                                | All five correct roots and no extra. Must have $-\pi < \theta \leq \pi$ . Allow e.g., $\sqrt{2}e^{\frac{10\pi}{15}i}$ for $\sqrt{2}e^{\frac{2\pi}{3}i}$                                               | A1    |
| <b>(3)</b>      |                                                                                                                                                                |                                                                                                                                                                                                       |       |
| <b>Total 6</b>  |                                                                                                                                                                |                                                                                                                                                                                                       |       |

Additional Notes:

(a)

B1: Accept positive or negative root. Allow if seen in part (b).

M1: Requires a method to achieve an argument in the correct quadrant for their  $r$ . Accept attempts at

$\arctan\left(\frac{2\sqrt{6}}{2\sqrt{2}}\right) \pm \pi$  or  $\arctan\left(\frac{2\sqrt{2}}{2\sqrt{6}}\right) \pm \pi$  for attempts where  $r > 0$ , or equivalent methods via arccos or arcsin

(must use  $r$  as the denominator). May be implied by sight of any of  $-\frac{2\pi}{3}, \frac{4\pi}{3}, \frac{7\pi}{6}$  or  $-\frac{5\pi}{6}$  if no method is shown.

In cases where a negative  $r$  is given, accept attempts at just  $\arctan\left(\frac{2\sqrt{6}}{2\sqrt{2}}\right)$  or  $\arctan\left(\frac{2\sqrt{2}}{2\sqrt{6}}\right)$  provided they

do not subsequently adjust to a different quadrant.

A1: Correct answer in the form specified, with  $-\pi < \theta < \pi$ . Must be given in correct form, but accept

answer with negative  $r$ , i.e.  $-4\sqrt{2}e^{\frac{\pi}{3}i}$  Accept with exact processed equivalents for  $r$ .

Note: Correct answer with no working shown will score B1M1(implied)A1.

(b)

M1: Applies de Moivre correctly with their answer to (a) which could be in trig form, to reach either simplified modulus with one correct argument for their (a), or multiple arguments with unsimplified modulus.

Condone missing "i" for the M.

Note: If  $\pm 2k\pi$  is not used or is used incorrectly this mean you can score M1A0A0 for just  $\sqrt{2}e^{-\frac{2\pi}{15}i}$

A1: Accept for three correct distinct roots allowing arguments in the range  $-\pi < \theta < \pi$  or  $0 < \theta < 2\pi$  for this mark. The modulus need not be simplified for this mark.

A1: All 5 correct roots in the form specified.

| Question Number | Scheme                                                                                                                                                   | Notes                                                                                         | Marks          |
|-----------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------|----------------|
| 2               | $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} - 5y^2 = e^{x^2-9}$                                                                                                  |                                                                                               |                |
|                 | $\frac{d^3y}{dx^3} - 4\frac{d^2y}{dx^2} - 10y\frac{dy}{dx} = 2xe^{x^2-9}$                                                                                | M1: Differentiates to obtain three correct terms<br>A1: Fully correct differentiated equation | M1 A1          |
|                 | $y(3) = 2, y'(3) = -1$<br>$y''(3) - 4(-1) - 5 \times 2^2 = 1 \Rightarrow y''(3) = 17$<br>$y'''(3) - 4(17) - 10(2)(-1) = 6 \Rightarrow y'''(3) = 54$      | Uses the given equation and their equation to obtain values for $y''(3)$ and $y'''(3)$        | M1             |
|                 | $(y =) y(3) + y'(3)(x-3) + \frac{y''(3)}{2}(x-3)^2 + \frac{y'''(3)}{3!}(x-3)^3 \Rightarrow$<br>$2 - 1(x-3) + \frac{17}{2}(x-3)^2 + \frac{54}{3!}(x-3)^3$ | A correct unsimplified Taylor series for their $y''(3)$ and $y'''(3)$                         | M1             |
|                 | $(y =) 2 - (x-3) + \frac{17}{2}(x-3)^2 + 9(x-3)^3$                                                                                                       | Correct simplified series. "y = ..." is not required                                          | A1             |
|                 |                                                                                                                                                          |                                                                                               | (5)            |
|                 |                                                                                                                                                          |                                                                                               | <b>Total 5</b> |

#### Additional Notes

M1: Attempts to differentiate using the chain rule at least once – at least three terms correct, allowing for slips in rearranging before differentiation.


A1: Any correct third derivative expression. They may have rearranged first.

M1: Finds the values of the second and third derivatives using the given equation and their third derivative expression (which must have at least three terms). Condone one slip in substitutions as long as evidence of the correct values used is seen. If substitution is not seen, correct values can imply the method.

M1: Applies the Taylor Series formula correctly with the given values of  $y$  and  $\frac{dy}{dx}$  and with their values for

the second and third derivatives – the first two terms must be correct. If the values of their second and third derivatives are not explicitly stated but embedded in the series, it must be clear that a correct attempt at the series has been used – the denominators 2 and 3! (or 6) must be seen – but their values need not be correct. Condone a miscopy of one of **their** values if it is clearly a correct attempt to apply their value to the series.

A1: Correct fully simplified series expansion. The  $y =$  is not required and condone if  $f(x)$  is used.

| Question Number                                                    | Scheme                                                                                                                                                              | Notes                                                                                                                                                                                                                                                                                     | Marks |
|--------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| 3(a)                                                               |                                                                                    | Circle centred at origin with radius of 1 correctly indicated in any way.<br>May use $\pm i$ or $\pm 1$ in appropriate position on imaginary axis.                                                                                                                                        | B1    |
| <b>(1)</b>                                                         |                                                                                                                                                                     |                                                                                                                                                                                                                                                                                           |       |
| <b>(b)</b>                                                         | $w = \frac{9iz - i}{z + 1}$ $\Rightarrow wz + w = 9iz - i \Rightarrow z(w - 9i) = -w - i$ $\Rightarrow z = \frac{-w - i}{w - 9i} = \frac{w + i}{9i - w} \text{ oe}$ | M1: Makes $z$ the subject<br>A1: Correct expression oe                                                                                                                                                                                                                                    | M1 A1 |
|                                                                    | $ z  = \left  \frac{-w - i}{w - 9i} \right  \Rightarrow  w - 9i  =  w - (-i) $                                                                                      | Uses $ z  = 1$ and obtains<br>$ \pm w \pm z_1  =  \pm w \pm z_2 $<br><b>Requires previous M mark.</b>                                                                                                                                                                                     | dM1   |
|                                                                    | $\Rightarrow v = "4" \quad \text{or} \quad \begin{aligned} u^2 + (v - 9)^2 &= u^2 + (v + 1)^2 \\ -18v + 81 &= 2v + 1 \\ 20v &= 80 \\ v &= 4 \end{aligned}$          | Obtains $v = k$ with $k$ correct for their<br>$ w - pi  =  w - qi $ oe<br>or substitutes $u + iv$ for $w$ , applies Pythagoras correctly and obtains a value for $v$<br>Allow “ $y = \dots$ ” for this mark.<br>Condone $v = 4i$ for this mark.<br><b>Requires both previous M marks.</b> | ddM1  |
|                                                                    | $v = 4$ only                                                                                                                                                        | Correct equation                                                                                                                                                                                                                                                                          | A1    |
|                                                                    | <b>(5)</b>                                                                                                                                                          |                                                                                                                                                                                                                                                                                           |       |
| <b>Total 6</b>                                                     |                                                                                                                                                                     |                                                                                                                                                                                                                                                                                           |       |
| <b>Alt 1:<br/>using<br/>points<br/>on<br/><math> z  = 1</math></b> | $z = 1 \Rightarrow w = \frac{9i - i}{1 + 1}, \quad z = i \Rightarrow w = \frac{9i^2 - i}{i + 1}$                                                                    | M1: Uses two correct values of $z$ in the formula to obtain two expressions for $w$<br>A1: Both expressions for $w$ correct                                                                                                                                                               | M1 A1 |
|                                                                    | $w = 4i, \quad w = \frac{-i - 9}{1 + i} \times \frac{1 - i}{1 - i} = \frac{-10 + 8i}{2} = -5 + 4i$                                                                  | Simplifies expressions to $a + bi$ form including correct method to rationalise denominators.<br><b>Requires previous M mark.</b>                                                                                                                                                         | dM1   |
|                                                                    | $\Rightarrow v = "4"$                                                                                                                                               | Obtains $v = b$ with $b$ correct for their $a + bi$ forms.<br>Allow “ $y = \dots$ ” for this mark.<br>Condone $v = 4i$ for this mark.<br><b>Requires both previous M marks.</b>                                                                                                           | ddM1  |
|                                                                    | $v = 4$ only                                                                                                                                                        | Correct equation                                                                                                                                                                                                                                                                          | A1    |
| <b>(5)</b>                                                         |                                                                                                                                                                     |                                                                                                                                                                                                                                                                                           |       |
| <b>Alt 2:<br/>using<br/><math>u + iv</math></b>                    | $w = \frac{9iz - i}{z + 1}$ $\Rightarrow wz + w = 9iz - i \Rightarrow z(w - 9i) = -w - i$                                                                           | M1: Makes $z$ the subject<br>A1: Correct expression oe                                                                                                                                                                                                                                    | M1 A1 |

|                                               |                                                                                                                                                                                                                  |                                                                                                        |              |
|-----------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------|--------------|
|                                               | $\Rightarrow z = \frac{-w-i}{w-9i} = \frac{w+i}{9i-w} \text{ oe}$                                                                                                                                                |                                                                                                        |              |
|                                               | $\Rightarrow z = \frac{-u-iv-i}{u+iv-9i} \times \frac{u-(v-9)i}{u-(v-9)i}$ $= \frac{-u^2-(v+1)(v-9)-10ui}{u^2+(v-9)^2}$ $\Rightarrow x = -\frac{u^2+(v+1)(v-9)}{u^2+(v-9)^2} \quad y = -\frac{10u}{u^2+(v-9)^2}$ | Replaces $w$ with $u + vi$ and rationalises the denominator and extracts real and imaginary parts      | <b>dM1</b>   |
|                                               | $x^2 + y^2 = 1 \Rightarrow \frac{(u^2+(v+1)(v-9))^2 + (10u)^2}{(u^2+(v-9)^2)^2} = 1$ $\Rightarrow \dots \Rightarrow (v-4)(u^2+v^2-18v+81) = 0$                                                                   | Applies $x^2 + y^2 = 1$ and simplifies to identify a factor $v-k=0$<br>Same conditions as main scheme. | <b>ddM1</b>  |
|                                               | $v = 4 \text{ only}$                                                                                                                                                                                             | Correct equation                                                                                       | <b>A1</b>    |
| <b>(5)</b>                                    |                                                                                                                                                                                                                  |                                                                                                        |              |
| <b>Alt 3:<br/>using<br/><math>x+iy</math></b> | $w = \frac{9i(x+iy)-i}{x+iy+1} \times \frac{x+1-iy}{x+1-iy} = \dots$ $= \frac{-9y(x+1) + (9x-1)y + ((9x-1)(x+1) + 9y^2)i}{(x+1)^2 + y^2}$                                                                        | M1: Replaces $z$ by $x + iy$ and rationalises the denominator.<br>A1: Correct expression oe            | <b>M1 A1</b> |
|                                               | $= \frac{-10y + (9x^2 + 8x - 1 + 9y^2)i}{1 + 2x + x^2 + y^2}$ $= \frac{-10y + (8x - 1 + 9)i}{1 + 2x + 1} \left[ = \frac{-5y + 4(x+1)i}{x+1} \right]$                                                             | Expands and applies $x^2 + y^2 = 1$ to simplify to linear coefficients                                 | <b>dM1</b>   |
|                                               | $= \frac{-5y}{x+1} + 4i \Rightarrow v = 4$                                                                                                                                                                       | Simplifies to a constant imaginary term and deduces the equation.<br>Same conditions as main scheme.   | <b>ddM1</b>  |
|                                               | $v = 4 \text{ only}$                                                                                                                                                                                             | Correct equation                                                                                       | <b>A1</b>    |
|                                               | <b>(5)</b>                                                                                                                                                                                                       |                                                                                                        |              |

## Additional Notes

(a)

B1: Draws the correct circle, allow tolerance with shape, look for the intent, and indicates the radius is 1. Accept  $\pm 1$  on the imaginary axis or incorrect notation of coordinates as long as the radius of 1 is implied.

May be drawn as the radius. Accept  $r=1$  stated next to the diagram, but not  $x^2 + y^2 = 1$  without further indication of the radius being 1.

(b)

Note for the main scheme ddM1 first deduction is from using the perpendicular bisector of  $9i$  and  $-i$ .

| Question Number | Scheme                                                                                                                                                                               | Notes                                                                                                                                                                                     | Marks       |
|-----------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>4</b>        | $4\frac{d^2y}{dx^2} - 11\frac{dy}{dx} - 3y = 78e^{3x} \quad y = \frac{9}{2}, \frac{dy}{dx} = 0 \text{ at } x = 0$                                                                    |                                                                                                                                                                                           |             |
| <b>(a)</b>      | $y = \lambda xe^{3x} \Rightarrow$<br>$y' = \lambda e^{3x} + 3\lambda xe^{3x}, \quad y'' = 3\lambda e^{3x} + 3\lambda e^{3x} + 9\lambda xe^{3x}$                                      | Correct first and second derivatives (need not be simplified).                                                                                                                            | B1          |
|                 | $4(6\lambda e^{3x} + 9\lambda xe^{3x}) - 11(\lambda e^{3x} + 3\lambda xe^{3x}) - 3(\lambda xe^{3x}) = 78e^{3x}$<br>$\Rightarrow 13\lambda e^{3x} = 78e^{3x} \Rightarrow \lambda = 6$ | M1: Substitutes their attempts at the derivatives of form $\alpha e^{3x} + \beta xe^{3x}$ into the differential equation and solves for $\lambda$<br>A1: Correct value from fully correct | M1<br>A1cso |
| <b>(3)</b>      |                                                                                                                                                                                      |                                                                                                                                                                                           |             |
| <b>(b)</b>      | $4m^2 - 11m - 3 = 0 \Rightarrow (4m+1)(m-3) = 0$<br>$\Rightarrow m = -\frac{1}{4}, 3$                                                                                                | Forms auxiliary equation and solves (usual rules)                                                                                                                                         | M1          |
|                 | CF: $y = Ae^{-\frac{1}{4}x} + Be^{3x}$                                                                                                                                               | For $Ae^{m_1x} + Be^{m_2x}$ with their real and different values of $m$                                                                                                                   | M1          |
|                 | $y = Ae^{-\frac{1}{4}x} + Be^{3x} + "6"xe^{3x}$                                                                                                                                      | Correct GS ft their non-zero $\lambda$ including "y = ..."                                                                                                                                | A1ft        |
| <b>(3)</b>      |                                                                                                                                                                                      |                                                                                                                                                                                           |             |
| <b>(c)</b>      | $\frac{dy}{dx} = -\frac{1}{4}Ae^{-\frac{1}{4}x} + 3Be^{3x} + 6e^{3x} + 18xe^{3x}$ or ft                                                                                              | M1: Differentiates to obtain an expression of the correct form<br>A1: Correct ft expression                                                                                               | M1 A1ft     |
|                 | $x = 0, y = \frac{9}{2} \Rightarrow \frac{9}{2} = A + B$<br>$x = 0, \frac{dy}{dx} = 0 \Rightarrow 0 = -\frac{1}{4}A + 3B + 6$<br>$\Rightarrow A = \dots, B = \dots$                  | Substitutes the given values into their equations and solves simultaneously for A and B                                                                                                   | M1          |
|                 | $y = 6e^{-\frac{1}{4}x} - \frac{3}{2}e^{3x} + 6xe^{3x}$                                                                                                                              | Fully correct particular solution (including "y = ..." unless this was penalised in part (b))                                                                                             | A1          |
| <b>(4)</b>      |                                                                                                                                                                                      |                                                                                                                                                                                           |             |
| <b>Total 10</b> |                                                                                                                                                                                      |                                                                                                                                                                                           |             |

## Additional Notes

You may mark this question as a whole. Do not be concerned about part labelling as many will answer out of order, e.g. finding CF before PI.

**(a) Note: This appears on ePEN as MMA but we are marking it as BMA**

B1: Correct first and second derivatives.

M1: Full process to find the constant. The derivatives need not have been correct for this mark but must have form  $\alpha e^{3x} + \beta xe^{3x}$  ( $\alpha, \beta \neq 0$ ), look for the correct process.

A1cso: Correct value from fully correct work. The  $xe^{3x}$  must have cancelled in their equation giving only  $\lambda = 6$ .

**(b)**

M1: Correct attempt to solve the auxiliary equation. If no method is shown the correct values can imply the attempt.

M1: Correct CF for their values, which must be real and distinct.

A1: Correct solution following through their value for  $\lambda$ . Must include the y =..

(c)

M1: Differentiates to correct form  $Pe^{\frac{x}{4}} + Qe^{3x} + Rxe^{3x} + "6"e^{3x}$  ( $P, Q, R \neq 0$ ), condoning at most one slip in powers. Must involve the product rule.

A1ft: Correct expression following through on their value for "6".

M1: For a full process to set up and solve simultaneous equations in  $A$  and  $B$  (or their variables) using the given values. Do not be concerned about algebraic slips, as long as two equations are set up then allow for any values following them.

A1: Fully correct, but only penalise a missing  $y = \dots$  in the first instance it would otherwise gain a mark.

Allow from solutions which gained  $\lambda = 6$  in (a) even if from incorrect work.

| Question Number | Scheme                                                                                                                                                                                                                                                                                                                                   | Notes                                                                                                                  | Marks |
|-----------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|-------|
| <b>5(a)</b>     | $\frac{2}{r(r+1)(r+2)} = \frac{A}{r} + \frac{B}{r+1} + \frac{C}{r+2}$ $2 = A(r+1)(r+2) + Br(r+2) + Cr(r+1)$ $r = 0: 2 = 2A \Rightarrow A = 1$ $r = -1: 2 = -B \Rightarrow B = -2$ $r = -2: 2 = 2C \Rightarrow C = 1$                                                                                                                     | Any valid method to obtain at least one constant with partial fractions of the correct form.<br>Could be implied       | M1    |
|                 | $\frac{1}{r} - \frac{2}{r+1} + \frac{1}{r+2}$                                                                                                                                                                                                                                                                                            | Correct partial fractions                                                                                              | A1    |
| <b>(2)</b>      |                                                                                                                                                                                                                                                                                                                                          |                                                                                                                        |       |
| <b>(b)</b>      | $\sum_{r=1}^n \frac{2}{r(r+1)(r+2)} = \left\{ \begin{array}{l} \frac{1}{1} - \frac{2}{2} + \frac{1}{3} + \\ \frac{1}{2} - \frac{2}{3} + \frac{1}{4} + \\ \frac{1}{3} - \frac{2}{4} + \frac{1}{5} + \\ \dots + \dots \\ \frac{1}{n-1} - \frac{2}{n} + \frac{1}{n+1} + \\ \frac{1}{n} - \frac{2}{n+1} + \frac{1}{n+2} \end{array} \right.$ | Attempts at least three rows with their partial fractions of a correct form (see notes). Implied by correct expression | M1    |
|                 | $= 1 - 1 + \frac{1}{2} + \frac{1}{n+1} - \frac{2}{n+1} + \frac{1}{n+2}$                                                                                                                                                                                                                                                                  | A1: Correct constant terms or algebraic terms extracted<br>A1: Fully correct expression                                | A1 A1 |
|                 | $\frac{1}{2} - \frac{1}{n+1} + \frac{1}{n+2} = \frac{(n+1)(n+2) - 2(n+2) + 2(n+1)}{2(n+1)(n+2)}$                                                                                                                                                                                                                                         | Combines terms of the form $p$ , $\frac{q}{n+b}$ , and $\frac{r}{n+c}$ and no others into a single fraction            | M1    |
|                 | $= \frac{n^2 + 3n + 2 - 2n - 4 + 2n + 2}{2(n+1)(n+2)} = \frac{n(n+3)}{2(n+1)(n+2)}$                                                                                                                                                                                                                                                      | Correct expression                                                                                                     | A1    |
| <b>(5)</b>      |                                                                                                                                                                                                                                                                                                                                          |                                                                                                                        |       |
| <b>(c)</b>      | $\frac{n(n+3)}{2(n+1)(n+2)} > \frac{7}{15}$ $15n^2 + 45n > 14n^2 + 42n + 28$                                                                                                                                                                                                                                                             | Forms inequality or equation, cross-multiplies or puts over common denominator and expands brackets                    | M1    |
|                 | $n^2 + 3n - 28 > 0$ $(n+7)(n-4) > 0$ $\Rightarrow n = \dots \text{ or } n > \dots \quad (n > 4)$                                                                                                                                                                                                                                         | Forms and solves 3TQ inequality or equation (usual rules)                                                              | dM1   |
|                 | $n = 5$                                                                                                                                                                                                                                                                                                                                  | 5 only                                                                                                                 | A1    |
| <b>(3)</b>      |                                                                                                                                                                                                                                                                                                                                          |                                                                                                                        |       |
| <b>Total 10</b> |                                                                                                                                                                                                                                                                                                                                          |                                                                                                                        |       |

Additional Note

(a)

M1: Must be correct partial fraction form with three fractions. Note that  $\frac{A}{r(r+1)} + \frac{B}{(r+1)(r+2)}$  is not a

complete partial fraction form so is M0, but this form can be used to score full marks in part (b).

A1: Correct form.

(b)

M1: For evidence of correctly applying the method of differences to the question. This may be implied by extraction of the correct terms if no/incomplete method is seen. If method is shown they should give at least three lines of fractions to show cancelling.

A1: For either the correct constant terms at the start or the correct algebraic terms at the end extracted from the differences.

A1: All correct terms extracted (and not others)

Note that  $\frac{1}{r(r+1)} - \frac{1}{(r+1)(r+2)}$  will give  $\frac{1}{2} - \frac{1}{6} + \frac{1}{6} - \frac{1}{12} + \dots - \frac{1}{(n+1)(n+2)} = \frac{1}{2} - \frac{1}{(n+1)(n+2)}$  which can

score M1A1A1, and then the next M for combining these.

M1: Combines the terms to a single fraction provided terms of suitable forms were extracted – as shown in scheme. The denominator must be correct but allow a slip in the numerator.

A1: Correct answer from correct work.

(c)

M1: Sets up and forms the required inequality, or accept as an equality formed, and proceeds to form an unsimplified quadratic equation or inequality by cross multiplying or by **correctly** putting terms over a common denominator and expanding the  $n(n+3)$  and  $(n+1)(n+2)$  brackets.

dM1: Solves the resulting 3 term quadratic (accept if just one root found). Note: answers which try to solve for the whole solution set for  $n$  will carry this out as part of the working, and can score full marks. Solutions to the quadratic by calculator are acceptable – you may need to check and accept if only one root of their quadratic is given.

A1: Correct answer only.

Note: Trial and Improvement or calculator methods which do not form a quadratic inequality or equation will score no marks.

| Question Number | Scheme                                                                                                                                                                                                                                                                                                        | Notes                                                                                                                                                                                                                          | Marks         |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| <b>6</b>        | $2 \frac{dy}{dx} + (\cot x)y + (\tan x \sec x)y^3 = 0$                                                                                                                                                                                                                                                        |                                                                                                                                                                                                                                |               |
| <b>(a)</b>      | $z = \frac{1}{y^2} \Rightarrow \frac{dz}{dx} = -\frac{2}{y^3} \frac{dy}{dx}$ or<br>$y^2 = \frac{1}{z} \Rightarrow 2y \frac{dy}{dx} = -\frac{1}{z^2} \frac{dz}{dx} \left\{ = -y^4 \frac{dz}{dx} \right\}$ or<br>$y = z^{-\frac{1}{2}} \Rightarrow \frac{dy}{dx} = -\frac{1}{2} z^{-\frac{3}{2}} \frac{dz}{dx}$ | Any correct equation in $\frac{dy}{dx}$ and $\frac{dz}{dx}$ (may be implied)                                                                                                                                                   | B1            |
|                 | $-y^3 \frac{dz}{dx} + (\cot x)y + (\tan x \sec x)y^3 = 0$                                                                                                                                                                                                                                                     | Substitutes to obtain an equation with $\frac{dz}{dx}$ as the only derivative                                                                                                                                                  | M1            |
|                 | $-\frac{dz}{dx} + \frac{\cot x}{y^2} + \tan x \sec x = 0$<br>$\frac{dz}{dx} - (\cot x)z = \tan x \sec x^*$                                                                                                                                                                                                    | Fully correct proof with intermediate step after substitution                                                                                                                                                                  | A1*           |
| <b>(3)</b>      |                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                                |               |
| <b>(b)</b>      | I.F. = $e^{\int -\cot x dx}$                                                                                                                                                                                                                                                                                  | I.F. = $e^{\int \pm \cot x dx}$                                                                                                                                                                                                | M1            |
|                 | $= e^{-\ln \sin x} = \frac{1}{\sin x}$                                                                                                                                                                                                                                                                        | Correct processed I.F. in any form                                                                                                                                                                                             | A1            |
|                 | $\frac{z}{\sin x} = \int \frac{1}{\sin x} \times \frac{\sin x}{\cos x} \times \frac{1}{\cos x} dx$                                                                                                                                                                                                            | Applies<br>I.F. $\times z = \int$ I.F. $\times \tan x \sec x dx$<br>Condone y for z                                                                                                                                            | M1            |
|                 | $\frac{z}{\sin x} = \int \sec^2 x dx \Rightarrow \frac{z}{\sin x} = \tan x (+c)$                                                                                                                                                                                                                              | Correct equation in z and x<br>Condone missing "+c"                                                                                                                                                                            | A1            |
|                 | $z = \sin x (\tan x + c) \Rightarrow y^2 = \frac{1}{\sin x (\tan x + c)}$ oe                                                                                                                                                                                                                                  | Fully correct general solution. Accept equivalent forms.                                                                                                                                                                       | A1            |
| <b>(5)</b>      |                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                                |               |
| <b>(c)</b>      | $\frac{4\sqrt{3}}{3} = \frac{1}{\frac{1}{2}(\frac{\sqrt{3}}{3} + c)} \Rightarrow \frac{\sqrt{3}}{3} + c = \frac{\sqrt{3}}{2} \Rightarrow c = \dots \left( \frac{\sqrt{3}}{6} \right)$                                                                                                                         | Correctly substitutes the given values to find a value for c                                                                                                                                                                   | M1            |
|                 | $x = \frac{\pi}{3} \Rightarrow y^2 = \frac{1}{\frac{\sqrt{3}}{2}(\sqrt{3} + \frac{\sqrt{3}}{6})} = \frac{4}{7}$<br>$\Rightarrow y = \pm \frac{2\sqrt{7}}{7}$ oe                                                                                                                                               | <b>dM1:</b> Substitutes $x = \frac{\pi}{3}$ and their c into their equation and reaches $y^2 = \dots$ or $y =$ with trig terms evaluated.<br><b>A1:</b> Both correct exact values.<br>The negative value must not be rejected. | <b>dM1 A1</b> |
| <b>(3)</b>      |                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                                |               |
| <b>Total 11</b> |                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                                |               |

Additional Notes:

Condone poor notation of  $\sin^{-1} x$  for cosec x as long as it is treated correctly.

(a) Note for A mark you may condone a missing "=0" in intermediate lines as long as it is implied by working.

(b) Note an equivalent form for the answer is  $y^2 = \frac{\cos x}{\sin^2 x + K \sin 2x}$  which will lead to  $c = \frac{\sqrt{3}}{12}$  in (c)

| Question Number | Scheme                                                                                                                                                                                                                                                                                                                                                          | Notes                                                                                                                                                                                                             | Marks |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| 7               | $r = \sqrt{3} \sin \theta \quad (0 \leq \theta \leq \pi) \quad r = 3 \cos \theta \quad \left(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}\right)$                                                                                                                                                                                                              |                                                                                                                                                                                                                   |       |
| (a)             | $\sqrt{3} \sin \theta = 3 \cos \theta \Rightarrow \tan \theta = \sqrt{3}$<br>$\Rightarrow \theta = \frac{\pi}{3}, \quad r = \frac{3}{2} \quad \text{or} \quad \left(\frac{3}{2}, \frac{\pi}{3}\right)$                                                                                                                                                          | B1: Either correct coordinate<br>B1: Both correct coordinates and no others                                                                                                                                       | B1 B1 |
| <b>(2)</b>      |                                                                                                                                                                                                                                                                                                                                                                 |                                                                                                                                                                                                                   |       |
| (b)             | $\int (\sqrt{3} \sin \theta)^2 d\theta = K \int \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta d\theta = \dots$ or<br>$\int (3 \cos \theta)^2 d\theta = M \int \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta d\theta = \dots$                                                                                                                                      | Uses one of<br>$\sin^2 \theta = \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta$ or<br>$\cos^2 \theta = \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta$ in an attempt to integrate $r^2$ for one of the equations.     | M1    |
|                 | $\int (\sqrt{3} \sin \theta)^2 d\theta = K \int \frac{1}{2} - \frac{1}{2} \cos 2\theta d\theta = \dots$ and<br>$\int (3 \cos \theta)^2 d\theta = M \int \frac{1}{2} + \frac{1}{2} \cos 2\theta d\theta = \dots$                                                                                                                                                 | Uses both<br>$\sin^2 \theta = \frac{1}{2} - \frac{1}{2} \cos 2\theta$ and<br>$\cos^2 \theta = \frac{1}{2} + \frac{1}{2} \cos 2\theta$ in an attempt to integrate $r^2$ for both of the equations to correct form. | M1    |
|                 | $\int (\sqrt{3} \sin \theta)^2 d\theta = \frac{3}{2} \left( \theta - \frac{1}{2} \sin 2\theta \right)$ and<br>$\int (3 \cos \theta)^2 d\theta = \frac{9}{2} \left( \theta + \frac{1}{2} \sin 2\theta \right)$                                                                                                                                                   | Fully correct expression after integration ignoring limits (accept equivalent expressions)                                                                                                                        | A1    |
|                 | $\frac{1}{2} \int_0^{\frac{\pi}{3}} (\sqrt{3} \sin \theta)^2 d\theta = \frac{3}{4} \left[ \theta - \frac{1}{2} \sin 2\theta \right]_0^{\frac{\pi}{3}} = \dots$ or<br>$\frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} (3 \cos \theta)^2 d\theta = \frac{9}{4} \left[ \theta + \frac{1}{2} \sin 2\theta \right]_{\frac{\pi}{3}}^{\frac{\pi}{2}} = \dots$        | Full process for at least one of the two areas required, including correct area formula, with their attempt to apply the double angle formula, and evidence of correct limits applied.                            | M1    |
|                 | Area of $R = \frac{1}{2} \int_0^{\frac{\pi}{3}} 3 \sin^2 \theta d\theta + \frac{1}{2} \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} 9 \cos^2 \theta d\theta$<br>$= \frac{3}{4} \left[ \frac{\pi}{3} - \frac{1}{2} \sin \frac{2\pi}{3} - (0 - 0) \right] + \frac{9}{4} \left[ \frac{\pi}{2} + 0 - \left( \frac{\pi}{3} + \frac{1}{2} \sin \frac{2\pi}{3} \right) \right]$ | Full and correct process for the area of $R$ . Formulae and limits must all be correct, and evidence of limits applied, with areas added.                                                                         | DM1   |
|                 | $\frac{\pi}{4} - \frac{3\sqrt{3}}{16} + \frac{3\pi}{8} - \frac{9\sqrt{3}}{16} = \frac{5\pi}{8} - \frac{3\sqrt{3}}{4}$                                                                                                                                                                                                                                           | Correct answer.<br>Allow $a = \frac{5}{8}, \quad b = -\frac{3}{4}$                                                                                                                                                | A1    |
| <b>(6)</b>      |                                                                                                                                                                                                                                                                                                                                                                 |                                                                                                                                                                                                                   |       |
| <b>Total 8</b>  |                                                                                                                                                                                                                                                                                                                                                                 |                                                                                                                                                                                                                   |       |

## Additional Notes

(a)

B1: One correct coordinate, do not be concerned about the method for this mark. Allow for  $\theta = \frac{\pi}{3}$  seen

and isw if they go on to use  $\theta = \frac{\pi}{3}$  for this mark.

B1: Both coordinates correct, and no extras. Again, do not be concerned about the method. Condone if given in the wrong order in a coordinate pair.

(b)

M1: Starts correct process to find one of the two areas. Must be using  $r^2$  for one of the two curves, attempt a double angle formula  $\sin^2 \theta = \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta$  or  $\cos^2 \theta = \pm \frac{1}{2} \pm \frac{1}{2} \cos 2\theta$  and any attempt to integrate the result (no matter how poor, ie a changed function). Condone errors with the coefficient not being squared.

M1: A full process to find the integral of  $r^2$  for **both** curves with **correct** double angle formulae used, so  $\sin^2 \theta = \frac{1}{2} - \frac{1}{2} \cos 2\theta$  and  $\cos^2 \theta = \frac{1}{2} + \frac{1}{2} \cos 2\theta$  and reaches correct for each, but again allow if slips occur in the coefficient.

A1: Fully correct integration for  $r^2$  for **both** curves. The coefficients must have been correctly dealt with.

Note the  $\frac{1}{2}$  may be included throughout.

M1: For a full correct process to find at least one of the areas. Must be using  $\frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$  including the  $\frac{1}{2}$

and with **correct** limits for the area applied (condone not showing subtracted 0's), but condone slips in the coefficients when squaring and condone if incorrect double angle formula attempts were made.

Note: Some candidates may attempt to find only one area and double the result. If the area formula is clearly seen or applied before doubling this is acceptable as an attempt for the full process for one area.

DM1: Depends on all previous method marks. Fully correct process for **both** areas and adds the results. This may be done separately or in one expression throughout the working.

A1: Correct answer.

| Question Number | Scheme                                                                                                                                            | Notes                                                                                                                                               | Marks |
|-----------------|---------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|-------|
| <b>8(a)</b>     | For intersection when $x > 0$ :<br>$\frac{15x}{x+4} = x - 2$ $15x = x^2 + 2x - 8$ $x^2 - 13x - 8 = 0$ $\Rightarrow x = \frac{13 + \sqrt{201}}{2}$ | M1: Solves 3TQ obtained from a correct equation (usual rules) for $x > 0$<br>A1: Correct exact positive value (condone negative root also included) | M1 A1 |
|                 | For intersections when $x < 0$ :<br>$\frac{15x}{-x+4} = x - 2$ $15x = -x^2 + 6x - 8$ $x^2 + 9x + 8 = 0$ $(x+8)(x+1) = 0$ $\Rightarrow x = -8, -1$ | M1: Solves 3TQ obtained from a correct equation (usual rules) for $x < 0$<br>A1: Correct values                                                     | M1 A1 |
|                 | $-8 < x < -1, \quad x > \frac{13 + \sqrt{201}}{2}$                                                                                                | M1: For $a < x < b, x > c$ oe with cvs $a < b < 0 < c$ from the correct equations.<br>A1: Fully correct set of values in any form                   | M1 A1 |
| <b>(6)</b>      |                                                                                                                                                   |                                                                                                                                                     |       |
| <b>(b)</b>      | $\frac{15x}{x+4} = -x + 2$ $15x = -x^2 - 2x + 8$ $x^2 + 17x - 8 = 0$ $\Rightarrow x = \frac{-17 + \sqrt{321}}{2}$                                 | M1: Solves 3TQ obtained from a correct equation (usual rules)<br>A1: Correct exact positive value (condone negative root also included)             | M1 A1 |
|                 | $x < "-8", \quad x > "\frac{13 + \sqrt{201}}{2}"$                                                                                                 | Correct regions ft their $a < 0$ and $c > 0$                                                                                                        | B1ft  |
|                 | $x < "-8", \quad "-1" < x < \frac{-17 + \sqrt{321}}{2}, \quad x > "\frac{13 + \sqrt{201}}{2}"$                                                    | Correct set of values in any form.<br>Allow ft on their cvs $a, b$ and $c$ provided $a < b < 0$ and $c > \frac{-17 + \sqrt{321}}{2} (= 0.458...)$   | A1ft  |
| <b>(4)</b>      |                                                                                                                                                   |                                                                                                                                                     |       |
| <b>Total 10</b> |                                                                                                                                                   |                                                                                                                                                     |       |

Additional notes.

“Usual Rules” : as well as the general information, we will accept here solutions by calculator having reached a quadratic equation but they must be correct solutions for the stated quadratic if no method is shown (you may need to check).

(a)

Accept with any inequalities used instead of equalities for the solving of the intersection points. For methods via putting over common denominators score the Ms for the process of putting over a common denominator and solving for the numerator = 0.

May see attempts via squaring

---

$\frac{15x}{|x|+4} = x-2 \Rightarrow |x| = \frac{15x}{x-2} - 4 = \frac{11x+8}{x-2} \Rightarrow x^2 = \left(\frac{11x+8}{x-2}\right)^2$  then cross multiplies (or puts over a common denominator  $(x-2)^2$ ) to get

$\Rightarrow x^2(x^2 - 4x + 4) - (121x^2 + 176x + 64) = x^4 - 4x^3 - 117x^2 - 176x - 64 = 0$ . This will probably then be solved by calculator, which as algebra has been used will be accepted. Score M1 for the process reaching quartic, A1 correct quartic, M1 for solving the quartic (look for attempt to factorise to linear terms with correct product of roots, or accept answer from calculator – may need to check), A1 for at least the three relevant roots (and condone if  $x=2$  is also include for this mark).

For the final M they must be producing a solution set of the correct form from their CVs from attempts at the correct equations (but may have had incorrect method for solving), so if extra values from denominators are used it will be M0 (regardless of whether the form of the answer is correct).

FYI  $\frac{-13-\sqrt{201}}{2} \approx -13.6$  is being used often as one of the limits, but note it is less than  $-8$ , so

$-8 < x < \frac{-13-\sqrt{201}}{2}$  used is M0 but you may condone the " $-8 < x$ " for the B1ft in (b).

Final A1: Accept any correct equivalent notation. Accept with "and" or "or" written between inequalities,

but if using set notation must use the union. E.g. in interval notation  $(-8, -1) \cup \left(\frac{13+\sqrt{201}}{2}, \infty\right)$ . Use of

square brackets is A0.

(b)

Attempts at squaring may be seen again here. They will need a full process leading solving a quartic. Again for the A, accept if other roots are also included at this stage.

B1ft: For the follow through same "outside" ends of the region matching their answer to (b) in their **final** answer.

A1ft: Accept answers of the correct form providing their answer to (a) was of the correct form as specified

in the scheme and their  $c$  is  $c > \frac{-17+\sqrt{321}}{2} (= 0.458...)$ .

| Question Number | Scheme                                                                                                                                                                                                                                                                      | Notes                                                                                                                                                                                                   | Marks  |
|-----------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------|
| <b>9(a)</b>     | $\cos 5\theta + i \sin 5\theta = (\cos \theta + i \sin \theta)^5$ $= \cos^5 \theta + {}^5C_1 \cos^4 \theta (i \sin \theta) + {}^5C_2 \cos^3 \theta (i \sin \theta)^2 + {}^5C_3 \cos^2 \theta (i \sin \theta)^3 + {}^5C_4 \cos \theta (i \sin \theta)^4 + (i \sin \theta)^5$ |                                                                                                                                                                                                         | M1     |
|                 | $(\sin 5\theta =) 5 \cos^4 \theta \sin \theta - 10 \cos^2 \theta \sin^3 \theta + \sin^5 \theta$                                                                                                                                                                             | M1: Equates imaginary parts<br>A1: Correct expression                                                                                                                                                   | M1 A1  |
|                 | $= 5 \sin \theta (1 - \sin^2 \theta)^2 - 10 \sin^3 \theta (1 - \sin^2 \theta) + \sin^5 \theta$ $= 5 \sin \theta - 10 \sin^3 \theta + 5 \sin^5 \theta - 10 \sin^3 \theta + 10 \sin^5 \theta + \sin^5 \theta$ $= 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta$         | M1: Uses $\cos^2 \theta = 1 - \sin^2 \theta$ and reaches an expression of the correct form<br>A1: Correct expression or<br>$a = 16, b = -20, c = 5$<br>$\sin 5\theta$ must have been seen at some point | M1 A1  |
| <b>(5)</b>      |                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                         |        |
| <b>(b)</b>      | $\left( \text{For } \theta = \frac{k\pi}{5}, k \in \mathbb{Z}, \sin 5\theta = \sin k\pi = 0 \Rightarrow \right)$ $16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta = 0$                                                                                                  | Sets their answer to (a) = 0<br>Could be implied                                                                                                                                                        | M1     |
|                 | $\sin \theta (16 \sin^4 \theta - 20 \sin^2 \theta + 5) = 0$ $\Rightarrow \sin^2 \theta = \frac{-(-20) \pm \sqrt{(-20)^2 - 4(16)(5)}}{2(16)} = \frac{20 \pm \sqrt{80}}{32}$                                                                                                  | Solves their quadratic in $\sin^2 \theta$ to find both roots (usual rules)                                                                                                                              | M1     |
|                 | $\Rightarrow \sin^2 \frac{k\pi}{5} = \frac{5 \pm \sqrt{5}}{8}, 0$                                                                                                                                                                                                           | A1: $\frac{5 \pm \sqrt{5}}{8}$ oe<br>B1: 0                                                                                                                                                              | A1, B1 |
| <b>(4)</b>      |                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                         |        |
| <b>Total 9</b>  |                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                         |        |

## Additional Notes

(a)

M1: Attempts to expand  $(\cos \theta + i \sin \theta)^5$  to obtain an expression of the correct form for the three imaginary terms. May use  $c + is$  notation. Need only see the imaginary terms, condone missing or incorrect real terms. Binomial coefficients may be as expressions or values, if values only used they must be correct. The powers of  $i$  may be processed in the initial expansion. Allow invisible brackets if recovered.

M1: Extracts the imaginary terms from the expansion. The equating to  $\sin 5\theta$  may be tacit at this stage.

A1: Correct expression for  $\sin 5\theta$  (though the  $\sin 5\theta$  need not be seen yet) in terms of  $\sin \theta$  and  $\cos \theta$ .

M1: Uses  $\cos^2 \theta = 1 - \sin^2 \theta$  to eliminate  $\cos \theta$  from the expression to get to an expanded equation in  $\sin \theta$  only. Terms need not be gathered at this stage, but the simplification may be done in one step.

A1: Correct answer including the  $\sin 5\theta$  having been seen identified with the expression at some stage if not included at the end. Accept values for  $a, b$  and  $c$  stated following reaching a suitable equation in  $\sin \theta$

(b)

M1: Recognition of the correct process seen by either their answer to (a) equal to zero, which by be implied, or by an attempt to solve it (the “=0” being implied, accept one correct root for their equation as evidence).

M1: Correct method to solve the quadratic in  $\sin^2 \theta$  after factoring out the  $\sin \theta$  - both roots must be given. Condone for this mark if they mistakenly mislabel the solution to the quadratic in  $\sin^2 \theta$  as  $\sin \theta$ , but do not

allow the A if they go on square their values. Note answer stated directly from the quintic score M0. They must at least identify the quadratic in  $\sin^2 \theta$  before reaching the answers (from calculator).

A1: Correct answers only but accept equivalent forms.

B1: For 0 identified as a value for  $\sin^2 \frac{k\pi}{5}$  - this can come from anywhere (method not required).

