

Write your name here

Surname	Other names
---------	-------------

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Further Pure
Mathematics F2
Advanced/Advanced Subsidiary

Sample Assessment Material
Time: 1 hour 30 minutes

Paper Reference
WFM02/01

You must have:
Mathematical Formulae and Statistical Tables (Blue)

Total Marks

--

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

S45002A

©2013 Pearson Education Ltd.



S 4 5 0 0 2 A 0 1 2 4

PEARSON

2. The displacement x metres of a particle at time t seconds is given by the differential equation

$$\frac{d^2x}{dt^2} + x + \cos x = 0$$

When $t = 0$, $x = 0$ and $\frac{dx}{dt} = \frac{1}{2}$.

Find a Taylor series solution for x in ascending powers of t , up to and including the term in t^3 .

(5)

5.




Figure 1

Figure 1 shows the curves given by the polar equations

$$r = 2, \quad 0 \leq \theta \leq \frac{\pi}{2},$$

$$\text{and } r = 1.5 + \sin 3\theta, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

- (a) Find the coordinates of the points where the curves intersect. (3)

The region S , between the curves, for which $r > 2$ and for which $r < (1.5 + \sin 3\theta)$, is shown shaded in Figure 1.

- (b) Find, by integration, the area of the shaded region S , giving your answer in the form $a\pi + b\sqrt{3}$, where a and b are simplified fractions. (7)

Leave
blank

6. A complex number z is represented by the point P in the Argand diagram.

(a) Given that $|z-6|=|z|$, sketch the locus of P . (2)

(b) Find the complex numbers z which satisfy both $|z-6|=|z|$ and $|z-3-4i|=5$. (3)

The transformation T from the z -plane to the w -plane is given by $w = \frac{30}{z}$.

(c) Show that T maps $|z-6|=|z|$ onto a circle in the w -plane and give the cartesian equation of this circle. (5)

Leave
blank

7. (a) Show that the transformation $z = y^{\frac{1}{2}}$ transforms the differential equation

$$\frac{dy}{dx} - 4y \tan x = 2y^{\frac{1}{2}} \quad (\text{I})$$

into the differential equation

$$\frac{dz}{dx} - 2z \tan x = 1 \quad (\text{II}) \quad (5)$$

- (b) Solve the differential equation (II) to find z as a function of x . (6)

- (c) Hence obtain the general solution of the differential equation (I). (1)
