



Mark Scheme (Results)

Summer 2014

Pearson Edexcel International A Level in
Further Pure Mathematics F2
(WFM02/01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2014

Publications Code IA038888

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
 7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Further Pure Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \dots$

2. Formula

Attempt to use the correct formula (with values for a, b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $\left(x \pm \frac{b}{2}\right)^2 \pm q \pm c = 0$, $q \neq 0$, leading to $x = \dots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small errors in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required.

| Question Number | Scheme | | Marks |
|-----------------|--|--|----------------|
| 1.(a) | $\frac{1}{2(r+1)(r+2)} - \frac{1}{2(r+2)(r+3)}$ $= \frac{r+3 - (r+1)}{2(r+1)(r+2)(r+3)}$ | Attempt common denominator | M1 |
| | $= \frac{2}{2(r+1)(r+2)(r+3)}$ $= \frac{1}{(r+1)(r+2)(r+3)}$ | Correct proof | A1 |
| | | | (2) |
| (a) Way 2 | $\frac{1}{(r+1)(r+2)(r+3)} = \frac{1}{r+2} \left(\frac{1}{(r+1)(r+3)} \right) = \frac{1}{r+2} \left(\frac{1}{2(r+1)} - \frac{1}{2(r+3)} \right)$ $\frac{1}{2(r+1)(r+2)} - \frac{1}{2(r+2)(r+3)}$ | | M1 A1 |
| | M1: Factor of $\frac{1}{r+2}$ and attempt partial fractions A1: Correct proof | | |
| | Other methods: Complete method scores All work correct inc final answer reached | M1 A1 | |
| (b) | $\sum_{r=1}^n \frac{1}{(r+1)(r+2)(r+3)} =$ $= \frac{1}{12} - \frac{1}{24} + \dots$ $+ \frac{1}{2(n+1)(n+2)} - \frac{1}{2(n+2)(n+3)}$ | Attempt at least the first pair and the last pair of terms as shown. Must start at 1 and end at n | M1 |
| | $= \frac{1}{12} - \frac{1}{2(n+2)(n+3)}$ | Identifies that the first and last terms do not cancel. | M1 |
| | $= \frac{n^2 + 5n + 6 - 6}{12(n+2)(n+3)}$ | Correctly combined fractions | A1 |
| | $= \frac{n^2 + 5n}{12(n+2)(n+3)} \text{ or } \frac{n(n+5)}{12(n+2)(n+3)} *$ | Allow either form isw attempts to multiply out the denominator | A1 |
| | | | (4) |
| | | | Total 6 |

| Question Number | Scheme | | Marks |
|-----------------|--|---|--------|
| 2. | $\frac{6}{x-3} \leq x+2$ | | |
| | Way 1 | | |
| | $\frac{6}{x-3} \leq x+2 \Rightarrow x+2 - \frac{6}{x-3} \geq 0$ | | |
| | $x+2 - \frac{6}{x-3} \geq 0 \Rightarrow \frac{(x+3)(x-4)}{x-3} \geq 0$ | Attempt to combine fractions and factorise the numerator | M1 |
| | $x = -3, x = 4$ | Correct critical values | A1, A1 |
| | $x \geq 4$ | Follow through their 4 | A1ft |
| | $x = 3$ | Identifies 3 as a critical value | B1 |
| | $-3 \leq x < 3$ | M1: Attempt inside region | M1A1 |
| | | A1: Correct inequality | |
| | | | (7) |
| | Way 2 | | |
| | $6(x-3) \leq (x+2)(x-3)^2$ $\Rightarrow (x-3)(4-x)(x+3) \geq 0$ | Multiplies both sides by $(x-3)^2$ and attempt to factorise | M1 |
| | $x = -3, x = 4$ | Correct critical values | A1, A1 |
| | $x \geq 4$ | Follow through their 4 | A1ft |
| | $x = 3$ | Identifies 3 as a critical value | B1 |
| | $-3 \leq x < 3$ | M1: Attempt inside region | M1A1 |
| | | A1: Correct inequality | |
| | | | (7) |
| | Way 3 | | |
| | $x-3 > 0 \Rightarrow 6 \leq (x+2)(x-3)$ $\Rightarrow (x-4)(x+3) \geq 0$ | Multiplies both sides by $(x-3)$ and attempt to factorise Must state $x-3 > 0$ | M1 |
| | $x = 4$ | Correct critical values | A1 |
| | $x \geq 4$ | Follow through their 4 | A1ft |
| | $x = 3$ | Identifies 3 as a critical value | B1 |
| | $(x-3 < 0) \Rightarrow 6 \geq (x+2)(x-3)$ $x = -3$ | Correct critical value | A1 |
| | $(x+2)(x-3) \leq 6 \Rightarrow (x-4)(x+3) \leq 0$ $\Rightarrow -3 \leq x < 3$ | M1: Attempt inside region | M1A1 |
| | | A1: Correct inequality | |
| | | | (7) |

| Question Number | Scheme | | Marks |
|-----------------|--|---|----------------------------|
| 3. | | | |
| | $r^5 = \sqrt{16^2 + (16\sqrt{3})^2} = 32 \Rightarrow r = 32^{\frac{1}{5}} (= 2)$ | Correct value for r | B1 |
| | $\arg(16 - 16i\sqrt{3}) = \frac{5\pi}{3}$ | Allow $\frac{5\pi}{3}$ or $-\frac{\pi}{3}$ | B1 |
| | $5\theta = \frac{11\pi}{3}, \frac{17\pi}{3}, \frac{23\pi}{3}, \frac{29\pi}{3}$ | $\left(\frac{5\pi}{3}\right) + 2n\pi, n = 1, 2, 3, 4$ At least 2 values which must be positive. May be implied by correct final answers. | M1 |
| | $z = \underline{2e^{\frac{\pi i}{3}}}, \underline{2e^{\frac{11\pi i}{15}}}, \underline{2e^{\frac{17\pi i}{15}}}, \underline{2e^{\frac{23\pi i}{15}}}, \underline{2e^{\frac{29\pi i}{15}}}$ | 2 or $32^{\frac{1}{5}}, e^{\frac{5\pi i}{15}}$ or $e^{\frac{\pi i}{3}}$ | <u>B1 A1(all 4 values)</u> |
| | | | (5) |
| | | | Total 5 |

| Question Number | Scheme | Marks |
|-----------------|--|----------------|
| 4 | $w = \frac{z}{z+3}$ | |
| | $w = \frac{z}{z+3} \Rightarrow z = \frac{3w}{1-w}$ | M1A1 |
| | $ z = 2 \Rightarrow \left \frac{3w}{1-w} \right = 2$ $ 3w = 2 1-w $ $9(u^2 + v^2) = 4(u-1)^2 + 4v^2$ | M1A1 |
| | $5u^2 + 5v^2 + 8u - 4 = 0$ | |
| | $\left(u + \frac{4}{5}\right)^2 + v^2 - \frac{16}{25} - \frac{4}{5} = 0$ | M1 |
| | Centre $\left(-\frac{4}{5}, 0\right)$ | A1 |
| | Radius $\frac{6}{5}$ | A1 |
| | | (7) |
| | | Total 7 |

| Question Number | Scheme | | Marks |
|-----------------|---|--|-----------------|
| 5 | $\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} + 2y = 0$ | | |
| (a) | $y''' - 2y' - 2xy'' + 2y' (= 0) (y''' = 2xy'')$ | M1: Attempt to differentiate including use of the product rule on $2x\frac{dy}{dx}$ Equation may have been re-written as $y'' = \dots$ before differentiating | M1A1 |
| | | A1: Correct differentiation | |
| | $y'''' - 2y'' - 2xy''' - 2y'' + 2y' (= 0)$ | M1: Second use of product rule. Dependent on first M1. | dM1A1 |
| | | A1: Correct differentiation NB A simpler form is obtained if $y''' - 2xy'' = 0$ is used. | |
| | $y'''' = 2xy''' + 2y''$ $= 2x(2xy'') + 2y'' = (4x^2 + 2)y''$ | Cao and cso | A1 |
| | | | (5) |
| (b) | $y_0'' = -2, y_0''' = 0, y_0^{(4)} = -4$ | B1: $y_0'' = -2$ | B1, M1A1 |
| | | M1: Attempts y_0''' and $y_0^{(4)}$ | |
| | | A1: All correct and obtained from correct expressions | |
| | $(y =) 1 + 3x - x^2 - \frac{x^4}{6}$ | M1: Correct use of Maclaurin series | M1A1 |
| | | A1: Fully correct expansion. | |
| | | | (5) |
| (c) | $(y =) 1 + 3(-0.2) - (-0.2)^2 - \frac{(-0.2)^4}{6}$ | Use of the correct Maclaurin series and substitution of $x = -0.2$ | M1 |
| | | $(y =) 0.3597$ | Allow awrt |
| | | | (2) |
| | | | Total 12 |

| Question Number | Scheme | | Marks |
|-----------------|---|---|----------------|
| 6. | $x \frac{dy}{dx} + (1 - 2x)y = x$ | | |
| | $\frac{dy}{dx} + \frac{(1 - 2x)}{x}y (= 1)$ | Divides by x - may be implied by subsequent working | M1 |
| | Integrating factor $I = e^{\int \frac{1-2x}{x} dx}$ | Correct attempt at I , including an attempt at the integration. In must be seen if not fully correct. | dM1 |
| | $= e^{\ln(x)-2x}$ | Correct expression | A1 |
| | $= xe^{-2x}$ | No errors in working allowed | A1 |
| | $xye^{-2x} = \int xe^{-2x} dx$ | Multiply through by their IF and integrate LHS. Can be given if y x their IF is seen | M1 |
| | $= -\frac{1}{2}xe^{-2x} + \frac{1}{2}\int e^{-2x} dx$ | M1: Correct application by parts ie differentiate x and attempt to integrate e^{-2x} | M1A1 |
| | | A1: Correct expression | |
| | $= -\frac{1}{2}xe^{-2x} - \frac{1}{4}e^{-2x} (+c)$ | A1: Complete the integration to a correct result. Constant not required. | A1 |
| | $y = \frac{ce^{2x}}{x} - \frac{1}{4x} - \frac{1}{2}$ | Oe Must have $y = \dots$ Must include a constant. ft their previous line | A1ft |
| | | | (9) |
| | | | Total 9 |

| Question Number | Scheme | | Marks |
|-----------------|--|--|----------------|
| 7 | $P : z + 1 = 2z - 1 $ | | |
| | $z = x + iy \Rightarrow x + iy + 1 = 2(x + iy) - 1 $ | | |
| | $(x+1)^2 + y^2 = (2x-1)^2 + (2y)^2$ | M1: Correct use of Pythagoras A1: Any correct equation | M1A1 |
| | $Q : w = w - 1 + i $ | | |
| | $w = x + iy \Rightarrow x + iy = x + iy - 1 + i $ | | |
| | $x^2 + y^2 = (x-1)^2 + (y+1)^2 (\Rightarrow y = x-1)$ | M1: Correct use of Pythagoras. Allow with u and v instead of x and y A1: Any correct equation Must have x and y now. | M1A1 |
| | Alternative – M1: identifies perpendicular bisector of (0, 0) and (1, -1) A1: $y = x - 1$ | | |
| | | | |
| | | | |
| | $(x-1)^2 + (x-1)^2 = 1$ or $y^2 + y^2 = 1$ | Attempt to solve simultaneously tie obtain an equation in one variable and get to $x = \dots$ or $y = \dots$ | M1 |
| | $x = 1 + \frac{1}{\sqrt{2}}, 1 - \frac{1}{\sqrt{2}}$ or $y = \frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}$ | Both (oe) | A1 |
| | $\left(1 + \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$ and $\left(1 - \frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}}\right)$ | Both (oe) Pairs must be clearly identifiable but coordinate brackets not needed. | A1 |
| | | | (7) |
| | | | Total 7 |

| Question Number | Scheme | Marks | |
|-----------------|--|--|-----------------|
| 8. | $x = e^t$ | | |
| (a) | $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx}$ | Attempt to use an appropriate version of the chain rule | M1 |
| | $\frac{dy}{dx} = \frac{dy}{dt} \times \frac{1}{e^t} \left(= \frac{1}{x} \frac{dy}{dt} \right)$ | Oe | A1 |
| | $\frac{d^2y}{dx^2} = -\frac{1}{x^2} \frac{dy}{dt} + \frac{1}{x} \frac{d^2y}{dt^2} \frac{dt}{dx}$ or $\frac{d^2y}{dx^2} = -\frac{1}{x^2} \frac{dy}{dt} + \frac{1}{x^2} \frac{d^2y}{dt^2}$ | M1: Use of the product rule (penalise chain rule errors by loss of A mark or marks) $\left(\text{Note } t = \ln x \Rightarrow \frac{dt}{dx} = \frac{1}{x} \right)$ | M1 A1, A1 |
| | $x^2 \frac{d^2y}{dx^2} + 5x \frac{dy}{dx} + 13y = 0$ $\Rightarrow x^2 \cdot \frac{1}{x^2} \left(\frac{d^2y}{dt^2} - \frac{dy}{dt} \right) + 5x \frac{1}{x} \frac{dy}{dt} + 13y = 0$ $\Rightarrow \frac{d^2y}{dt^2} + 4 \frac{dy}{dt} + 13y = 0^*$ | M1: Substitutes their first and second derivatives into the given differential equation Depends on both M marks above A1: Correct completion to printed answer | ddM1A1 |
| | | | (7) |
| (b) | $m^2 + 4m + 13 = 0$ $\Rightarrow (m =) \frac{-4 \pm \sqrt{16 - 52}}{2}$ | Attempt to solve the auxiliary equation | M1 |
| | $(m =) -2 \pm 3i$ | Correct roots May be implied by a correct GS | A1 |
| | $y = e^{-2t} (A \cos 3t + B \sin 3t)$ or $y = Ae^{(-2+3i)t} + Be^{(-2-3i)t}$ | Correct GS | A1 |
| | $t = \ln x$ | | B1 |
| | $y = \frac{A \cos(3 \ln x) + B \sin(3 \ln x)}{x^2}$ or $y = Ae^{(-2+3i) \ln x} + Be^{(-2-3i) \ln x}$ | | A1 |
| | | | (5) |
| | | | Total 12 |

| Question Number | Scheme | | Marks |
|-----------------|---|---|-----------------|
| 9. | | | |
| (a) | $a = 2a \sin 2\theta \Rightarrow \sin 2\theta = \frac{1}{2} \Rightarrow 2\theta = \dots$ | $C_1 = C_2$ and attempt to solve for 2θ | M1 |
| | $\sin 2\theta = \frac{1}{2} \Rightarrow 2\theta = \frac{\pi}{6}, \frac{5\pi}{6}$ | $2\theta = \frac{\pi}{6}$ or $\frac{5\pi}{6}$ or both Decimals allowed (min 3 sf). | A1 |
| | $\left(a, \frac{\pi}{12}\right), \left(a, \frac{5\pi}{12}\right)$ | Both points Can be written $r = a, \theta = \frac{\pi}{12}, \frac{5\pi}{12}$ Decimals allowed (min 3 sf). | A1 |
| | | | (3) |
| (b) | $\frac{1}{2} \times a^2 \times \frac{\pi}{3}$ oe | Correct expression for the sector | B1 |
| | $\frac{1}{2} \int r^2 d\theta = \frac{1}{2} \int (2a \sin 2\theta)^2 d\theta$ | Use of correct formula Limits not needed (ignore any shown) | M1 |
| | $\cos 4\theta = 1 - 2\sin^2 2\theta$ $\Rightarrow \sin^2 2\theta = \frac{1}{2}(1 - \cos 4\theta)$ | Uses $\sin^2 2\theta = \frac{\pm 1 \pm \cos 4\theta}{2}$ | M1 |
| | $\int (1 - \cos 4\theta) d\theta = \theta - \frac{1}{4} \sin 4\theta$ | Correct integration Limits not needed (ignore any shown) | A1 |
| | $I = a^2 \left[\theta - \frac{1}{4} \sin 4\theta \right]_0^{\frac{\pi}{12}}$ $= a^2 \left\{ \left(\frac{\pi}{12} - \frac{1}{4} \sin 4 \cdot \frac{\pi}{12} \right) - (0) \right\}$ | An attempt to find one or both of the regions either side of the sector. ie uses limits $0, \frac{\pi}{12}$ and/or $\frac{5\pi}{12}, \frac{\pi}{2}$, limits to be substituted and subtracted (if non-zero after substitution). Limits to be used the correct way round. If two integrals seen award mark if either correct. Both previous method marks must have been scored. | ddM1 |
| | $R = 2I + \frac{a^2 \pi}{6} = 2a^2 \left(\frac{\pi}{12} - \frac{\sqrt{3}}{8} \right) + \frac{a^2 \pi}{6}$ | Correct strategy for the complete area (sector + $2I$). All areas must be positive. | M1 |
| | $R = \frac{1}{12} a^2 (4\pi - 3\sqrt{3})$ | If decimals seen anywhere (either in rt 3 or the limits) this mark is lost. | A1 |
| | | | (7) |
| | | | Total 10 |

