Please check the examination details below before entering your candidate information					
Candidate surname			Other name	25	
Pearson Edexcel International Advanced Level	Centre	Number		Candidate N	umber
Sample Assessment Materials for	r first te	aching S	eptember	2018	
(Time: 1 hour 30 minutes)		Paper R	eference V	WFM03/0	1
Mathematics International Advanced Subsidiary/Advanced Level Further Pure Mathematics FP3					
You must have: Mathematical Formulae and Stat	istical T	ables, cal	culator	To	otal Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided

 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over

\$59760A©2018 Pearson Education Ltd.
1/1/1/1/1/1/





Answer ALL questions. Write your answers in the spaces provided.

1.	The	curve	Ch	as ec	quation

$$y = 9\cosh x + 3\sinh x + 7x$$

Use differentiation to find the exact <i>x</i> cocanswer as a natural logarithm.	ordinate of the	Stational y	point of C,	giving you
answer as a natural logarithm.				(6

(6

	į	>		
			₹	

Question 1 continued		blan
zucstion i continucu		
		Q1
	(Total for Question 1 is 6 marks)	

(5)

(6)

2. An ellipse has equation

$$\frac{x^2}{25} + \frac{y^2}{4} = 1$$

The point P lies on the ellipse and has coordinates $(5\cos\theta, 2\sin\theta)$, $0 < \theta < \frac{\pi}{2}$

The line L is a normal to the ellipse at the point P.

(a) Show that an equation for L is

$$5x\sin\theta - 2y\cos\theta = 21\sin\theta\cos\theta$$

Given that the line L crosses the y-axis at the point Q and that M is the midpoint of PQ,

(b) find the exact area of triangle OPM, where O is the origin, giving your answer as a multiple of $\sin 2\theta$


uestion 2 continued	

uestion 2 continued	

Ķ	
	Š
ř	
	ď
	ă
	Ž
Ç	

	blank
Question 2 continued	
	Q2
(Total for Question 2 is 11 marks)	

3. Without using a calculator, find



(b)
$$\int_{-1}^{4} \frac{1}{\sqrt{4x^2 - 12x + 34}} dx$$
, giving your answer in the form $p \ln(q + r\sqrt{2})$,

where p, q and r are rational numbers to be found.

ľ	7	
•	•	

_
_
_
_
_
_
_
_
_
_
_
_
_

Ķ	
	Š
ř	
	ď
	ă
	Ž
Ç	

Question 3 continued	b
	—

estion 3 continued		

К	
	ż
Y V V	

Question 3 continued	blank
Question 5 continued	
	Q3
(Total for Question 3 is 12 marks)	

4.

$$\mathbf{M} = \begin{pmatrix} 1 & k & 0 \\ -1 & 1 & 1 \\ 1 & k & 3 \end{pmatrix}, \text{ where } k \text{ is a constant}$$

(a) Find M^{-1} in terms of k.

(5)

(4)

Hence, given that k = 0

(b) find the matrix N such that

$$\mathbf{MN} = \begin{pmatrix} 3 & 5 & 6 \\ 4 & -1 & 1 \\ 3 & 2 & -3 \end{pmatrix}$$

	L
uestion 4 continued	

estion 4 continued	

			ì		
			į		
	ĕ				
			Ė		
			þ		
	é	ě	ī	Z	Š
	á	ě	ě	ś	ì
	é			į	į
	6			Ž	Ì
	í			Ž	į
	í			į	į
	4				
	i				
	i				
	i				
	(

(Total for Question 4 is 9 marks)	uestion 4 continued	

(2)

5. Given that $y = \operatorname{artanh}(\cos x)$

(a) show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\mathrm{cosec}\,x$$

(b) Hence find the exact value of

$$\int_0^{\frac{\pi}{6}} \cos x \, \operatorname{artanh}(\cos x) \, \mathrm{d}x$$

giving your answer in the form $a \ln(b + c\sqrt{3}) + d\pi$, where a, b, c and d are rational numbers to be found.

(5)

	blank
Question 5 continued	Olalik
Question 5 continued	

uestion 5 continued		
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_

				ø
			ĺ	
				i
	į			

Question 5 continued		Leave
Question 5 continued		
		0.5
		Q5
	(Total for Question 5 is 7 marks)	

6.	The coordinates of the points A , B and C relative to a fixed origin O are $(1, 2, 3)$, $(-1, 3, 4)$ and $(2, 1, 6)$ respectively. The plane Π contains the points A , B and C .
	(a) Find a cartesian equation of the plane Π . (5)
	The point D has coordinates $(k, 4, 14)$ where k is a positive constant.
	Given that the volume of the tetrahedron ABCD is 6 cubic units,
	(b) find the value of k . (4)

	b
uestion 6 continued	

estion 6 continued	

Question 6 continued	blank
	Q6
(Total for Question 6 is 9 marks)	

7. The curve C has parametric equations

$$x = 3t^4, \quad y = 4t^3, \qquad 0 \leqslant t \leqslant 1$$

The curve C is rotated through 2π radians about the x-axis. The area of the curved surface generated is S.

(a) Show that

$$S = k\pi \int_0^1 t^5 (t^2 + 1)^{\frac{1}{2}} dt$$

where k is a constant to be found.

(4)

(b) Use the substitution $u^2 = t^2 + 1$ to find the value of S, giving your answer in the form $p\pi(11\sqrt{2} - 4)$ where p is a rational number to be found.

(7)

		į	

	blank
Question 7 continued	Olalik
Question / continued	

estion 7 continued	

	g	
	M	
	ŝ	
	Ń	
	P	
×.		

Question 7 continued Question 7 continued	Overtion 7 continued		blank
	Question 7 continued		
			07
(Total for Question 7 is 11 marks)			
		(Total for Question 7 is 11 marks)	

(5)

(5)

 $I_n = \int_0^{\ln 2} \tanh^{2n} x \, \mathrm{d}x, \quad n \geqslant 0$ 8.

$$I_n = \int_0^\infty \tanh^{2n} x \, \mathrm{d}x, \quad n \geqslant 0$$

(a) Show that, for $n \ge 1$

$$I_n = I_{n-1} - \frac{1}{2n-1} \left(\frac{3}{5}\right)^{2n-1}$$

(b) Hence show that

$$\int_0^{\ln 2} \tanh^4 x \, \mathrm{d}x = p + \ln 2$$

where p is a rational number to be found.

		`

	į			
			È	
			2	
			7	
			2	

estion 8 continued	

uestion 8 continued	

	Leave
	blank
Question 8 continued	

Question 8 continued	Leave blank
	Q8
(T-1-18 01 0:-10 1)	
(Total for Question 8 is 10 marks) TOTAL FOR PAPER IS 75 MARKS	