

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Monday 21 January 2019

Afternoon (Time: 1 hour 30 minutes)

Paper Reference **WME01/01**

Mechanics M1
Advanced/Advanced Subsidiary

You must have:

Mathematical Formulae and Statistical Tables (Blue)

Total Marks

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P54829A

©2019 Pearson Education Ltd.

1/1/1/



Pearson

Leave blank

1. Two particles, *A* and *B*, have masses $2m$ and $3m$ respectively. They are moving towards each other in opposite directions along the same straight line on a smooth horizontal plane when they collide directly. Immediately before they collide, the speed of *A* is $3u$ and the speed of *B* is u . As a result of the collision, the speed of *A* is halved and the direction of motion of each particle is reversed.

(i) Find the speed of *B* immediately after the collision.

(ii) Find the magnitude of the impulse exerted on *A* by *B* in the collision.

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

2. [In this question \mathbf{i} and \mathbf{j} are horizontal unit vectors due east and due north respectively and position vectors are given relative to a fixed origin O .]

At time $t = 0$, a bird A leaves its nest, that is located at the point with position vector $(20\mathbf{i} - 17\mathbf{j})\text{m}$, and flies with constant velocity $(-6\mathbf{i} + 7\mathbf{j})\text{m s}^{-1}$. At the same time a second bird B leaves its nest which is located at the point with position vector $(-8\mathbf{i} + 9\mathbf{j})\text{m}$ and flies with constant velocity $(p\mathbf{i} + 2p\mathbf{j})\text{m s}^{-1}$, where p is a constant. At time $t = 4$ s, bird B is south west of bird A .

- (a) Find the direction of motion of A , giving your answer as a bearing to the nearest degree. **(3)**

- (b) Find the speed of B . **(10)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave blank

Question 2 continued

A large rectangular area with horizontal lines for writing.

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

3.

**Figure 1**

A lift of mass M kg is being raised by a vertical cable attached to the top of the lift. A person of mass m kg stands on the floor inside the lift, as shown in Figure 1. The lift ascends vertically with constant acceleration 1.4 m s^{-2} . The tension in the cable is 2800 N and the person experiences a constant normal reaction of magnitude 560 N from the floor of the lift. The cable is modelled as being light and inextensible, the person is modelled as a particle and air resistance is negligible.

(a) Write down an equation of motion for the person only. (2)

(b) Write down an equation of motion for the lift only. (2)

(c) Hence, or otherwise, find
(i) the value of m ,
(ii) the value of M . (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

6. A train travels for a total of 270 s along a straight horizontal track between two stations *A* and *B*. The train starts from rest at *A* and moves with constant acceleration for 60 s until it reaches a speed of $V \text{ m s}^{-1}$. The train then travels at this constant speed $V \text{ m s}^{-1}$ before it moves with constant deceleration for 30 s, coming to rest at *B*.

(a) Sketch below a speed-time graph for the journey of the train between the two stations *A* and *B*.

(2)

Given that the distance between the two stations is 4.5 km,

(b) find the value of V ,

(3)

(c) find how long it takes the train to travel from station *A* to the point that is exactly halfway between the two stations.

(4)

The train is travelling at speed $\frac{1}{4} V \text{ m s}^{-1}$ at times T_1 seconds and T_2 seconds after leaving station *A*.

(d) Find the value of T_1 and the value of T_2

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



