

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

Pearson Edexcel International Advanced Level

Thursday 11 January 2024

Afternoon (Time: 1 hour 30 minutes)

Paper
reference

WME02/01

Mathematics

**International Advanced Subsidiary/Advanced Level
Mechanics M2**

You must have:

Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P74318A

©2024 Pearson Education Ltd.
S:1/1/1/



Pearson

1. A particle P moves along a straight line. The fixed point O is on the line. At time t seconds, $t > 0$, the displacement of P from O is x metres, where

$$x = 2t^3 - 21t^2 + 60t$$

Find

- (a) the values of t for which P is instantaneously at rest (4)
- (b) the distance travelled by P in the interval $1 \leq t \leq 3$ (2)
- (c) the magnitude of the acceleration of P at the instant when $t = 3$ (2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2. [In this question, \mathbf{i} and \mathbf{j} are horizontal perpendicular unit vectors.]

A particle Q of mass 0.5 kg is moving on a smooth horizontal surface. Particle Q is moving with velocity $(3\mathbf{i} + \mathbf{j}) \text{ m s}^{-1}$ when it receives an impulse of $(2\mathbf{i} + 5\mathbf{j}) \text{ N s}$.

(a) Find the speed of Q immediately after receiving the impulse. (4)

As a result of receiving the impulse, the direction of motion of Q is turned through an angle θ°

(b) Find the value of θ (2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



5.




Figure 4

A van of mass 600 kg is moving up a straight road inclined at an angle α to the horizontal, where $\sin \alpha = \frac{1}{14}$. The van is towing a trailer of mass 200 kg . The trailer is attached to the van by a rigid towbar which is parallel to the direction of motion of the van and the trailer, as shown in Figure 4.

The resistance to the motion of the van from non-gravitational forces is modelled as a constant force of magnitude 250 N .

The resistance to the motion of the trailer from non-gravitational forces is modelled as a constant force of magnitude 150 N .

The towbar is modelled as a light rod.

At the instant when the speed of the van is 16 ms^{-1} , the engine of the van is working at a rate of 10 kW .

(a) Find the deceleration of the van at this instant. (5)

(b) Find the tension in the towbar at this instant. (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



6.




Figure 5

A uniform beam AB , of weight 40 N and length 7 m , rests with end A on rough horizontal ground.

The beam rests on a smooth horizontal peg at C , with $AC = 5\text{ m}$, as shown in Figure 5.

The beam is inclined at an angle θ to the ground, where $\sin \theta = \frac{3}{5}$

The beam is modelled as a rod that lies in a vertical plane perpendicular to the peg.

The normal reaction between the beam and the peg at C has magnitude P newtons.

Using the model,

- (a) show that $P = 22.4$ (3)
- (b) find the magnitude of the resultant force acting on the beam at A . (6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the answer to Question 6.



P 7 4 3 1 8 A 0 1 9 2 8

7. Particle P has mass m and particle Q has mass $5m$.

The particles are moving in the same direction along the same straight line on a smooth horizontal surface.

Particle P collides directly with particle Q .

Immediately **before** the collision, the speed of P is $6u$ and the speed of Q is u .

Immediately **after** the collision, the speed of P is x and the speed of Q is y .

The direction of motion of P is reversed as a result of the collision.

The coefficient of restitution between P and Q is e .

(a) Find the complete range of possible values of e . (7)

Given that $e = \frac{3}{5}$

(b) find the total kinetic energy lost in the collision between P and Q . (4)

After the collision, Q hits a smooth fixed vertical wall that is perpendicular to the direction of motion of Q .

Particle Q rebounds.

The coefficient of restitution between Q and the wall is f .

Given that there is a second collision between P and Q ,

(c) find the complete range of possible values of f . (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



8.

[In this question \mathbf{i} and \mathbf{j} are unit vectors, with \mathbf{i} horizontal and \mathbf{j} vertical.]




Figure 6

The fixed points A and B lie on horizontal ground.

At time $t = 0$, a particle P is projected from A with velocity $(4\mathbf{i} + 7\mathbf{j}) \text{ m s}^{-1}$

Particle P moves freely under gravity and hits the ground at B , as shown in Figure 6.

- (a) Find the distance AB . (4)

The speed of P is less than 5 m s^{-1} for an interval of length T seconds.

- (b) Find the value of T (3)

At the instant when the direction of motion of P is perpendicular to the initial direction of motion of P , the particle is h metres above the ground.

- (c) Find the value of h . (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



