

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--

Thursday 21 May 2020

Afternoon (Time: 1 hour 30 minutes)

Paper Reference **WME02/01**

Mathematics

International Advanced Subsidiary/Advanced Level Mechanics M2

You must have:

Mathematical Formulae and Statistical Tables (Blue), calculator

Total Marks

**Candidates may use any calculator permitted by Pearson regulations.
Calculators must not have the facility for symbolic algebra manipulation,
differentiation and integration, or have retrievable mathematical
formulae stored in them.**

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need*.
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
– *use this as a guide as to how much time to spend on each question*.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ▶

P65265A

©2020 Pearson Education Ltd.

1/1/1



P 6 5 2 6 5 A 0 1 3 2



Pearson

Leave
blank

1. A particle of mass 2 kg is moving with velocity $(5\mathbf{i} + 3\mathbf{j}) \text{ m s}^{-1}$ when it receives an impulse $\mathbf{I} \text{ N s}$, such that $\mathbf{I} = a\mathbf{i} + b\mathbf{j}$

Immediately after receiving the impulse, the particle is moving with velocity $\lambda(\mathbf{i} + \mathbf{j}) \text{ m s}^{-1}$, where λ is a constant.

Given that the magnitude of \mathbf{I} is $\sqrt{40}$, find the two possible impulses.

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q1

(Total 5 marks)



Leave
blank

2. A truck of weight 9000N is travelling up a hill on a straight road that is inclined at an angle θ to the horizontal, where $\sin \theta = \frac{1}{15}$

When the truck travels up the hill with the engine working at $3P$ watts, the truck is moving at a constant speed of 12 m s^{-1}

Later on, the truck travels down the hill along the same road, with the engine working at P watts. At the instant when the speed of the truck is 12 m s^{-1} , the acceleration of the truck is $\frac{g}{20}$

The resistance to motion of the truck from non-gravitational forces is a constant force of magnitude R newtons in all circumstances.

Find (i) the value of P ,

(ii) the value of R .

(9)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 5 3 2

Leave
blank

Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 2 continuedLeave
blank

Q2


(Total 9 marks)



P 6 5 2 6 5 A 0 7 3 2

Leave
blank

3.

**Figure 1**

A uniform rod AB , of mass 25 kg and length 3 m, has end A resting on rough horizontal ground. The end B rests against a rough vertical wall.

The rod is in a vertical plane perpendicular to the wall.

The coefficient of friction between the rod and the ground is $\frac{4}{5}$

The coefficient of friction between the rod and the wall is $\frac{3}{5}$

The rod rests in limiting equilibrium.

The rod is at an angle of θ to the ground, as shown in Figure 1.

Find the exact value of $\tan \theta$.

(9)



Leave
blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 9 3 2

Leave
blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q3


(Total 9 marks)



P 6 5 2 6 5 A 0 1 1 3 2

Leave
blank

4.

**Figure 2**

The uniform lamina L , shown shaded in Figure 2, is formed by removing the square $PQRV$, of side $2a$, and the square $RSTU$, of side $4a$, from a uniform square lamina $ABCD$, of side $8a$. The lines QRU and VRS are straight. The side AD is parallel to PV and the side AB is parallel to PQ . The distance between AD and PV is a and the distance between AB and PQ is a . The centre of mass of L is at the point G .

- (a) Show that the distance of G from the side AD is $\frac{42}{11}a$ (5)

The mass of L is M . A particle of mass kM is attached to L at C .

The lamina, with the attached particle, is freely suspended from B and hangs in equilibrium with BC making an angle of 45° with the horizontal.

- (b) Find the value of k . (4)
-
-
-
-
-
-
-
-
-
-
-
-
-
-
-
-
-
-
-
-

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 1 3 3 2

Leave
blank

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q4

(Total 9 marks)

15

Turn over ►



P 6 5 2 6 5 A 0 1 5 3 2

Leave
blank

5. At time t seconds ($t \geq 0$), a particle P has velocity $\mathbf{v} \text{ m s}^{-1}$, where

$$\mathbf{v} = (3t^2 - 9t + 6)\mathbf{i} + (t^2 + t - 6)\mathbf{j}$$

- (a) Find the acceleration of P when $t = 3$

(3)

When $t = 0$, P is at the fixed point O .

The particle comes to instantaneous rest at the point A .

- (b) Find the distance OA .

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 1 7 3 2

Leave
blank

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q5

(Total 10 marks)



P 6 5 2 6 5 A 0 1 9 3 2

Leave
blank

6.




Figure 3

A rough straight ramp is fixed to horizontal ground. The ramp has length 15 m and is inclined at an angle α to the ground, where $\tan \alpha = \frac{5}{12}$. The line AB is a line of greatest slope of the ramp, where A is at the bottom of the ramp, and B is at the top of the ramp, as shown in Figure 3.

A particle P of mass 6 kg is projected up the ramp with speed 14 m s^{-1} from A in a straight line towards B . The coefficient of friction between P and the ramp is 0.25

- (a) Find the work done against friction as P moves from A to B .

(3)

At the instant P reaches B , the speed of P is $v \text{ m s}^{-1}$. After leaving the ramp at B , the particle P moves freely under gravity until it hits the horizontal ground at the point C . Immediately before hitting the ground at C , the speed of P is $w \text{ m s}^{-1}$

- (b) Use the work-energy principle to find

- (i) the value of v ,
- (ii) the value of w .

(7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 2 1 3 2

Leave
blank

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q6

(Total 10 marks)



P 6 5 2 6 5 A 0 2 3 3 2

Leave
blank

7. Particle *A* of mass $3m$ is moving in a straight line with speed $2u$ on a smooth horizontal surface. Particle *A* collides directly with particle *B* of mass m , which is moving along the same straight line and in the same direction as *A*.

Immediately before the collision, the speed of *B* is u .

As a result of the collision, the direction of motion of *B* is unchanged and the kinetic energy gained by *B* is $\frac{48}{25}mu^2$

- (a) Find the coefficient of restitution between *A* and *B*.

(8)

After the collision, *B* hits a smooth fixed vertical wall that is perpendicular to the direction of motion of *B*. The coefficient of restitution between *B* and the wall is f .

Given that the speed of *B* immediately after first hitting the wall is equal to the speed of *A* immediately after its first collision with *B*,

- (b) find the value of f .

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q7

(Total 10 marks)



P 6 5 2 6 5 A 0 2 7 3 2

Leave
blank

8. [In this question, the unit vectors \mathbf{i} and \mathbf{j} are in a vertical plane, with \mathbf{i} being horizontal and \mathbf{j} being vertically upwards.]




Figure 4

At time $t = 0$, a small ball is projected from a fixed point O on horizontal ground. The ball is projected from O with velocity $(p\mathbf{i} + q\mathbf{j}) \text{ m s}^{-1}$, where p and q are positive constants. The ball moves freely under gravity.

At time $t = 3$ seconds, the ball passes through the point A with velocity $(8\mathbf{i} - 12\mathbf{j}) \text{ m s}^{-1}$, as shown in Figure 4.

- (a) Find the speed of the ball at the instant it is projected from O .

(5)

For an interval of T seconds the speed, $v \text{ m s}^{-1}$, of the ball is such that $v \leqslant 10$

- (b) Find the value of T .

(4)

At the point B on the path of the ball, the direction of motion of the ball is perpendicular to the direction of motion of the ball at A .

- (c) Find the vertical height of B above A .

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 2 9 3 2

Leave
blank

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 5 2 6 5 A 0 3 1 3 2

Leave
blank

Question 8 continued

Q8

(Total 13 marks)

END

TOTAL FOR PAPER: 75 MARKS



P 6 5 2 6 5 A 0 3 2 3 2

DO NOT WRITE IN THIS AREA