

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--

--	--	--	--

Pearson Edexcel International Advanced Level

Thursday 26 October 2023

Afternoon (Time: 1 hour 30 minutes)

Paper
reference

WME02/01



Mathematics

International Advanced Subsidiary/Advanced Level Mechanics M2

You must have:

Mathematical Formulae and Statistical Tables (Yellow), calculator

Total Marks

**Candidates may use any calculator permitted by Pearson regulations.
Calculators must not have the facility for symbolic algebra manipulation,
differentiation and integration, or have retrievable mathematical
formulae stored in them.**

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need*.
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question*.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P74323A

©2023 Pearson Education Ltd.
Z:1/1/1/



Pearson

1. At time t seconds, $t > 0$, a particle P is at the point with position vector \mathbf{r} m, where

$$\mathbf{r} = \left(t^4 - 8t^2 \right) \mathbf{i} + \left(6t^2 - 2t^{\frac{3}{2}} \right) \mathbf{j}$$

(a) Find the velocity of P when P is moving in a direction parallel to the vector \mathbf{j}

(4)

(b) Find the acceleration of P when $t = 4$

(3)

DO NOT WRITE IN THIS AREA



Question 1 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 1 is 7 marks)

P 7 4 3 2 3 A 0 3 2 8

2.


**Figure 1**

Figure 1 shows a template where

- $PQUY$ is a uniform square lamina with sides of length $4a$
- $RSTU$ is a uniform square lamina with sides of length $2a$
- $VWXY$ is a uniform square lamina with sides of length $2a$
- the three squares all lie in the same plane
- the mass per unit area of $VWXY$ is **double** the mass per unit area of $PQUY$
- the mass per unit area of $RSTU$ is **double** the mass per unit area of $PQUY$
- the distance of the centre of mass of the template from PX is d

$$(a) \text{ Show that } d = \frac{5}{2}a \quad (5)$$

The template is freely pivoted about Q and hangs in equilibrium with PQ at an angle of θ to the downward vertical.

$$(b) \text{ Find the value of } \tan \theta \quad (6)$$

The mass of the template is M

The template is still freely pivoted about Q , but it is now held in equilibrium, with PQ vertical, by a horizontal force of magnitude F which acts on the template at X . The line of action of the force lies in the same plane as the template.

$$(c) \text{ Find } F \text{ in terms of } M \text{ and } g \quad (3)$$



Question 2 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 5 2 8

Question 2 continued

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 2 is 14 marks)

P 7 4 3 2 3 A 0 7 2 8

3.

**Figure 2**

A particle Q of mass 0.25 kg is moving in a straight line on a smooth horizontal surface with speed 8 m s^{-1} when it receives an impulse of magnitude $I \text{ N s}$.

The impulse acts parallel to the horizontal surface and at 60° to the original direction of motion of Q .

Immediately after receiving the impulse, the speed of Q is 12 m s^{-1}

As a result of receiving the impulse, the direction of motion of Q is turned through α° , as shown in Figure 2.

Find the value of I

(6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 9 2 8

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 3 is 6 marks)

P 7 4 3 2 3 A 0 1 1 2 8

4.

[In this question \mathbf{i} and \mathbf{j} are unit vectors, with \mathbf{i} horizontal and \mathbf{j} vertical.]




Figure 3

The fixed points A and B lie on horizontal ground.

At time $t = 0$, a particle P is projected from A with velocity $(4\mathbf{i} + 4\mathbf{j}) \text{ m s}^{-1}$

Particle P moves freely under gravity and hits the ground at B , as shown in Figure 3.

At time T_1 seconds, P is at its highest point above the ground.

(a) Find the value of T_1

(2)

At time $t = 0$, a particle Q is also projected from A but with velocity $(5\mathbf{i} + 7\mathbf{j}) \text{ m s}^{-1}$

Particle Q moves freely under gravity.

(b) Find the vertical distance between Q and P at time T_1 seconds, giving your answer to 2 significant figures.

(3)

At the instant when particle P reaches B , particle Q is moving at α° below the horizontal.

(c) Find the value of α .

(4)

At time T_2 seconds, the direction of motion of Q is perpendicular to the initial direction of motion of Q .

(d) Find the value of T_2

(3)



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 1 3 2 8

Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 is 12 marks)

P 7 4 3 2 3 A 0 1 5 2 8

5. A cyclist is travelling on a straight horizontal road and working at a constant rate of 500W.

The total mass of the cyclist and her cycle is 80kg.

The total resistance to the motion of the cyclist is modelled as a constant force of magnitude 60N.

- (a) Using this model, find the acceleration of the cyclist at the instant when her speed is 6 m s^{-1}

(4)

On the following day, the cyclist travels up a straight road from a point *A* to a point *B*.

The distance from *A* to *B* is 20km.

Point *A* is 500m above sea level and point *B* is 800m above sea level.

The cyclist starts from rest at *A*.

At the instant she reaches *B* her speed is 8 m s^{-1}

The total resistance to the motion of the cyclist from non-gravitational forces is modelled as a constant force of magnitude 60N.

- (b) Using this model, find the total work done by the cyclist in the journey from *A* to *B*.

(5)

Later on, the cyclist is travelling up a straight road which is inclined at an angle α to the horizontal, where $\sin \alpha = \frac{1}{20}$

The cyclist is now working at a constant rate of P watts and has a constant speed of 7 m s^{-1}

The total resistance to the motion of the cyclist from non-gravitational forces is again modelled as a constant force of magnitude 60N.

- (c) Using this model, find the value of P

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 1 7 2 8

Question 5 continued

DO NOT WRITE IN THIS AREA



Question 5 continued

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 5 is 13 marks)

P 7 4 3 2 3 A 0 1 9 2 8

6.

**Figure 4**

A uniform rod AB has length $8a$ and weight W .

The end A of the rod is freely hinged to a fixed point on a vertical wall.

A particle of weight $\frac{1}{4}W$ is attached to the rod at B .

A light inelastic string of length $5a$ has one end attached to the rod at the point C , where $AC = 5a$.

The other end of the string is attached to the wall at the point D , where D is above A and $AD = 5a$, as shown in Figure 4.

The rod rests in equilibrium.

The tension in the string is T .

- (a) Show that $T = \frac{6}{5}W$ (3)

- (b) Find, in terms of W , the magnitude of the force exerted on the rod by the hinge at A . (6)



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 2 1 2 8

Question 6 continued

DO NOT WRITE IN THIS AREA



Question 6 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 6 is 9 marks)

P 7 4 3 2 3 A 0 2 3 2 8

7. Particle P has mass $4m$ and particle Q has mass $2m$.

The particles are moving in opposite directions along the same straight line on a smooth horizontal surface.

Particle P collides directly with particle Q .

Immediately **before** the collision, the speed of P is $2u$ and the speed of Q is $3u$.

Immediately **after** the collision, the speed of P is x and the speed of Q is y .

The direction of motion of each particle is reversed as a result of the collision.

The total kinetic energy of P and Q after the collision is half of the total kinetic energy of P and Q before the collision.

- (a) Show that $y = \frac{8}{3}u$ (6)

The coefficient of restitution between P and Q is e .

- (b) Find the value of e . (3)

After the collision, Q hits a smooth fixed vertical wall that is perpendicular to the direction of motion of Q .

Particle Q rebounds.

The coefficient of restitution between Q and the wall is f .

Given that there is no second collision between P and Q ,

- (c) find the range of possible values of f . (3)

Given that $f = \frac{1}{4}$

- (d) find, in terms of m and u , the magnitude of the impulse received by Q as a result of its impact with the wall. (2)



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 2 5 2 8

Question 7 continued

DO NOT WRITE IN THIS AREA



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 4 3 2 3 A 0 2 7 2 8

Question 7 continued

DO NOT WRITE IN THIS AREA

(Total for Question 7 is 14 marks)

TOTAL FOR PAPER IS 75 MARKS