

2. A curve C has the equation

$$x^3 - 3xy - x + y^3 - 11 = 0$$

Find an equation of the tangent to C at the point $(2, -1)$, giving your answer in the form $ax + by + c = 0$, where a , b and c are integers.

(6)

$$\frac{d}{dx}(x^3 - 3xy - x + y^3 - 11)$$

$$3x^2 - 3x \frac{dy}{dx} - 3y - 1 + 3y^2 \frac{dy}{dx} = 0$$

$$\Rightarrow (3y^2 - 3x) \frac{dy}{dx} = 1 + 3y - 3x^2$$

$$\therefore \frac{dy}{dx} = \frac{1 + 3y - 3x^2}{3y^2 - 3x} \quad \text{at } (2, -1)$$

$$m_t = \frac{1 + 3(-1) - 12}{3(-1)^2 - 3(2)} = \frac{14}{3}$$

$$y + 1 = \frac{14}{3}(x - 2) \quad \Rightarrow \quad 3y + 3 = 14x - 28$$

$$14x - 3y - 31 = 0$$

3. Given that

$$y = \frac{\cos 2\theta}{1 + \sin 2\theta}, \quad -\frac{\pi}{4} < \theta < \frac{3\pi}{4}$$

show that

$$\frac{dy}{d\theta} = \frac{a}{1 + \sin 2\theta}, \quad -\frac{\pi}{4} < \theta < \frac{3\pi}{4}$$

where a is a constant to be determined.

(4)

$$u = \cos 2\theta \quad v = 1 + \sin 2\theta$$

$$u' = -2\sin 2\theta \quad v' = 2\cos 2\theta$$

$$\therefore \frac{du}{d\theta} = \frac{-2\sin 2\theta(1 + \sin 2\theta) - 2(\cos^2 2\theta)}{(1 + \sin 2\theta)^2}$$

$$= \frac{-2\sin 2\theta - 2(\sin^2 2\theta + \cos^2 2\theta)}{(1 + \sin 2\theta)^2}$$

$$= \frac{-2(1 + \sin 2\theta)}{(1 + \sin 2\theta)^2} = \frac{-2}{1 + \sin 2\theta}$$

4. Find

$$(a) \int (2x + 3)^{12} dx$$

(2)

$$(b) \int \frac{5x}{4x^2 + 1} dx$$

(2)

$$a) \frac{1}{13} (2x+3)^{13} \div 2 = \frac{1}{26} (2x+3)^{13} + C$$

$$b) \frac{5}{8} \int \frac{8x}{4x^2+1} dx = \frac{5}{8} \ln(4x^2+1) + C$$

5.

$$f(x) = (8 + 27x^3)^{\frac{1}{3}}, \quad |x| < \frac{2}{3}$$

Find the first three non-zero terms of the binomial expansion of $f(x)$ in ascending powers of x . Give each coefficient as a simplified fraction.

$$8^{\frac{1}{3}} \left(1 + \frac{27}{8}x^3\right)^{\frac{1}{3}} \quad (5)$$

$$= 2 \left[1 + \left(\frac{1}{3}\right)\left(\frac{27}{8}x^3\right) + \frac{\left(\frac{1}{3}\right)\left(-\frac{2}{3}\right)}{2} \left(\frac{27}{8}x^3\right)^2 \right]$$

$$= 2 + \frac{9}{4}x^3 + -\frac{81}{32}x^6$$

6. (a) Express $\frac{5-4x}{(2x-1)(x+1)}$ in partial fractions.

(3)

(b) (i) Find a general solution of the differential equation

$$(2x-1)(x+1)\frac{dy}{dx} = (5-4x)y, \quad x > \frac{1}{2}$$

Given that $y = 4$ when $x = 2$,

(ii) find the particular solution of this differential equation.
Give your answer in the form $y = f(x)$.

(7)

$$5-4x = \frac{A}{2x-1} + \frac{B}{x+1}$$

$$5-4x = A(x+1) + B(2x-1)$$

$$x = -1 \Rightarrow 9 = -3B \quad \therefore B = -\frac{3}{2}$$

$$x = \frac{1}{2} \Rightarrow 3 = \frac{1}{2}A \quad A = \frac{2}{2} = 1$$

$$\therefore \frac{1}{2x-1} - \frac{3}{x+1}$$

$$b) \int \frac{1}{y} dy = \int \frac{(5-4x)}{(2x-1)(x+1)} dx$$

$$\therefore \ln y = \ln(2x-1) - 3\ln(x+1) + C$$

$$(2, 4) \quad \ln 4 = \ln 3 - 3\ln 3 + C \quad \ln 4 = -2\ln 3 + C$$

$$C = \ln 4 + 2\ln 3 = \ln 4 + \ln 9 = \ln 36$$

$$\therefore \ln y = \ln(2x-1) - \ln(x+1)^3 + \ln 36$$

$$\therefore y = \frac{36(2x-1)}{(x+1)^3}$$

7. The function f is defined by

$$f: x \mapsto \frac{3x-5}{x+1}, \quad x \in \mathbb{R}, x \neq -1$$

(a) Find an expression for $f^{-1}(x)$

(3)

(b) Show that

$$ff(x) = \frac{x+a}{x-1}, \quad x \in \mathbb{R}, x \neq -1, x \neq 1$$

where a is an integer to be determined.

(4)

The function g is defined by

$$g: x \mapsto x^2 - 3x, \quad x \in \mathbb{R}, 0 \leq x \leq 5$$

(c) Find the value of $fg(2)$

(2)

(d) Find the range of g

(3)

$$x = \frac{3y-5}{y+1} \Rightarrow xy + x = 3y - 5 \Rightarrow 3y - xy = x + 5$$

$$y(3-x) = x+5 \quad \therefore y = \frac{x+5}{3-x}$$


$$b) \quad ff(x) = \frac{3\left(\frac{3x-5}{x+1}\right) - 5}{\left(\frac{3x-5}{x+1}\right) + 1} = \frac{9x - 15 - 5(x+1)}{(x+1)} = \frac{9x - 15 - 5x - 5}{(x+1)}$$

$$= \frac{4x - 20}{(x+1)}$$

$$= \frac{4x - 20}{4x - 4} = \frac{4(x-5)}{4(x-1)} = \frac{x-5}{x-1}$$

$$c) \quad fg(2) = f(2^2 - 3(2)) = f(-2) = \frac{-6-5}{-2+1} = \frac{-11}{-1} = 11$$

$$d) \quad g(x) = x(x-3)$$



$$g(1.5) = \frac{-9}{4} \quad -\frac{9}{4} \leq g(x) \leq 10$$

$$g(3) = 10$$

8. The volume V of a spherical balloon is increasing at a constant rate of $250 \text{ cm}^3 \text{ s}^{-1}$. Find the rate of increase of the radius of the balloon, in cm s^{-1} , at the instant when the volume of the balloon is $12\,000 \text{ cm}^3$. Give your answer to 2 significant figures.

(5)

[You may assume that the volume V of a sphere of radius r is given by the formula $V = \frac{4}{3}\pi r^3$.]

$$\frac{dV}{dt} = 250 \quad \text{find } \frac{dr}{dt} \text{ when } V = 12000$$

$$\frac{dV}{dr} = 4\pi r^2 \quad \frac{dr}{dt} = \frac{dr}{dV} \times \frac{dV}{dt}$$

$$\frac{dr}{dt} = \left(\frac{1}{4\pi r^2}\right)(250) = \frac{250}{4\pi r^2} = 0.099 \text{ cm/sec}$$

$$V = 12000 = \frac{4}{3}\pi r^3$$

$$r = 14.2 \dots$$

9.




Figure 1

Figure 1 shows a sketch of part of the curve with equation $y = e^{\sqrt{x}}$, $x > 0$

The finite region R , shown shaded in Figure 1, is bounded by the curve, the x -axis and the lines $x = 4$ and $x = 9$

(a) Use the trapezium rule, with 5 strips of equal width, to obtain an estimate for the area of R , giving your answer to 2 decimal places.

(4)

(b) Use the substitution $u = \sqrt{x}$ to find, by integrating, the exact value for the area of R .

(7)

$$\begin{array}{c|c|c|c|c|c|c}
 x & 4 & 5 & 6 & 7 & 8 & 9 \\
 \hline
 y & e^2 & e^{\sqrt{5}} & e^{\sqrt{6}} & e^{\sqrt{7}} & e^{\sqrt{8}} & e^3
 \end{array}$$

$$\text{Area} \approx \frac{1}{2} (e^2 + 2(e^{\sqrt{5}} + \dots) + e^3) \approx 65.69$$

$$b) \quad u = x^{\frac{1}{2}} \quad \frac{du}{dx} = \frac{1}{2} x^{-\frac{1}{2}} \quad dx = 2x^{\frac{1}{2}} du = 2u du$$

$$\therefore \int_4^9 y dx = \int_2^3 e^u \times 2u du \quad \begin{array}{l} u = 2u \\ u' = 2 \end{array} \quad \begin{array}{l} v = e^u \\ v' = e^u \end{array}$$

$$= [2ue^u - 2e^u]_2^3 = [2ue^u - 2e^u]_2^3$$

$$= [2e^u (u - 1)]_2^3 = \frac{4e^3 - 2e^2}{2}$$

10. (a) Use the identity for $\sin(A + B)$ to prove that

$$\sin 2A \equiv 2 \sin A \cos A \quad (2)$$

(b) Show that

$$\frac{d}{dx} [\ln(\tan(\frac{1}{2}x))] = \operatorname{cosec} x \quad (4)$$

A curve C has the equation

$$y = \ln(\tan(\frac{1}{2}x)) - 3 \sin x, \quad 0 < x < \pi$$

(c) Find the x coordinates of the points on C where $\frac{dy}{dx} = 0$

Give your answers to 3 decimal places.

(Solutions based entirely on graphical or numerical methods are not acceptable.)

(6)

a) let $A=B$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A+A) = \sin A \cos A + \cos A \sin A$$

$$\sin 2A \equiv 2 \sin A \cos A \quad \#$$

b) $\frac{d}{dx} (\ln[\tan(\frac{1}{2}x)]) = \frac{\frac{1}{2} \sec^2(\frac{1}{2}x)}{\tan(\frac{1}{2}x)} = \frac{1}{2(\cos(\frac{1}{2}x))^2} \times \frac{\cos(\frac{1}{2}x)}{\sin(\frac{1}{2}x)}$

$$= \frac{1}{2 \sin(\frac{1}{2}x) \cos(\frac{1}{2}x)} = \frac{1}{\sin x} = \operatorname{cosec} x \quad \#$$

c) $\frac{dy}{dx} = 0 \Rightarrow \operatorname{cosec} x - 3 \cos x = 0 \Rightarrow 3 \cos x = \frac{1}{\sin x}$

$$\Rightarrow 3 \sin x \cos x = 1 \Rightarrow 2 \sin x \cos x = \frac{2}{3}$$

$$\Rightarrow \sin 2x = \frac{2}{3} \Rightarrow 2x = \sin^{-1}(\frac{2}{3}) = 0.7297...; \quad 2.412$$

$$\therefore x = 0.3648...$$

$$x = 0.365^\circ$$

$$x = 1.206^\circ$$




Figure 2

Figure 2 shows a sketch of part of the curve C with equation

$$y = e^{a-3x} - 3e^{-x}, \quad x \in \mathbb{R}$$

where a is a constant and $a > \ln 4$

The curve C has a turning point P and crosses the x -axis at the point Q as shown in Figure 2.

(a) Find, in terms of a , the coordinates of the point P .

(6)

(b) Find, in terms of a , the x coordinate of the point Q .

(3)

(c) Sketch the curve with equation

$$y = |e^{a-3x} - 3e^{-x}|, \quad x \in \mathbb{R}, \quad a > \ln 4$$

Show on your sketch the exact coordinates, in terms of a , of the points at which the curve meets or cuts the coordinate axes.

(3)

$$\frac{dy}{dx} = -3e^{a-3x} + 3e^{-x} = 0 \quad 3e^{-x} = 3e^{a-3x}$$

$$\therefore -x = a - 3x$$

$$2x = a$$

$$x = \frac{1}{2}a$$

$$y = e^{a - \frac{3}{2}a} - 3e^{-\frac{a}{2}} = e^{-\frac{a}{2}} - 3e^{-\frac{a}{2}}$$

$$\left(\frac{1}{2}a, -2e^{-\frac{a}{2}} \right)$$

$$b) \quad y=0 \Rightarrow e^{a-3x} = 3e^{-x}$$


$$\ln e^{a-3x} = \ln 3e^{-x}$$

$$a-3x = \ln 3 + \ln e^{-x}$$

$$a-3x = \ln 3 - x \quad \therefore 2x = a - \ln 3$$

$$x = \frac{1}{2}(a - \ln 3)$$

c)



$$y = |e^{a-3x} - 3e^{-x}|$$

$$x=0 \quad y = e^a - 3$$

12.




Figure 3

Figure 3 shows a sketch of part of the curve C with parametric equations

$$x = \tan t, \quad y = 2\sin^2 t, \quad 0 \leq t < \frac{\pi}{2}$$

The finite region S , shown shaded in Figure 3, is bounded by the curve C , the line $x = \sqrt{3}$ and the x -axis. This shaded region is rotated through 2π radians about the x -axis to form a solid of revolution.

(a) Show that the volume of the solid of revolution formed is given by

$$4\pi \int_0^{\frac{\pi}{3}} (\tan^2 t - \sin^2 t) dt$$

(6)

(b) Hence use integration to find the exact value for this volume.

(6)

$$12a) \text{ Volume} = \pi \int y^2 dx = \pi \int_{x=0}^{x=\sqrt{3}} y^2 \frac{dx}{dt} dt$$

$$\tan t = \sqrt{3} \therefore t = \frac{\pi}{3}$$

$$\tan t = 0 \therefore t = 0$$

$$x = \tan t \quad y^2 = 4 \sin^4 t$$

$$\frac{dx}{dt} = \sec^2 t$$

$$\therefore \text{Volume} = 4\pi \int_0^{\frac{\pi}{3}} \sin^4 t \sec^2 t dt$$

$$= 4\pi \int_0^{\frac{\pi}{3}} \frac{\sin^2 t (1 - \cos^2 t)}{\cos^2 t} dt = 4\pi \int_0^{\frac{\pi}{3}} \tan^2 t (1 - \cos^2 t) dt$$

$$= 4\pi \int_0^{\frac{\pi}{3}} \tan^2 t - \frac{\sin^2 t}{\cos^2 t} dt = 4\pi \int_0^{\frac{\pi}{3}} \tan^2 t - \sin^2 t dt$$

b)

$$\frac{\sin^2 + \cos^2}{\cos^2} = \frac{1}{\cos^2} \Rightarrow \tan^2 + 1 = \sec^2$$

$$-1 + \sec^2 x = \tan^2 x$$

$$\cos 2t = 1 - 2\sin^2 t$$

$$\sin^2 t = \frac{1}{2} - \frac{1}{2} \cos 2t$$

$$= 4\pi \int_0^{\frac{\pi}{3}} \sec^2 x - 1 - \frac{1}{2} + \frac{1}{2} \cos 2t dt$$

$$= 4\pi \int_0^{\frac{\pi}{3}} \sec^2 x + \frac{1}{2} \cos 2t - \frac{3}{2} dt = 2\pi \int_0^{\frac{\pi}{3}} 2\sec^2 x + \cos 2t - 3 dt$$

$$= 2\pi \left[2\tan x + \frac{1}{2} \sin 2t - 3t \right]_0^{\frac{\pi}{3}}$$

$$= 2\pi \left[\left(2\sqrt{3} + \frac{\sqrt{3}}{4} - \pi \right) - (0) \right]$$

$$= 2\pi \left[\frac{9\sqrt{3}}{4} - \pi \right] = \frac{1}{2} \pi (9\sqrt{3} - 4\pi)$$

13. (a) Express $2\sin\theta + \cos\theta$ in the form $R\sin(\theta + \alpha)$, where R and α are constants, $R > 0$ and $0 < \alpha < 90^\circ$. Give your value of α to 2 decimal places. (3)

$$R\sin(\theta + \alpha) = R\sin\theta\cos\alpha + R\cos\theta\sin\alpha$$

$$\frac{R\sin\alpha}{R\cos\alpha} = \frac{1}{2}$$

$$\tan\alpha = \frac{1}{2} \quad \alpha = 0.463647 \quad \alpha = 0.46^\circ \quad R^2 = 1^2 + 2^2 \Rightarrow R = \sqrt{5}$$

$$\sqrt{5}\sin(\theta + 0.46^\circ) \approx \sqrt{5}\sin(\theta + 26.57^\circ)$$




Figure 4

Figure 4 shows the design for a logo that is to be displayed on the side of a large building. The logo consists of three rectangles, C , D and E , each of which is in contact with two horizontal parallel lines l_1 and l_2 . Rectangle D touches rectangles C and E as shown in Figure 4.

Rectangles C , D and E each have length 4 m and width 2 m. The acute angle θ between the line l_2 and the longer edge of each rectangle is shown in Figure 4.

Given that l_1 and l_2 are 4 m apart,

(b) show that

$$2\sin\theta + \cos\theta = 2 \tag{2}$$

Given also that $0 < \theta < 45^\circ$,

(c) solve the equation

$$2\sin\theta + \cos\theta = 2$$


giving the value of θ to 1 decimal place. (3)

Rectangles C and D and rectangles D and E touch for a distance h m as shown in Figure 4.

Using your answer to part (c), or otherwise,

(d) find the value of h , giving your answer to 2 significant figures. (3)

b)



$$\begin{aligned} \therefore 4 \sin \theta + 2 \cos \theta &= 4 \\ \therefore 2 \sin \theta + \cos \theta &= 2 \end{aligned}$$

c) $\sqrt{5} \sin(\theta + 0.469) = 2$

$$\theta + 26.57 = \sin^{-1}\left(\frac{2}{\sqrt{5}}\right) = 63.4, 116.56 \dots$$

$$\therefore \theta = \underline{36.9^\circ}$$

d)



$$\tan \theta = \frac{2}{x} \therefore x = \frac{2}{\tan \theta} = \frac{8}{3}$$

$$\therefore h = 4 - \frac{8}{3} = \frac{4}{3} \text{ m} = \underline{1.3 \text{ m}}$$

14. Relative to a fixed origin O , the line l has vector equation

$$\mathbf{r} = \begin{pmatrix} -1 \\ -4 \\ 6 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 1 \\ -1 \end{pmatrix}$$

where λ is a scalar parameter.

Points A and B lie on the line l , where A has coordinates $(1, a, 5)$ and B has coordinates $(b, -1, 3)$.

(a) Find the value of the constant a and the value of the constant b .

(3)

(b) Find the vector \vec{AB} .

(2)

The point C has coordinates $(4, -3, 2)$

(c) Show that the size of the angle CAB is 30°

(3)

(d) Find the exact area of the triangle CAB , giving your answer in the form $k\sqrt{3}$, where k is a constant to be determined.

(2)

The point D lies on the line l so that the area of the triangle CAD is twice the area of the triangle CAB .

(e) Find the coordinates of the two possible positions of D .

(4)

$$\text{a) } \begin{pmatrix} -1+2\lambda \\ -4+\lambda \\ 6-\lambda \end{pmatrix} = \begin{pmatrix} 1 \\ a \\ 5 \end{pmatrix} \quad \lambda=1 \quad \therefore \underline{a=-3} \quad \begin{pmatrix} -1+2\lambda \\ -4+\lambda \\ 6-\lambda \end{pmatrix} = \begin{pmatrix} b \\ -1 \\ 3 \end{pmatrix} \quad \lambda=3 \quad \therefore \underline{b=5}$$


$$\text{b) } \vec{AB} = b - a = \begin{pmatrix} 5 \\ -1 \\ 3 \end{pmatrix} - \begin{pmatrix} 1 \\ -3 \\ 5 \end{pmatrix} = \begin{pmatrix} 4 \\ 2 \\ -2 \end{pmatrix}$$

$$\text{c) } \vec{AC} = c - a = \begin{pmatrix} 4 \\ -3 \\ 2 \end{pmatrix} - \begin{pmatrix} 1 \\ -3 \\ 5 \end{pmatrix} = \begin{pmatrix} 3 \\ 0 \\ -3 \end{pmatrix}$$

$$\cos \theta = \frac{\begin{pmatrix} 4 \\ 2 \\ -2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 0 \\ -3 \end{pmatrix}}{\left| \begin{pmatrix} 4 \\ 2 \\ -2 \end{pmatrix} \right| \cdot \left| \begin{pmatrix} 3 \\ 0 \\ -3 \end{pmatrix} \right|} \Rightarrow \cos \theta = \frac{18}{\sqrt{24} \times 18} \therefore \cos \theta = \frac{18}{12\sqrt{3}}$$


$$\therefore \cos \theta = \frac{18\sqrt{3}}{36} = \frac{\sqrt{3}}{2} \therefore \underline{\theta = 30^\circ}$$

d)



$$\text{Area} = \frac{1}{2} \sqrt{24} \sqrt{18} \sin 30 = \frac{12\sqrt{3}}{4} = \frac{3\sqrt{3}}{1}$$

e)



$$\text{Area} = 6\sqrt{3}$$

$$\frac{1}{2} \sqrt{24} |\vec{AD}| \sin 30 = 6\sqrt{3}$$

$$\therefore |\vec{AD}| = \frac{24\sqrt{3}}{\sqrt{24}} = 6\sqrt{2}$$

$$AB = \sqrt{18} = 3\sqrt{2}$$

$$\therefore \vec{AD} = 2 \times \vec{AB}$$

$$d = a \pm 2\vec{AB}$$

$$d = \begin{pmatrix} 1 \\ -3 \\ 5 \end{pmatrix} \pm 2 \begin{pmatrix} 4 \\ 2 \\ 2 \end{pmatrix} = \begin{pmatrix} 9 \\ 1 \\ 1 \end{pmatrix} \text{ or } \begin{pmatrix} -7 \\ -7 \\ 9 \end{pmatrix}$$

$$(9, 1, 1) \text{ or } (-7, 7, 9)$$