



Mark Scheme (FINAL)

October 2019

Pearson Edexcel IAL Mathematics (WMA12)

Pure Mathematics P2

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

October 2019

Publications Code WMA12\_01\_1910\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2019

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Pearson Mathematics mark schemes use the following types of marks:
  - **M** marks: Method marks are awarded for ‘knowing a method and attempting to apply it’, unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  or ft will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - d... or dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper or ag- answer given
  - $\square$  or d... The second mark is dependent on gaining the first mark
4. All A marks are ‘correct answer only’ (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
  5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by ‘MR’ in the body of the script.

6. If a candidate makes more than one attempt at any question:
- If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.
8. Marks for each question are scored by clicking in the marking grids that appear below each student response on ePEN. The maximum mark allocation for each question/part question(item) is set out in the marking grid and you should allocate a score of '0' or '1' for each mark, or "trait", as shown:

	0	1
aM		•
aA	•	
bM1		•
bA1	•	
bB	•	
bM2		•
bA2		•

9. Be careful when scoring a response that is either all correct or all incorrect. It is very easy to click down the '0' column when it was meant to be '1' and all correct.

## General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

### Method mark for solving 3 term quadratic:

#### 1. Factorisation

$$(x^2 + bx + c) = (x + p)(x + q), \text{ where } |pq| = |c|, \text{ leading to } x = \dots$$

$$(ax^2 + bx + c) = (mx + p)(nx + q), \text{ where } |pq| = |c| \text{ and } |mn| = |a|, \text{ leading to } x = \dots$$

#### 2. Formula

Attempt to use correct formula (with values for  $a$ ,  $b$  and  $c$ ).

#### 3. Completing the square

$$\text{Solving } x^2 + bx + c = 0: \quad (x \pm \frac{b}{2})^2 \pm q \pm c, \quad q \neq 0, \quad \text{leading to } x = \dots$$

### Method marks for differentiation and integration:

#### 1. Differentiation

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

#### 2. Integration

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

### Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

**Method mark** for quoting a correct formula and attempting to use it, even if there are small mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

### Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

### Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice.



Question Number	Scheme	Marks
<b>1 (a)</b>	$y = 2x^2(x-5) = 2x^3 - 10x^2$ $\frac{dy}{dx} = 6x^2 - 20x$ Sets $\frac{dy}{dx} = 0 \Rightarrow 6x^2 - 20x = 0 \Rightarrow x = 0, \frac{10}{3}$ oe	B1 M1 dM1 A1 <b>(4)</b>
<b>(b)</b>	$x \leq 0, \quad x \geq \frac{10}{3}$	M1 A1 <b>(2)</b> <b>(6 marks)</b>
<b>Alt (a)</b>	$\left( \frac{d(uv)}{dx} = uv' + vu' \right) \quad u = 2x^2, u' = 4x, v = x - 5, v' = 1$ $\left( \frac{dy}{dx} = \right) Ax(x-5) + Bx^2$ Sets $\frac{dy}{dx} = 0 \Rightarrow 4x(x-5) + 2x^2 = 0 \Rightarrow x = 0, \frac{10}{3}$	B1 M1 dM1 A1

(a)

B1 Expands to a correct form ( $y =$ )  $2x^2(x-5) = 2x^3 - 10x^2$

(may be implied by sight of  $\frac{dy}{dx} = 6x^2 - 20x$ )

If a candidate uses the product rule then award this mark for correct terms for  $u, u', v$  and  $v'$  stated or implied.

M1 Differentiated reducing a power by one on one of their terms.

If a candidate uses the product rule then look for  $\left( \frac{dy}{dx} = \right) Ax(x-5) + Bx^2$ . (Do not award for just differentiating the factor in front of the bracket and the  $x-5$ )

dM1  $\frac{dy}{dx} = 0$  and solves a quadratic equation to find at least one solution. (Usual rules for solving a quadratic). Values such as 0 and 3.33 to imply this mark.

It is dependent on the previous method mark. Dividing by  $x$  to find one of the solutions is also acceptable for this mark.

Allow answers to be just written down but you may need to check these on your calculator.



A1  $x = 0, \frac{10}{3}$  oe (Must be seen in (a))

Allow if embedded within pairs of coordinates for the stationary points of C

SC If no calculus work shown then 0001 for achieving the correct  $x$  coordinates.

SC If no credit worthy work seen in (a) but they use calculus in (b) then B1M1M1 can be awarded but withhold the A mark.

(b) Note we are now marking this on open as M1A1

M1 One of  $x \leq 0$  or  $x \geq \frac{10}{3}$ . Allow for  $x < 0$  or  $x > \frac{10}{3}$ .

They must have only achieved a maximum of two  $x$  coordinates in (a).

Do not award for inequalities using solutions from  $y = 0$  or if they have integrated instead.

A1 Both  $x \leq 0, x \geq \frac{10}{3}$ . Allow both  $x < 0, x > \frac{10}{3}$ . Ignore any joining statements such as "and", "or" or any

set notation eg.  $\cup$  or  $\cap$  and allow statements such as  $(-\infty, 0], \left[\frac{10}{3}, \infty\right)$ .  $\frac{10}{3}$  does not need to be in its

lowest terms. Isw after two correct inequality statements unless they contradict. (Eg ignore after two correct inequalities  $x < 0, x > \frac{10}{3}$  a statement such as  $0 > x > \frac{10}{3}$  but withhold the final mark if they

subsequently write  $0 < x < \frac{10}{3}$ )

Question Number	Scheme	Marks
<b>2 (a)</b>	States or uses $r = 1.02$ Attempts $25000 \times 1.02^{13} = 32340$ or 32341 or 32300	B1 M1 A1  <b>(3)</b>
<b>(b)</b>	Attempts $\frac{a(r^n - 1)}{(r - 1)} = \frac{25000(1.02^{14} - 1)}{1.02 - 1}$ or $\frac{125000(1.02^{14} - 1)}{1.02 - 1}$  £1 997 000	M1 A1  A1  <b>(3)</b> <b>(6 marks)</b>

(a)

B1 States or uses  $r = 1.02$ , 102%, (1+2%) oe

M1 Attempts  $25000 \times r^{n2}$  or  $25000 \times r^{n3}$ . Allow eg  $r = 2$  for this mark. They may show the increase each year which is acceptable. Expect to see at least one correct increase for their  $r$  and attempting 12 or 13 increases by the same percentage. Allow a misread if 2500 or 250000 is used instead. Awrt 31706 which is  $25000 \times 1.02^{12}$  is usually sufficient evidence for this mark.

A1 32340 (or 32341). Must be an integer. Allow 32300 which is the answer rounded to 3sf.

(b)

M1 Attempts  $\frac{a(r^n - 1)}{r - 1}$  or  $\frac{a(1 - r^n)}{1 - r}$  with  $a = 25000$  or 125000,  $n = 13$  or 14 and  $r = 1.02$  or their  $r$  from part (a).

A1 For  $\frac{25000(1.02^{14} - 1)}{1.02 - 1}$  or  $\frac{125000(1.02^{14} - 1)}{1.02 - 1}$  or equivalent. Awrt 1 997 000 implies this mark.

A1 1 997 000 only

.....  
Alt(b)

M1 If they attempt to add year by year then look for 13 or 14 values being added with the first year 25000 and at least one increase year to year of 2% or their  $r$ .

A1 All 14 values seen or implied from either column 2 or 3 of the table on the next page. Allow values to be truncated or rounded to 3 significant figures. Awrt 1 997 000 also implies this mark.

A1 1 997 000 only

n	$x1.02^{(n-1)}$	x5
1	25000	125000
2	25500	127500
3	26010	130050
4	26530	132651
5	27061	135304
6	27602	138010
7	28154	140770
8	28717	143586
9	29291	146457
10	29877	149387
11	30475	152374
12	31084	155422
13	31706	158530
14	32340	161701
	399348	1996742

Question Number	Scheme	Marks
<b>3. (a)</b>	$\left(1 + \frac{x}{4}\right)^{12} = 1 + 12\left(\frac{x}{4}\right) + \frac{12 \times 11}{2}\left(\frac{x}{4}\right)^2 + \frac{12 \times 11 \times 10}{3!}\left(\frac{x}{4}\right)^3 + \dots$ $= 1 + 3x + \frac{33}{8}x^2 + \frac{55}{16}x^3$	M1 B1, A1 <b>(3)</b>
<b>(b)</b>	$\left(\frac{x^2 + 8}{x^5}\right)\left(1 + \frac{x}{4}\right)^{12} =$ <p>Sight of a term independent of <math>x = \frac{55}{16}</math> or <math>8 \times {}^{12}C_5 \left(\frac{1}{4}\right)^5 (= \frac{99}{16})</math></p> $\frac{55}{16} + 8 \times {}^{12}C_5 \left(\frac{1}{4}\right)^5 = \frac{55}{16} + \frac{99}{16} = \frac{77}{8}$	B1ft, M1 A1 <b>(3)</b> <b>(6 marks)</b>

(a)

M1 For an attempt at the binomial expansion. Score for a correct attempt at term 3 or 4.

Accept sight of  ${}^{12}C_2 \left(\frac{x}{4}\right)^2$  or  ${}^{12}C_3 \left(\frac{x}{4}\right)^3$  condoning the omission of brackets. Accept any relevant coefficient appearing from Pascal's triangle. FYI  ${}^{12}C_2 = 66$ ,  ${}^{12}C_3 = 220$

B1 For  $1 + 3x$  Must be simplified. Also allow  $1, 3x$  if written as a list.A1 For  $+\frac{33}{8}x^2 + \frac{55}{16}x^3$  Accept  $+4.125x^2 + 3.4375x^3$ 

(All four terms do not need to be written as a summation and may be written as a list).

(b)

B1ft Sight of  $\frac{55}{16}$  or  $8 \times {}^{12}C_5 \left(\frac{1}{4}\right)^5 (= \frac{99}{16})$  which must be independent of  $x$ 

Most candidates do not find the additional term in the expansion of  $\left(1 + \frac{x}{4}\right)^{12}$  so typically would only score a maximum of 100 in (b)

M1 For attempting to add their  $\frac{55}{16}$  to  $8 \times {}^{12}C_5 \left(\frac{1}{4}\right)^5$  or  $8 \times 792 \left(\frac{1}{4}\right)^5$ A1 For  $\frac{77}{8}$  oe eg 9.625



Question Number	Scheme	Marks
4 (a)	-35	B1 (1)
(b)	Attempts $f\left(\pm\frac{2}{3}\right) = 0 \rightarrow \left(\frac{2}{3}-3\right)\left(3\left(\frac{2}{3}\right)^2 + \left(\frac{2}{3}\right) + a\right) - 35 = 0$ $-\frac{7}{3}(2+a) = 35 \Rightarrow a = -17 *$	M1 A1* (2)
(c)	$f(x) = (x-3)(3x^2 + x - 17) - 35 = 3x^3 - 8x^2 - 20x + 16$ $= (3x-2)(x^2 - 2x - 8)$ $= (3x-2)(x+2)(x-4)$	B1 M1 A1 M1 A1 (5) (8 marks)

(a)

B1 -35 stated

(b)

M1 Attempts to set  $f\left(\pm\frac{2}{3}\right) = 0$ . Must see  $\pm\frac{2}{3}$  substituted into the expression and set equal to zero.

Condone bracket errors for this mark. They may multiply out the expression before substituting so allow errors in the manipulation for this mark. (FYI  $3x^3 + x^2 + ax - 9x^2 - 3x - 3a$ )  
“=0” may be implied by their working.

A1\* Achieves the given answer with no incorrect work (including brackets) and at least one correct intermediate line before proceeding to  $a = -17$ . “=0” must be seen somewhere for this mark.Eg.  $-\frac{7}{3}(2+a) = 35 \Rightarrow a = -17$ .....  
Alt (b)M1 Assumes  $a = -17$  and attempts  $f\left(\pm\frac{2}{3}\right) = 0$  where  $f(x) = (x-3)(3x^2 + x - 17) - 35$ A1\* Achieves  $f\left(\frac{2}{3}\right) = 0$  with no incorrect work, at least one correct intermediate line and states hence proven or some other acknowledgement of the “proof”.  
.....

(c)

B1 Multiplies out to achieve  $f(x) = 3x^3 - 8x^2 - 20x + 16$ . This may be implied by eg  $(3x-2)(x^2 - 2x - 8)$

M1 Attempts to divide or factorise out  $(3x-2)$  from their cubic.

By factorisation look for  $3x^3 \pm \dots x^2 \pm \dots x \pm 2\alpha = (3x-2)(x^2 + kx \pm \alpha)$   $k \neq 0$  or they may equate coefficients correctly to find  $k$ .

By division look for 
$$3x-2 \overline{) \begin{array}{r} x^2 \pm \frac{(\beta+2)}{3}x \dots\dots\dots \\ 3x^3 \pm \beta x^2 \dots\dots\dots \\ \underline{3x^3 - 2x^2} \end{array}} \quad \text{(Usually } x^2 \pm 6x \dots)$$

There are various other methods but we should be seeing a correct method to find at least two of the coefficients (allow  $\pm$ ) of their quadratic for the method mark. Send to review if unsure how to mark a response.

A1 Correct factors  $(3x-2)(x^2 - 2x - 8)$  The two factors do not need to be written together eg this mark can be awarded if the quadratic factor is seen at the top of their algebraic division.

M1 Factorises their quadratic factor of the form  $Ax^2 + Bx + C$  with  $A, B, C \neq 0$ . Score for  $(Ax^2 + Bx + C) = (dx + e)(fx + g)$  where  $|A| = |d \times f|$  or  $|C| = |e \times g|$

A1  $(3x-2)(x+2)(x-4)$  on one line following B1M1A1M1

isw  $(3x-2)(x+2)(x-4) \Rightarrow x = \frac{2}{3}, -2, 4$  and allow  $3(x - \frac{2}{3})(x+2)(x-4)$

**Note** the question says using algebra and showing each step.


$f(x) = 3x^3 - 8x^2 - 20x + 16 \Rightarrow (3x-2)(x+2)(x-4)$  is 10000.

Beware  $f(x) = 3x^3 - 8x^2 - 20x + 16 = 0 \Rightarrow x = \frac{2}{3}, -2, 4 \Rightarrow (3x-2)(x+2)(x-4)$  is 10000

$f(x) = 3x^3 - 8x^2 - 20x + 16 = 0 \Rightarrow (x - \frac{2}{3})(x+2)(x-4)$  is 10000

isw  $(3x-2)(x+2)(x-4) \Rightarrow x = \frac{2}{3}, -2, 4$

isw  $3(x - \frac{2}{3})(x+2)(x-4) \Rightarrow (x - \frac{2}{3})(x+2)(x-4)$

Question Number	Scheme	Marks
5.(a)		Shape or intercept at 1 M1 Fully correct A1  h = 0.5 B1  (2)
(b)	$\text{Area} \approx \frac{0.5}{2} \{4.25 + 16.06 + 2 \times (6.427 + 9.125 + 12.34)\}$ $= \text{awrt } 19.0$	M1 A1  (3)
(c)	$\int_2^4 \left( x(x-3) + \left(\frac{1}{2}\right)^x \right) dx = \int_2^4 \left( x^2 + \left(\frac{1}{2}\right)^x - 3x \right) dx = (b) - \left[ \frac{3}{2}x^2 \right]_2^4$ $= \text{awrt } 1.0$	M1 A1ft  (2)
<b>(7 marks)</b>		

(a)

M1 For either the shape in quadrants 1 and 2 only or the  $y$ -intercept at 1. Condone slips of the pen but the gradient should be tending to 0 as  $x$  increases. Condone (1,0) marked on the correct axis.

A1 Fully correct. Shape in quadrants 1 and 2 only with the  $y$ -intercept at 1. Condone slips of the pen towards the  $x$ -axis but the graph should not cross the  $x$ -axis. Condone (1,0) marked on the correct axis. For a fully correct graph its asymptote should not appear to be  $y=1$ . As a rule of thumb look for the asymptote to be at least half way below its  $y$ -intercept.

Do not be too concerned about relative position of the curve in the second quadrant as long as the curve gets steeper as  $x \rightarrow -\infty$

(b)

B1 For  $h = 0.5$  seen or implied by sight of  $\frac{0.5}{2}$  in front of the bracket

M1 For the correct bracket structure  $\{4.25 + 16.06 + 2 \times (6.427 + 9.125 + 12.34)\}$  condoning slips copying from the table or the omission of the final bracket.

eg  $\frac{1}{2} \times 0.5 \times (4.25 + 16.06 + 2 \times (6.427 + 9.125 + 12.34))$

Recovery from missing brackets to achieving the correct answer can still score full marks.

A1 19.0235 but awrt 19.0 will usually score full marks

**Note: The calculator answer for the integral is 18.937... which is A0.**



Examples of mark traits where full marks are not scored:

$$\frac{1}{2} \times 0.5 \times 4.25 + 16.06 + 2 \times (6.427 + 9.125 + 12.34) \quad (= 72.9065) \text{ is B1M0A0}$$

$$\frac{1}{2} \times 0.5 \times 4.25 + 16.06 + 2 \times 6.427 + 9.125 + 12.34 \quad (= 51.4415) \text{ is B1M0A0}$$

(c)

M1 For realising that  $\int_2^4 \left( x(x-3) + \left(\frac{1}{2}\right)^x \right) dx = \int_2^4 \left( x^2 + \left(\frac{1}{2}\right)^x - 3x \right) dx = (b) - \left[ \frac{3}{2}x^2 \right]_2^4$

i.e.  $(b) - \left[ \frac{3}{2}x^2 \right]_2^4$  must be seen or  $(b) - \int_2^4 (3x) dx \Rightarrow (b) - 18$  for this mark. The  $3x$  term must be

isolated from the rest of the integral that is identified in some way to be their part (b). **There must be some integration carried out** so look for  $3x \rightarrow \dots x^2$ . Condone poor notation.

A1ft 1.0235 but accept awrt 1.0 or follow through on their answer or their unrounded answer from (b) - 18 **provided** M1 has been scored.

**Note: The calculator answer for the integral is 0.937 (awrt 0.9) is M0A0**

Question Number	Scheme	Marks
<b>6(a)</b>	Attempts line with gradient $-2$ and point $(4, -1)$ $y+1 = -2(x-4)$ $y = -2x+7$	M1 A1 <b>(2)</b>
<b>(b)</b>	$y = \frac{1}{2}x$ meets $y = -2x+7$ when $\frac{1}{2}x = -2x+7 \Rightarrow x = \frac{14}{5}, y = \frac{7}{5}$ oe Attempts $r^2 = \left(4 - \frac{14}{5}\right)^2 + \left(-1 - \frac{7}{5}\right)^2 = \frac{36}{5}$ oe $(x-4)^2 + (y+1)^2 = \frac{36}{5}$ oe	M1 A1 dM1 A1 A1 <b>(5)</b> <b>(7 marks)</b>

(a)

M1 Attempts the equation of the line with gradient  $-2$  and point  $(4, -1)$   
Condone a slip on one of the signs of the  $(4, -1)$ . For example  $y-1 = -2(x-4)$   
If the form  $y = mx + c$  is used they must proceed as far as  $c = \dots$

A1  $y = -2x + 7$

.....

Alt (a)

M1 Differentiates implicitly the equation of the circle  $(x-4)^2 + (y+1)^2 = r^2 \Rightarrow 2(x-4) + 2(y+1) \frac{dy}{dx} = 0$   
and substitutes in  $\frac{dy}{dx} = \frac{1}{2}$

A1  $y = -2x + 7$

.....

(b)

M1 Attempts to find where  $y = \frac{1}{2}x$  meets their  $y = -2x + 7$ . Expect to see candidates proceed to  $x = \dots$

A1  $\left(\frac{14}{5}, \frac{7}{5}\right)$  or any other form eg  $(2.8, 1.4)$  or  $x = \frac{14}{5}, y = \frac{7}{5}$

dM1 Attempts to find the distance between their  $\left(\frac{14}{5}, \frac{7}{5}\right)$  and  $(4, -1)$  using Pythagoras' theorem. Look for an attempt to subtract the coordinates, square and add. It is dependent on the previous method mark. Condone a slip on one of the signs.

A1 For  $(r =) \sqrt{\frac{36}{5}}$  or  $(r^2 =) \frac{36}{5}$  oe

A1  $(x-4)^2 + (y+1)^2 = \frac{36}{5}$  or  $(x-4)^2 + (y+1)^2 = 7.2$  or  $5(x-4)^2 + 5(y+1)^2 = 36$  only

.....

Alt (b)

M1 Substitutes  $y = \frac{1}{2}x$  into  $(x-4)^2 + (y+1)^2 = r^2$ , multiplies out and rearranges ... = 0

A1  $5x^2 - 28x + 68 - 4r^2 = 0$  or  $5y^2 - 14y + 17 - r^2$

dM1 Uses the discriminant  $b^2 - 4ac = 0$  and proceeds to  $r = \dots$  or  $r^2 = \dots$  It is dependent on the previous method mark.

A1 For  $r = \sqrt{\frac{36}{5}}$  or  $r^2 = \frac{36}{5}$

A1  $(x-4)^2 + (y+1)^2 = \frac{36}{5}$  oe eg  $(x-4)^2 + (y+1)^2 = 7.2$  or  $5(x-4)^2 + 5(y+1)^2 = 36$

**Note: Candidates who substitute  $y = -2x + "7"$  into  $(x-4)^2 + (y+1)^2 = r^2$  and attempt the discriminant approach will score 0 marks in (b)**

Question Number	Scheme	Marks
7. (i)	$\log_a \left( \frac{\sqrt{a}}{b} \right) = \frac{1}{2} \log_a a - \log_a b = \frac{1}{2} - k$	M1 A1 (2)
(ii)	$\frac{\log_a a^2 b}{\log_a b^3} = \frac{2 \log_a a + \log_a b}{3 \log_a b} = \frac{2+k}{3k}$	M1 A1 (2)
(iii)	$\sum_{n=1}^{50} (k + \log_a b^n) = 50k + (1k + 2k + 3k + \dots + 50k) \text{ or } (2k + 3k + 4k + \dots + 51k)$ <p>Uses the sum formula an AP with <math>n = 50, d = k</math></p> $S = 50k + \frac{50}{2}(2k + 49k) \qquad S = \frac{50}{2}(2k + 51k)$ $= 1325k$	M1 A1 A1 (3) (7 marks)

(i)

M1 Uses log laws to achieve  $\frac{1}{2} \log a - \log b$  or eg  $0.5 \log a - \log b$ A1  $\frac{1}{2} - k$  oe

(ii)

M1 Uses correct log laws on the numerator or denominator. Score for  $\frac{\dots}{3 \log b}$  or  $\frac{\dots}{3k}$  or sight of the numerator  $2 \log a + \log b$  which does not have to be part of a fraction (May be implied by  $2+k$ )A1  $\frac{2+k}{3k}$  or  $\frac{2}{3k} + \frac{1}{3}$  do not isw

(iii)

M1 **Either** uses the sum formula for an AP in terms of  $k$  on part or all of the expression with  $n = 50, d = k, a = k$  or  $n = 50, d = k, a = 2k$ . It is sufficient to see the terms substituted in for this mark.

$$\sum_{n=1}^{50} nk = \frac{50}{2}(2k + 49k) \text{ or } \sum_{n=1}^{50} (k + nk) = \frac{50}{2}(2k + 51k).$$

You may see the equivalent sum  $\frac{n}{2}(a + L)$  eg  $\frac{50}{2}(k + 50k)$

**Or** uses the sum formula for an AP in terms of  $\log_a b$  on part of the expression with  $n = 50, d = \log b, a = \log b$

$$\sum_{n=1}^{50} \log b^n = \frac{50}{2}(2 \log b + 49 \log b)$$

You may see the equivalent sum  $\frac{n}{2}(a + L) = \sum_{n=1}^{50} \log b^n = \frac{50}{2}(\log b + 50 \log b)$

A1 A correct unsimplified answer in terms of  $k$ .

Eg.  $S = 50k + \frac{50}{2}(2k + 49k)$  or  $S = \frac{50}{2}(2k + 51k)$

A1  $1325k$

Question Number	Scheme	Marks
8 (i)	$\int \frac{8\sqrt{x}-5}{2x^2} dx = \int 4x^{-\frac{3}{2}} - \frac{5}{2}x^{-2} dx$	B1
	$= -8x^{-\frac{1}{2}} + \frac{5}{2}x^{-1} (+C)$	M1 A1
(ii)	$\int_2^4 \left( 4x^{-\frac{3}{2}} - \frac{5}{2}x^{-2} \right) dx = \left( -4 + \frac{5}{8} \right) - \left( -4\sqrt{2} + \frac{5}{4} \right) = 4\sqrt{2} - \frac{37}{8}$	dM1 A1
	$\int \left( \frac{1}{2}x^2 + k \right) dx = \left[ \frac{1}{6}x^3 + kx \right]$	M1 A1
	$\int_{-3}^6 \left( \frac{1}{2}x^2 + k \right) dx = 55 \Rightarrow \left[ \frac{1}{6}x^3 + kx \right]_{-3}^6 = 55 \Rightarrow (36 + 6k) - \left( -\frac{9}{2} - 3k \right) = 55 \Rightarrow k = \dots$	dM1
	$k = \frac{29}{18}$	A1
		(4) (9 marks)

(i)

B1 For  $\frac{8\sqrt{x}-5}{2x^2} \rightarrow 4x^{-\frac{3}{2}} - \frac{5}{2}x^{-2}$  (seen or implied). Ignore any +C

You may also see a factor of  $\frac{1}{2}$  taken out eg  $\frac{8\sqrt{x}-5}{2x^2} \rightarrow \frac{1}{2} \left( 8x^{-\frac{3}{2}} - 5x^{-2} \right)$

M1 For raising any index by one on one of their terms. Follow through on incorrect indices but do not allow for candidates who attempt to integrate the numerator and denominators separately.

A1 Correct terms (may be unsimplified but the indices must be processed)

dM1 Substitutes 4 and 2 into an integrated function and subtracts either way round. Dependent on the previous method mark. awrt 1.03 following B1M1A1 is sufficient evidence to award this mark.

A1  $4\sqrt{2} - \frac{37}{8}$  or exact equivalent (eg  $4\sqrt{2} - 4.625$  or  $\frac{32\sqrt{2} - 37}{8}$ )

**Note: Answer only scores 0 marks.**

(ii)

M1 Attempts to integrate  $\left( \frac{1}{2}x^2 + k \right)$  with one term correct unsimplified so allow eg  $\frac{1}{2}x^3$  or  $kx^1$  but the index must be processed.

A1  $\frac{1}{6}x^3 + kx$  (allow unsimplified equivalent expressions as above)

dM1 Substitutes 6 and -3 into  $Ax^3 + kx$ , subtracts, sets = 55 and proceeds to  $k = \dots$ . Do not be concerned with the mechanics of their rearrangement.

A1  $k = \frac{29}{18}$  or exact equivalent eg 1.61 but do not allow 1.6 or 1.61 or 1.61....



Question Number	Scheme	Marks
9.(i)	$3 \sin(2\theta - 10^\circ) = 1 \Rightarrow (2\theta - 10^\circ) = \arcsin\left(\frac{1}{3}\right)$ $\theta = \frac{19.47 + 10}{2}, \frac{160.53 + 10}{2}$ $\theta = \text{awrt } 14.7^\circ, 85.3^\circ$	M1 dM1 A1, A1 <b>(4)</b>
(ii) (a)	<p>Writes <math>\frac{1}{\tan \alpha} - \sin \alpha = 2 \sin \alpha - \frac{1}{\tan \alpha}</math> oe</p> $\frac{2}{\tan \alpha} = 3 \sin \alpha \Rightarrow \frac{2 \cos \alpha}{\sin \alpha} = 3 \sin \alpha \Rightarrow 2 \cos \alpha = 3 \sin^2 \alpha *$	M1 dM1 A1* <b>(3)</b>
(b)	$2 \cos \alpha = 3 \sin^2 \alpha \Rightarrow 2 \cos \alpha = 3(1 - \cos^2 \alpha)$ $3 \cos^2 \alpha + 2 \cos \alpha - 3 = 0$ <p>Attempts to solve <math>3 \cos^2 \alpha + 2 \cos \alpha - 3 = 0 \Rightarrow \cos \alpha = \frac{-2 \pm \sqrt{40}}{6}</math> oe</p> $\alpha = 5.517 \text{ radians}$	M1 A1 dM1 A1 A1 <b>(5)</b>
		<b>(12 marks)</b>

(i)

M1 For proceeding to  $x = \arcsin\left(\frac{1}{3}\right)$  which may be implied by the sight of awrt  $19.5^\circ$  or awrt  $160.5^\circ$   
(Allow awrt 0.340 (radians) for this mark)

dM1 For correct order of operations leading to one answer for  $\theta$ . May be implied by  $14.7^\circ/14.8^\circ$  but cannot be scored by adding 10 to an angle in radians but may be implied by awrt 0.257/0.258 rad

A1 One of  $\theta = \text{awrt } 14.7^\circ, 85.3^\circ$  ignore any others.

A1 Both of  $\theta = \text{awrt } 14.7^\circ, 85.3^\circ$  and no others in the range.

**Note that solutions based entirely on graphical or numerical methods are not acceptable so answers only will score 0 marks.**

(ii)(a)

M1 Uses the terms of an AP to set up a correct equation. Eg  $\frac{1}{\tan \alpha} - \sin \alpha = 2 \sin \alpha - \frac{1}{\tan \alpha}$ ,  
 $\frac{\cos \alpha}{\sin \alpha} - \sin \alpha = 2 \sin \alpha - \frac{\cos \alpha}{\sin \alpha}$ . They may also do  $2\left(\frac{1}{\tan \alpha} - \sin \alpha\right) = 2 \sin \alpha - \sin \alpha$ . Condone mixed variables and poor notation for the method marks.



dM1 Uses  $\tan \alpha = \frac{\sin \alpha}{\cos \alpha}$  within an equation involving  $\frac{1}{\tan \alpha}$ ,  $\sin \alpha$  &  $2 \sin \alpha$ . This mark can be implied but do not award if they just proceed straight to the final answer. It is dependent on the previous method mark. A candidate starting with  $\frac{\cos \alpha}{\sin \alpha} - \sin \alpha = 2 \sin \alpha - \frac{\cos \alpha}{\sin \alpha}$  scores M1dM1 straight away.

A1\* Proceeds to given answer with no errors or omissions. The equation must start with an equation involving  $\tan \alpha$  but see the note below for further guidance. Withhold this mark for incorrect notation eg  $\sin \alpha^2$  or if they had mixed variables on the same line. Eg  $\alpha$  and  $\theta$ .

Note: In the example below they can score full marks as  $\tan \alpha = \frac{\sin \alpha}{\cos \alpha}$  is implied by their second term listed on the first line. Had they not written these first three terms or stated somewhere in their solution  $\tan \alpha = \frac{\sin \alpha}{\cos \alpha}$  in their working then we would withhold the final mark.

$\tan \alpha = \frac{\sin \alpha}{\cos \alpha}$  implied here

(ii)(b)

M1 Attempts to use  $\sin^2 \alpha + \cos^2 \alpha = 1$  Eg.  $2 \cos \alpha = 3 \sin^2 \alpha \Rightarrow 2 \cos \alpha = 3(\pm 1 \pm \cos^2 \alpha)$ . Beware that a candidate may use this identity in part (ii)(a) which would not gain the mark in this part.

A1  $3 \cos^2 \alpha + 2 \cos \alpha - 3 = 0$  The “=0” may be implied by later work but the terms must be collected on one side. Evidence of this may be awarded for correct coefficients substituted into the quadratic formula.

dM1 Attempts to solve their  $3 \cos^2 \alpha + 2 \cos \alpha - 3 = 0$  by the formula/completing the square, usual rules for solving a 3TQ apply but do not award for attempted factorisation unless their quadratic factorises.

Award for  $(\cos \alpha =) \frac{-1 \pm \sqrt{10}}{3}$  or  $(\cos \alpha =)$  awrt 0.72 or awrt -1.4 or equivalent. You may need to check decimal values on your calculator.

A1  $(\cos \alpha =) \frac{-2 + \sqrt{40}}{6}$  oe (typically  $(\cos \alpha =) \frac{-1 + \sqrt{10}}{3}$ ). Implied by  $(\cos \alpha =)$  awrt 0.721 or  $(\alpha =)$  awrt 5.52 radians.

A1  $(\alpha =)$  awrt 5.517 radians and no others in the given range

Question Number	Scheme	Marks
<b>10.(a)</b>	$x = 2, y = 5 \Rightarrow 5 = 8a - 12 + 6 + b$ $\frac{dy}{dx} = 3ax^2 - 6x + 3$ AND $x = 2, \frac{dy}{dx} = 7 \Rightarrow 7 = 12a - 12 + 3$ Solves $11 = 8a + b$ and $7 = 12a - 9 \Rightarrow a = \frac{4}{3}, b = \frac{1}{3}$	M1 M1 A1 A1 <b>(4)</b>
<b>(b)</b>	Sets $\frac{dy}{dx} = 3ax^2 - 6x + 3 = 0$ with their value of $a$ and $b$ $4x^2 - 6x + 3 = 0$ and attempts " $b^2 - 4ac$ " $b^2 - 4ac = -12 < 0$ hence there are no turning points oe	M1 dM1 A1* <b>(3)</b> <b>(7 marks)</b>

(a)  
M1 Substitutes  $x = 2, y = 5$  into  $y = ax^3 - 3x^2 + 3x + b$  to get an equation in  $a$  and  $b$  (condone slips)

M1 Substitutes  $x = 2, \frac{dy}{dx} = 7$  into  $\frac{dy}{dx} = 3ax^2 - 6x + 3$  to get an equation in  $a$  ( $\frac{dy}{dx}$  must be correct)

A1  $a = \frac{4}{3}$  or exact equivalent

A1  $b = \frac{1}{3}$  or exact equivalent

(b)  
M1 Sets their  $\frac{dy}{dx} = 0$  with their value of  $a$ . This may be implied by later working.

dM1 Attempts to find  $b^2 - 4ac$  or roots via the formula

A1\* Achieves  $4x^2 - 6x + 3 = 0, b^2 - 4ac = -12$  and states  $-12 < 0$  and so there are **no turning points** or equivalent. They may attempt to solve the equation and either state that **no real roots so no turning points** or achieve complex roots and state that **no real roots so no turning points**. Note full marks can be scored with an incorrect value for  $b = \frac{1}{3}$  but cannot be scored from an incorrect value for  $a = \frac{4}{3}$  from part (a)

.....  
Alt (b)

M1 Attempts to complete the square for  $\frac{dy}{dx} = 3ax^2 - 6x + 3$  for their value of  $a$  to achieve eg  $4(x \pm \dots)^2$  or  $(2x \pm \dots)^2$

dM1  $4x^2 - 6x + 3 = 4\left(x \pm \frac{3}{4}\right)^2 \pm \dots$  or  $4x^2 - 6x + 3 = \left(2x \pm \frac{3}{2}\right)^2 \pm \dots$

A1\* Achieves  $4x^2 - 6x + 3 = 4\left(x - \frac{3}{4}\right)^2 + \frac{3}{4}$  or  $4x^2 - 6x + 3 = \left(2x - \frac{3}{2}\right)^2 + \frac{3}{4}$  and states there are **no turning points** as  $\frac{dy}{dx} > 0$  (for all  $x$ ) or equivalent.

If you see any other ways that may be credit worthy then send to review



