

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel  
Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

**Wednesday 13 May 2020**

Morning (Time: 2 hours)

Paper Reference **8MA0/01**

**Mathematics**  
**Advanced Subsidiary**  
**Paper 1: Pure Mathematics**

**You must have:**

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.  
Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 14 questions in this question paper. The total mark for this paper is 100.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P62645RA

©2020 Pearson Education Ltd.

1/1/1/1/1/



P 6 2 6 4 5 R A 0 1 4 0



Pearson

1. A curve has equation

$$y = 2x^3 - 4x + 5$$

Find the equation of the tangent to the curve at the point  $P(2, 13)$ .

Write your answer in the form  $y = mx + c$ , where  $m$  and  $c$  are integers to be found.

**Solutions relying on calculator technology are not acceptable.**

(5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





2. [In this question the unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are due east and due north respectively.]

A coastguard station  $O$  monitors the movements of a small boat.

At 10:00 the boat is at the point  $(4\mathbf{i} - 2\mathbf{j})$  km relative to  $O$ .

At 12:45 the boat is at the point  $(-3\mathbf{i} - 5\mathbf{j})$  km relative to  $O$ .

The motion of the boat is modelled as that of a particle moving in a straight line at constant speed.

(a) Calculate the bearing on which the boat is moving, giving your answer in degrees to one decimal place.

(3)

(b) Calculate the speed of the boat, giving your answer in  $\text{km h}^{-1}$

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





3.

**In this question you must show all stages of your working.****Solutions relying on calculator technology are not acceptable.**

(i) Solve the equation

$$x\sqrt{2} - \sqrt{18} = x$$

writing the answer as a surd in simplest form.

(3)

(ii) Solve the equation

$$4^{3x-2} = \frac{1}{2\sqrt{2}}$$

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



















7. Given that  $k$  is a positive constant and  $\int_1^k \left( \frac{5}{2\sqrt{x}} + 3 \right) dx = 4$

(a) show that  $3k + 5\sqrt{k} - 12 = 0$

(4)

(b) Hence, using algebra, find any values of  $k$  such that

$$\int_1^k \left( \frac{5}{2\sqrt{x}} + 3 \right) dx = 4$$

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





8. The temperature,  $\theta^\circ\text{C}$ , of a cup of tea  $t$  minutes after it was placed on a table in a room, is modelled by the equation

$$\theta = 18 + 65e^{-\frac{t}{8}} \quad t \geq 0$$

Find, according to the model,

- (a) the temperature of the cup of tea when it was placed on the table, (1)
- (b) the value of  $t$ , to one decimal place, when the temperature of the cup of tea was  $35^\circ\text{C}$ . (3)
- (c) Explain why, according to this model, the temperature of the cup of tea could not fall to  $15^\circ\text{C}$ . (1)




Figure 2

The temperature,  $\mu^\circ\text{C}$ , of a second cup of tea  $t$  minutes after it was placed on a table in a different room, is modelled by the equation

$$\mu = A + Be^{-\frac{t}{8}} \quad t \geq 0$$

where  $A$  and  $B$  are constants.

Figure 2 shows a sketch of  $\mu$  against  $t$  with two data points that lie on the curve.

The line  $l$ , also shown on Figure 2, is the asymptote to the curve.

Using the equation of this model and the information given in Figure 2

- (d) find an equation for the asymptote  $l$ . (4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA











9.




Figure 3

Figure 3 shows part of the curve with equation  $y = 3 \cos x^\circ$ .

The point  $P(c, d)$  is a minimum point on the curve with  $c$  being the smallest negative value of  $x$  at which a minimum occurs.

(a) State the value of  $c$  and the value of  $d$ . (1)

(b) State the coordinates of the point to which  $P$  is mapped by the transformation which transforms the curve with equation  $y = 3 \cos x^\circ$  to the curve with equation

(i)  $y = 3 \cos \left( \frac{x^\circ}{4} \right)$

(ii)  $y = 3 \cos (x - 36)^\circ$  (2)

(c) Solve, for  $450^\circ \leq \theta < 720^\circ$ ,

$$3 \cos \theta = 8 \tan \theta$$

giving your solution to one decimal place.

**In part (c) you must show all stages of your working.**

**Solutions relying entirely on calculator technology are not acceptable.** (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



















11. (i) A circle  $C_1$  has equation

$$x^2 + y^2 + 18x - 2y + 30 = 0$$

The line  $l$  is the tangent to  $C_1$  at the point  $P(-5, 7)$ .

Find an equation of  $l$  in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers to be found.

(5)

(ii) A different circle  $C_2$  has equation

$$x^2 + y^2 - 8x + 12y + k = 0$$

where  $k$  is a constant.

Given that  $C_2$  lies entirely in the 4th quadrant, find the range of possible values for  $k$ .

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA























14. A curve has equation  $y = g(x)$ .

Given that

- $g(x)$  is a cubic expression in which the coefficient of  $x^3$  is equal to the coefficient of  $x$
- the curve with equation  $y = g(x)$  passes through the origin
- the curve with equation  $y = g(x)$  has a stationary point at  $(2, 9)$

(a) find  $g(x)$ ,

(7)

(b) prove that the stationary point at  $(2, 9)$  is a maximum.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





