



Oxford Cambridge and RSA

AS Level Further Mathematics A

Y531/01 Pure Core

Monday 14 May 2018 – Afternoon
Time allowed: 1 hour 15 minutes



You must have:

- Printed Answer Booklet
- Formulae AS Level Further Mathematics A

You may use:

- a scientific or graphical calculator

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Complete the boxes provided on the Printed Answer Booklet with your name, centre number and candidate number.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- Do **not** write in the barcodes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- **You are reminded of the need for clear presentation in your answers.**
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **4** pages.

Answer **all** the questions.

1 (i) Find a vector which is perpendicular to both $\begin{pmatrix} 1 \\ 3 \\ -2 \end{pmatrix}$ and $\begin{pmatrix} -3 \\ -6 \\ 4 \end{pmatrix}$. [2]

(ii) The cartesian equation of a line is $\frac{x}{2} = y - 3 = 2z + 4$.

Express the equation of this line in vector form. [3]

2 **In this question you must show detailed reasoning.**

The cubic equation $2x^3 + 3x^2 - 5x + 4 = 0$ has roots α , β and γ . By making an appropriate substitution, or otherwise, find a cubic equation with integer coefficients whose roots are $\frac{1}{\alpha}$, $\frac{1}{\beta}$ and $\frac{1}{\gamma}$. [3]

3 **In this question you must show detailed reasoning.**

The complex numbers z_1 and z_2 are given by $z_1 = 2 - 3i$ and $z_2 = a + 4i$ where a is a real number.

(i) Express z_1 in modulus-argument form, giving the modulus in exact form and the argument correct to 3 significant figures. [3]

(ii) Find $z_1 z_2$ in terms of a , writing your answer in the form $c + id$. [2]

(iii) The real and imaginary parts of a complex number on an Argand diagram are x and y respectively. Given that the point representing $z_1 z_2$ lies on the line $y = x$, find the value of a . [2]

(iv) Given instead that $z_1 z_2 = (z_1 z_2)^*$ find the value of a . [2]

4 The matrix \mathbf{A} is given by $\mathbf{A} = \begin{pmatrix} 2 & 1 & 2 \\ 1 & -1 & 1 \\ 2 & 2 & a \end{pmatrix}$.

(i) Show that $\det \mathbf{A} = 6 - 3a$. [2]

(ii) State the value of a for which \mathbf{A} is singular. [1]

(iii) Given that \mathbf{A} is non-singular find \mathbf{A}^{-1} in terms of a . [4]

5 In this question you must show detailed reasoning.

(i) Express $(2 + 3i)^3$ in the form $a + ib$. [3]

(ii) Hence verify that $2 + 3i$ is a root of the equation $3z^3 - 8z^2 + 23z + 52 = 0$. [3]

(iii) Express $3z^3 - 8z^2 + 23z + 52$ as the product of a linear factor and a quadratic factor with real coefficients. [4]

6 The matrices **A** and **B** are given by $\mathbf{A} = \begin{pmatrix} t & 6 \\ t & -2 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 2t & 4 \\ t & -2 \end{pmatrix}$ where t is a constant.

(i) Show that $|\mathbf{A}| = |\mathbf{B}|$. [2]

(ii) Verify that $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$. [3]

(iii) Given that $|\mathbf{AB}| = -1$ explain what this means about the constant t . [2]

7 Prove by induction that $2^{n+1} + 5 \times 9^n$ is divisible by 7 for all integers $n \geq 1$. [6]

8 The 2×2 matrix **A** represents a transformation T which has the following properties.

- The image of the point $(0, 1)$ is the point $(3, 4)$.
- An object shape whose area is 7 is transformed to an image shape whose area is 35.
- T has a line of invariant points.

(i) Find a possible matrix for **A**. [8]

The transformation S is represented by the matrix **B** where $\mathbf{B} = \begin{pmatrix} 3 & 1 \\ 2 & 2 \end{pmatrix}$.

(ii) Find the equation of the line of invariant points of S. [2]

(iii) Show that any line of the form $y = x + c$ is an invariant line of S. [3]

END OF QUESTION PAPER

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.