

**OCR**

Oxford Cambridge and RSA

**Thursday 13 June 2019 – Afternoon****AS Level Further Mathematics B (MEI)****Y415/01 Mechanics b****Time allowed: 1 hour 15 minutes****You must have:**

- Printed Answer Booklet
- Formulae Further Mathematics B (MEI)

**You may use:**

- a scientific or graphical calculator

MODEL  
ANSWERS

**INSTRUCTIONS**

- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- **Write your answer to each question in the space provided in the Printed Answer Booklet.** If additional space is required, you should use the lined page(s) at the end of the Printed Answer Booklet. The question number(s) must be clearly shown.
- You are permitted to use a scientific or graphical calculator in this paper.
- Final answers should be given to a degree of accuracy appropriate to the context.
- The acceleration due to gravity is denoted by  $g \text{ m s}^{-2}$ . Unless otherwise instructed, when a numerical value is needed, use  $g = 9.8$ .

**INFORMATION**

- The total number of marks for this paper is **60**.
- The marks for each question are shown in brackets [ ].
- You are advised that an answer may receive **no marks** unless you show sufficient detail of the working to indicate that a correct method is used. You should communicate your method with correct reasoning.
- The Printed Answer Booklet consists of **12** pages. The Question Paper consists of **8** pages.

- 1 A small object of mass 5 kg is attached to one end of each of two identical parallel light elastic strings. The upper ends of both strings are attached to a horizontal ceiling. The object hangs in equilibrium at R, with the extension of each string being 0.1 m, as shown in Fig. 1.




Fig. 1

- (a) Find the stiffness of each string. [3]

One of the strings is now removed and the object initially falls downwards. The object does not return to R at any point in the subsequent motion.

- (b) Suggest a reason why the object does not return to R. [1]

a)  $T = kx$   
*stiffness* *total tension*  
 for equilibrium:  $2k \times 0.1 = 5g$

$$\Rightarrow 2k = 50g$$

$$k = 25g \text{ Nm}^{-1}$$

*writing units is a good habit to get into*

$$= 245 \text{ Nm}^{-1}$$

b) the string may have stretched beyond its elastic limit ('new' unextended length), or the mass may have lost energy in a non-elastic collision

- 2 A particle P of mass  $m$  travels in a straight line on a smooth horizontal surface. At time  $t$ , P is a distance  $x$  from a fixed point O and is moving with speed  $v$  away from O. A horizontal force of magnitude  $3mt$  acts on P, in a direction away from O.
- (a) Show that  $\frac{d^2x}{dt^2} = 3t$ . [1]
- (b) Verify that the general solution of this differential equation is  $x = \frac{1}{2}t^3 + At + k$ , where  $A$  and  $k$  are constants. [2]
- (c) Given that  $x = 6$  and  $v = 12$  when  $t = 1$ , find the values of  $A$  and  $k$ . [4]

a) Newton's 2<sup>nd</sup> law for constant mass:  $F = ma$

$$\text{so } 3mt = ma = m \frac{d^2x}{dt^2} \Rightarrow \frac{d^2x}{dt^2} = 3t$$

b) differentiate given solution:  $\dot{x} = \frac{3}{2}t^2 + A$

$$\ddot{x} = 3t \Rightarrow \text{consistent}$$

c)  $x = 6, v = 12, t = 1: 6 = \frac{1}{2}(1) + A(1) + k \leftarrow x$

$$12 = \frac{3}{2}(1) + A \leftarrow v$$

$$\therefore \frac{21}{2} = A$$

$$k = 6 - \frac{1}{2} - \frac{21}{2}$$

$$k = -5$$

- 3 A particle Q of mass  $m$  moves in a horizontal plane under the action of a single force  $\mathbf{F}$ .

At time  $t$ , Q has velocity  $\begin{pmatrix} 2 \\ 3t-2 \end{pmatrix}$ .

- (a) Find an expression for  $\mathbf{F}$  in terms of  $m$ . [2]

At time  $t$ , the displacement of Q is given by  $\mathbf{r} = \begin{pmatrix} x \\ y \end{pmatrix}$ . When  $t = 1$ , Q is at the point with position vector  $\begin{pmatrix} 4 \\ -4 \end{pmatrix}$ .

- (b) Find the equation of the path of Q, giving your answer in the form  $y = ax^2 + bx + c$ , where  $a$ ,  $b$  and  $c$  are constants to be determined. [7]

- (c) What can you deduce about the path of Q from the value of the constant  $c$  you found in part (b)? [1]

$$a) \underline{v} = \begin{pmatrix} 2 \\ 3t-2 \end{pmatrix} \Rightarrow \underline{a} = \frac{d\underline{v}}{dt} = \begin{pmatrix} 0 \\ 3 \end{pmatrix} \Rightarrow \underline{F} = m\underline{a} = m \begin{pmatrix} 0 \\ 3 \end{pmatrix}$$

$$b) \underline{r} = \begin{pmatrix} x \\ y \end{pmatrix} = \int \underline{v} dt \Rightarrow \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2t+c \\ \frac{3}{2}t^2-2t+d \end{pmatrix} \begin{matrix} \swarrow \\ \nwarrow \end{matrix} \begin{matrix} \text{different} \\ \text{constants} \end{matrix}$$

$$t=1: \underline{r} = \begin{pmatrix} 4 \\ -4 \end{pmatrix} \Rightarrow 2(1)+c=4 \text{ so } c=2$$

$$\frac{3}{2}(1)-2(1)+d=-4 \text{ so } d=-\frac{7}{2}$$

$$\cdot x=2t+2 \text{ \& } y=\frac{3}{2}t^2-2t-\frac{7}{2}$$

rearrange to eliminate  $t$ :

$$t = \frac{1}{2}x - 1$$

$$\text{sub in expression for } y: y = \frac{3}{2} \left( \frac{1}{2}x - 1 \right)^2 - 2 \left( \frac{1}{2}x - 1 \right) - \frac{7}{2}$$

$$= \frac{3}{2} \left( \frac{1}{4}x^2 - x + 1 \right) - x + 2 - \frac{7}{2}$$

$$= \frac{3}{8}x^2 - \frac{3}{2}x + \frac{3}{2} - x - \frac{3}{2}$$

$$y = \frac{3}{8}x^2 - \frac{5}{2}x$$

c)  $C=0$  so particle passes through the origin  
(NOT starts @ origin)

- 4 Two uniform discs, A of mass  $0.2\text{ kg}$  and B of mass  $0.5\text{ kg}$ , collide with smooth contact while moving on a smooth horizontal surface. Immediately before the collision, A is moving with speed  $0.5\text{ ms}^{-1}$  at an angle  $\alpha$  with the line of centres, where  $\sin \alpha = 0.6$ , and B is moving with speed  $0.3\text{ ms}^{-1}$  at right angles to the line of centres. A straight smooth vertical wall is situated to the right of B, perpendicular to the line of centres, as shown in Fig. 4. The coefficient of restitution between A and B is  $0.75$ .





Fig. 4

- (a) Find the speeds of A and B immediately after the collision. [8]
- (b) Explain why there could be a second collision between A and B if B rebounds from the wall with sufficient speed. [1]
- (c) Find the range of values of the coefficient of restitution between B and the wall for which there will be a second collision between A and B. [3]
- (d) How does your answer to part (b) change if the contact between B and the wall is not smooth? [1]

a) conservation of momentum in  $x$  direction:



$$0.2 \times 0.5 \cos \alpha + 0 = 0.2 v_a + 0.5 v_b$$

$$\text{coefficient of restitution: } -e = \frac{v_b - v_a}{u_b - u_a}$$

$$\text{so } v_b - v_a = -0.75(0 - 0.5 \cos \alpha)$$

$$\cos \alpha = \sqrt{1 - \sin^2 \alpha} = \sqrt{1 - 0.6^2} = 0.8 \Rightarrow v_b - v_a = 0.3 \quad \textcircled{1}$$

$$0.2 v_a + 0.5 v_b = 0.08 \quad \textcircled{2}$$

Solve simultaneous equations:

$$5 \times \textcircled{2}: v_a + 2.5v_b = 0.4$$

$$\textcircled{1} + 5 \times \textcircled{2}: 3.5v_b = 0.7$$

$$v_b = 0.2$$

$$\text{sub in: } 0.2 - v_a = 0.3$$

$$v_a = -0.1$$

parallel to x

Momentum only transferred in x-direction, so after collision,

Velocities in y direction are:


$$v_A = 0.5 \sin \alpha = 0.3, v_B = 0.3$$

$$\therefore \text{speed of A} = \sqrt{0.3^2 + 0.1^2} = \sqrt{0.1} \approx 0.316 \text{ ms}^{-1}$$

$$\text{" " B} = \sqrt{0.3^2 + 0.2^2} = \sqrt{0.13} \approx 0.361 \text{ ms}^{-1}$$

b) they could collide again  $\because$  the velocity perpendicular to the line of centres is 0.3 for both A & B, so they

will remain in line over time



c) to 'catch up' with A, after hitting wall

B's speed must  $> 0.1 \text{ ms}^{-1}$  towards A

$$e = \frac{v_B}{u_B} \left. \begin{array}{l} \text{before \&} \\ \text{after colliding} \\ \text{with wall} \end{array} \right\} \Rightarrow v_B > 0.1 \text{ so } e > \frac{0.1}{0.2} \therefore e > 0.5$$

d) B's speed along the wall wouldn't be conserved, so

wouldn't be the same as A's,  $\therefore$  no 2<sup>nd</sup> collision

- 5 Fig. 5 shows the curve with equation  $y = -x^2 + 4x + 2$ .  
 The curve intersects the  $x$ -axis at P and Q. The region bounded by the curve, the  $x$ -axis, the  $y$ -axis and the line  $x = 4$  is occupied by a uniform lamina L. The horizontal base of L is OA, where A is the point (4, 0).




Fig. 5

- (a) (i) Explain why the centre of mass of L lies on the line  $x = 2$ . [1]

(ii) In this question you must show detailed reasoning.

Find the  $y$ -coordinate of the centre of mass of L. [7]

- (b) L is freely suspended from A. Find the angle AO makes with the vertical. [2]

The region bounded by the curve and the  $x$ -axis is now occupied by a uniform lamina M. The horizontal base of M is PQ.

- (c) Explain how the position of the centre of mass of M differs from the position of the centre of mass of L. [2]

a) i. the lamina's curve is symmetrical about  $x=2$ ,

so L is, too

$$\text{ii. } \bar{y} = \frac{\int_a^b \frac{1}{2} y^2 dx}{\int_a^b y dx}, \quad M \bar{y} = \int_a^b \frac{1}{2} \rho y^2 dx, \quad M = \int_a^b \rho y dx$$

$$\begin{aligned} \text{area} &= \int_0^4 (-x^2 + 4x + 2) dx = \left[ -\frac{1}{3}x^3 + 2x^2 + 2x \right]_0^4 \\ &= \frac{56}{3} \end{aligned}$$

$$\frac{1}{2} y^2 \text{ integral: } \frac{1}{2} \int_0^4 (-x^2 + 4x + 2)^2 dx = \frac{1}{2} \int_0^4 (x^4 - 8x^3 + 12x^2 + 16x + 4) dx$$



$$= \frac{1}{2} \left[ \frac{1}{5}x^5 - 2x^4 + 4x^3 + 8x^2 + 4x \right]_0^4$$


$$= \frac{232}{5}$$

$$\Rightarrow \bar{y} = \frac{232}{5} \div \frac{56}{3}$$

$$= \frac{87}{35}$$

$$= 2.49$$

b) freely suspended  $\Rightarrow$  C.O.M. directly below point of suspension



C.O.M. (2, 2.49)

symmetric about  $x=2 \therefore \angle OAG = \angle AOG$  (isosceles)

$$\tan OAG = \frac{2.49}{2} \therefore \angle OAG = \text{angle to vertical} = 51.2^\circ$$

c)  $\bar{x}$  remains the same (same symmetry)

$\bar{y}$  will decrease  $\because$  part of M not in L (added mass)

is closer to the x-axis, adding weight below 2.49

- 6 A smooth solid hemisphere of radius  $a$  is fixed with its plane face in contact with a horizontal surface. The highest point on the hemisphere is H, and the centre of its base is O. A particle of mass  $m$  is held at a point S on the surface of the hemisphere such that angle HOS is  $30^\circ$ , as shown in Fig. 6. The particle is projected from S with speed  $0.8\sqrt{ag}$  along the surface of the hemisphere towards H.




Fig. 6

- (a) Show that the particle passes through H without leaving the surface of the hemisphere. [4]

After passing through H, the particle passes through a point Q on the surface of the hemisphere, where angle HOQ =  $\theta^\circ$ .

- (b) State, in terms of  $g$  and  $\theta$ , the tangential component of the acceleration of the particle when it is at Q. [1]


The particle loses contact with the hemisphere at Q and subsequently lands on the horizontal surface at a point L.

- (c) Find the value of  $\cos \theta$  correct to 3 significant figures. [4]  
 (d) Show that  $OL = ka$ , where  $k$  is to be found correct to 3 significant figures. [5]

END OF QUESTION PAPER

a) to remain on hemisphere,  $R > 0$ . particle must have enough energy to reach H. *centripetal*

@ S, F towards centre =  $\frac{mV^2}{r} = \frac{m(0.8\sqrt{ag})^2}{a} = 0.64mg$



$W \text{ towards centre} = mg \cos 30^\circ$   
 $= 0.866mg$


$$R = W_{50} - F_{50}$$

$0.866mg > 0.64mg \Rightarrow R > 0$ , P stays on hemisphere

Energy @ S = K.E. + G.P.E. =  $\frac{1}{2}m(0.64ag) + mga \cos 30^\circ$

E @ S is > G.P.E. @ H,  $mga$ , so P reaches

H & passes through H



b)  $g \sin \theta$  'state' implies you shouldn't need to do any calculations



c) loses contact  $\Rightarrow R = 0$

@ Q,  $mg \cos \theta = \frac{mv^2}{a}$  ①

E conserved:  $\frac{1}{2}m(0.64ag) + mga \cos 30^\circ = \frac{1}{2}mv^2 + mg a \cos \theta$

sub in ①:  $0.32amg + mga \cos 30^\circ = a \underline{mg \cos \theta} + mg a \cos \theta$

$\Rightarrow \underline{\cos \theta = \frac{1}{3}(0.64 + 2 \cos 30^\circ)}$

keep in exact form for as long as possible

d)  $v^2 = 7.748664a$  using  $v^2 = ag \cos \theta$

vertical motion (SUVAT):  $a \cos \theta = 2.7836\sqrt{a} \sin \theta t + \frac{1}{2}gt^2$

from quadratic formula &  $\sin \theta = \sqrt{1 - \cos^2 \theta} = \sqrt{1 - 0.791^2} = 0.611$

$t = 0.2638\sqrt{a}$

OL = OQ +  $v \cos \theta t = a \sin \theta + 2.7836\sqrt{a} \cos \theta (0.2638\sqrt{a})$

=  $1.19a$

