



# Monday 4 October 2021 – Afternoon

# A Level Further Mathematics B (MEI)

Y420/01 Core Pure

Time allowed: 2 hours 40 minutes

#### You must have:

- the Printed Answer Booklet
- the Formulae Booklet for Further Mathematics B (MEI)
- a scientific or graphical calculator

#### **INSTRUCTIONS**

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the Printed Answer
  Booklet. If you need extra space use the lined pages at the end of the Printed Answer
  Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

### **INFORMATION**

- The total mark for this paper is 144.
- The marks for each question are shown in brackets [ ].
- This document has 8 pages.

# **ADVICE**

Read each question carefully before you start your answer.

# Section A (31 marks)

Answer all the questions.

1 (a) Express 
$$\frac{1}{(2r-1)(2r+1)}$$
 in partial fractions. [3]

(b) Hence find 
$$\sum_{r=1}^{n} \frac{1}{(2r-1)(2r+1)}$$
, expressing the result as a single fraction. [4]

# 2 In this question you must show detailed reasoning.

Find the gradient of the curve  $y = 6 \arcsin(2x)$  at the point with x-coordinate  $\frac{1}{4}$ . Express the result in the form  $m\sqrt{n}$ , where m and n are integers.

## 3 In this question you must show detailed reasoning.

The complex numbers  $z_1$  and  $z_2$  are given by  $z_1 = -2 + 2i$  and  $z_2 = 2\left(\cos\frac{1}{6}\pi + i\sin\frac{1}{6}\pi\right)$ .

(a) Find the modulus and argument of 
$$z_1$$
. [2]

(b) Hence express 
$$\frac{z_1}{z_2}$$
 in exact modulus-argument form. [4]

# 4 In this question you must show detailed reasoning.

Determine the mean value of  $\frac{1}{1+4x^2}$  between x=-1 and x=1. Give your answer to 3 significant figures. [4]

5 (a) Use a Maclaurin series to find a quadratic approximation for 
$$ln(1+2x)$$
. [1]

- (b) Find the percentage error in using the approximation in part (a) to calculate ln(1.2). [3]
- (c) Jane uses the Maclaurin series in part (a) to try to calculate an approximation for ln 3.

Explain whether her method is valid. [2]

6 Given that y = mx is an invariant line of the transformation with matrix  $\begin{pmatrix} 1 & 2 \\ 2 & -2 \end{pmatrix}$ , determine the possible values of m.


# Section B (113 marks)

Answer all the questions.

7 Prove that 
$$\sum_{r=1}^{n} \frac{r}{2^{r-1}} = 4 - \frac{n+2}{2^{n-1}}$$
 for all  $n \ge 1$ .

- 8 The equation  $4x^4 4x^3 + px^2 + qx 9 = 0$ , where p and q are constants, has roots  $\alpha$ ,  $-\alpha$ ,  $\beta$  and  $\frac{1}{\beta}$ .
  - (a) Determine the exact roots of the equation. [5]
  - (b) Determine the values of p and q. [4]
- **9** The transformation T of the plane has associated matrix **M**, where  $\mathbf{M} = \begin{pmatrix} -1 & 0 \\ -2 & 1 \end{pmatrix}$ .
  - (a) On the grid in the Printed Answer Booklet, plot the image OA'B'C' of the unit square OABC under the transformation T. [2]
  - (b) (i) Calculate the value of det M. [1]
    - (ii) Explain the significance of the value of det M in relation to the image OA'B'C'. [2]
  - (c) T is equivalent to a sequence of two transformations of the plane.
    - (i) Specify fully **two** transformations equivalent to T. [3]
    - (ii) Use matrices to verify your answer. [3]
- 10 (a) Show on an Argand diagram the points representing the three cube roots of unity. [2]
  - (b) (i) Find the exact roots of the equation  $z^3 1 = \sqrt{3}i$ , expressing them in the form  $re^{i\theta}$ , where r > 0 and  $-\pi < \theta < \pi$ . [5]
    - (ii) The points representing the cube roots of unity form a triangle  $\Delta_1$ . The points representing the roots of the equation  $z^3 1 = \sqrt{3}i$  form a triangle  $\Delta_2$ .
      - State a sequence of two transformations that maps  $\Delta_1$  onto  $\Delta_2$ . [2]
    - (iii) The three roots in part (b)(i) are  $z_1$ ,  $z_2$  and  $z_3$ .
      - By simplifying  $z_1 + z_2 + z_3$ , verify that the sum of these roots is zero. [2]
    - (iv) Hence show that  $\sin 20^\circ + \sin 140^\circ = \sin 100^\circ$ . [2]

- 11 (a) Given that  $\mathbf{u} = \lambda \mathbf{i} + \mathbf{j} 3\mathbf{k}$  and  $\mathbf{v} = \mathbf{i} + 2\mathbf{j} 2\mathbf{k}$ , find the following, giving your answers in terms of  $\lambda$ .
  - (i) u.v [1]
  - (ii)  $\mathbf{u} \times \mathbf{v}$  [2]
  - (b) Hence determine
    - (i) the acute angle between the planes 2x+y-3z=10 and x+2y-2z=10, [3]
    - (ii) the shortest distance between the lines  $\frac{x-3}{3} = \frac{y}{1} = \frac{z-2}{-3}$  and  $\frac{x}{1} = \frac{y-4}{2} = \frac{z+2}{-2}$ , giving your answer as a multiple of  $\sqrt{2}$ .
- **12 Fig. 12** shows a rhombus OACB in an Argand diagram. The points A and B represent the complex numbers *z* and *w* respectively.



**Fig. 12** 

Prove that  $arg(z+w) = \frac{1}{2}(arg z + arg w)$ .

[A copy of **Fig. 12** is provided in the Printed Answer Booklet.] [4]

13 Find the general solution of the differential equation 
$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - 3y = 2e^x.$$
 [7]

- 14 A curve has polar equation  $r = a(\cos\theta + 2\sin\theta)$ , where a is a positive constant and  $0 \le \theta \le \pi$ .
  - (a) Determine the polar coordinates of the point on the curve which is furthest from the pole. [7]
  - (b) (i) Show that the curve is a circle whose radius should be specified. [6]
    - (ii) Write down the polar coordinates of the centre of the circle. [1]
- 15 The equations of three planes are

$$-4x + ky + 7z = 4,$$
  
$$x - 2y + 5z = l,$$

$$2x + 3y + z = 2$$
.

Given that the planes form a sheaf, determine the values of k and l. [6]

16 (a) Show using exponentials that  $\cosh 2u = 1 + 2 \sinh^2 u$ . [4]

**(b)** Show that 
$$\int_0^2 \frac{x^2}{\sqrt{4+x^2}} \, dx = 2\sqrt{2} - 2\ln(1+\sqrt{2}).$$
 [10]

- 17 In a chemical process, a vessel contains 1 litre of pure water. A liquid chemical is then passed into the top of the vessel at a constant rate of *a* litres per minute and thoroughly mixed with the water. At the same time, the resulting mixture is drawn from the bottom of the vessel at a constant rate of *b* litres per minute. You may assume that the chemical mixes instantly and uniformly with the water. After *t* minutes, the mixture in the vessel contains *x* litres of the chemical.
  - (a) (i) Show that the proportion of chemical present in the vessel after t minutes is  $\frac{x}{1+(a-b)t}.$  [2]

(ii) Hence show that 
$$\frac{dx}{dt} + \frac{bx}{1 + (a-b)t} = a$$
. [2]

- **(b)** First, consider the case where b = a.
  - (i) Solve the differential equation to find x in terms of a and t. [4]
  - (ii) Given that after 1 minute the vessel contains equal amounts of water and chemical, find the rate of inflow of chemical. [2]
- (c) Now consider the case where b = 2a.
  - (i) Explain why the differential equation in part (a)(ii) is now invalid for  $t \ge \frac{1}{a}$ . [1]
  - (ii) Find the maximum amount of chemical in the vessel. [9]

#### **END OF QUESTION PAPER**

6

# **BLANK PAGE**

7 BLANK PAGE



#### Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.