



Oxford Cambridge and RSA

**Thursday 18 May 2023 – Afternoon**

**AS Level Mathematics B (MEI)**

**H630/01 Pure Mathematics and Mechanics**

**Time allowed: 1 hour 30 minutes**



**You must have:**

- the Printed Answer Booklet
- a scientific or graphical calculator



**INSTRUCTIONS**

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the **Printed Answer Booklet**. If you need extra space use the lined pages at the end of the Printed Answer Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give your final answers to a degree of accuracy that is appropriate to the context.
- The acceleration due to gravity is denoted by  $g \text{ m s}^{-2}$ . When a numerical value is needed use  $g = 9.8$  unless a different value is specified in the question.
- Do **not** send this Question Paper for marking. Keep it in the centre or recycle it.

**INFORMATION**

- The total mark for this paper is **70**.
- The marks for each question are shown in brackets [ ].
- This document has **8** pages.

**ADVICE**

- Read each question carefully before you start your answer.

## Formulae AS Level Mathematics B (MEI) (H630)

### Binomial series

$$(a+b)^n = a^n + {}^n C_1 a^{n-1} b + {}^n C_2 a^{n-2} b^2 + \dots + {}^n C_r a^{n-r} b^r + \dots + b^n \quad (n \in \mathbb{N}),$$

$$\text{where } {}^n C_r = {}_n C_r = \binom{n}{r} = \frac{n!}{r!(n-r)!}$$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!} x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!} x^r + \dots \quad (|x| < 1, n \in \mathbb{R})$$

### Differentiation from first principles

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

### Sample variance

$$s^2 = \frac{1}{n-1} S_{xx} \text{ where } S_{xx} = \sum (x_i - \bar{x})^2 = \sum x_i^2 - \frac{(\sum x_i)^2}{n} = \sum x_i^2 - n\bar{x}^2$$

Standard deviation,  $s = \sqrt{\text{variance}}$

### The binomial distribution

If  $X \sim B(n, p)$  then  $P(X = r) = {}^n C_r p^r q^{n-r}$  where  $q = 1 - p$

Mean of  $X$  is  $np$

### Kinematics

Motion in a straight line

$$v = u + at$$

$$s = ut + \frac{1}{2} at^2$$

$$s = \frac{1}{2}(u+v)t$$

$$v^2 = u^2 + 2as$$

$$s = vt - \frac{1}{2} at^2$$

## 3

- 1 A particle moves along a straight line. Its velocity  $v \text{ m s}^{-1}$  at time  $t \text{ s}$  is given by  $v = 2t + 0.6t^2$ .

Find an expression for the acceleration of the particle at time  $t$ . [2]

- 2 The height of the first part of a rollercoaster track is  $h \text{ m}$  at a horizontal distance of  $x \text{ m}$  from the start. A student models this using the equation  $h = 17 + 15 \cos 6x$ , for  $0 \leq x \leq 40$ , using the values of  $h$  given when their calculator is set to work in degrees.

(a) Find the height that the student's model predicts when the horizontal distance from the start is 40 m. [1]

(b) The student argues that the model predicts that the rollercoaster track will achieve a maximum height of 32 m more than once because the cosine function is periodic.

Comment on the validity of the student's argument. [2]

- 3 The points A and B have position vectors  $\begin{pmatrix} 2 \\ -1 \end{pmatrix}$  and  $\begin{pmatrix} 5 \\ 4 \end{pmatrix}$  respectively. The vector  $\overrightarrow{AC}$  is  $\begin{pmatrix} -2 \\ 2 \end{pmatrix}$ .


(a) Write down the position vector of C as a column vector. [1]

(b) Show that B is equidistant from A and C. [3]

- 4 **In this question you must show detailed reasoning.**


Solve the equation  $6 \cos^2 x + \sin x = 5$ , giving all the roots in the interval  $-180^\circ \leq x \leq 180^\circ$ . [5]

- 5 The graph shows displacement  $s$  m against time  $t$  s for a model of the motion of a bead moving along a straight wire. The points  $(0, 4)$ ,  $(2, 7)$ ,  $(5, 7)$  and  $(9, -7)$  are the endpoints of the line segments.



- (a) Find an expression for the displacement of the bead for  $0 \leq t \leq 2$ . [2]
- (b) Sketch the velocity-time graph for this model. [2]
- (c) Explain why the model may not be suitable at  $t = 2$  and  $t = 5$ . [1]
- 6 Show that the expression  $3x^3 + x^2 - 6x - 5$  can be written in the form  $(x+2)(ax^2 + bx + c) + d$  where  $a$ ,  $b$ ,  $c$  and  $d$  are constants to be determined. [5]

7 In this question you must show detailed reasoning.



Find the exact area of the shaded region shown in the diagram, enclosed by the  $x$ -axis and the curve  $y = -3x^2 + 7x - 2$ .


[6]

8 In this question you must show detailed reasoning.

(a) Find the centre and radius of the circle with equation  $x^2 + y^2 - 2x + 4y - 20 = 0$ . [4]

(b) Find the points of intersection of the circle with the line  $x + 3y - 10 = 0$ . [5]

- 9 The graph shows the function  $y = e^{2x}$ .



- (a) Describe the transformation of the graph of  $y = e^x$  that gives the graph of  $y = e^{2x}$ . [2]

A second function is defined by  $y = k + e^x$ .

- (b) A copy of the graph of  $y = e^{2x}$  is given in the Printed Answer Booklet.

Add a sketch of the graph of  $y = k + e^x$  in a case where  $k$  is a positive constant. [2]

- (c) Show that the two graphs do not intersect for values of  $k$  less than  $-\frac{1}{4}$ . [3]

- (d) In the case where  $k = 2$ , show that the only point of intersection occurs when  $x = \ln 2$ . [2]

- 10 Layla invests money in the bank and receives compound interest. The amount £ $L$  that she has after  $t$  years is given by the equation  $L = 2800 \times 1.023^t$ .

- (a) (i) State the amount she invests. [1]

- (ii) State the annual rate of interest. [1]


Amit invests £3000 and receives 2% compound interest per year. The amount £ $A$  that he has after  $t$  years is given by the equation  $A = ab^t$ .

- (b) Determine the values of the constants  $a$  and  $b$ . [2]

- (c) Layla and Amit invest their money in the bank at the same time.

Determine the value of  $t$  for which Layla and Amit have equal amounts in the bank. Give your answer correct to 1 decimal place. [3]

- 11** A block of mass 3 kg is at rest on a smooth horizontal table. It is attached to a light inextensible string which passes over a smooth pulley. This part of the string is horizontal. A sphere of mass 1.2 kg is attached to the other end of the string. The sphere hangs with this part of the string vertical as shown in the diagram. A horizontal force of magnitude  $F$  N is applied to the block to prevent motion.



- (a) Complete the copy of the diagram in the Printed Answer Booklet to show all the forces acting on the block and the sphere. [2]
- (b) Find the value of  $F$ . [2]

The force  $F$  N is removed, and the system begins to move.

- (c) The equation of motion of the block is  $T = 3a$ , where  $T$  N is the tension in the string and  $a$   $\text{ms}^{-2}$  is the acceleration of the block.

Write down the equation of motion of the sphere. [1]

- (d) Find the value of  $T$ . [2]

- 12** Points A, B and C lie in a straight line in that order on horizontal ground. A box of mass 5 kg is pushed from A to C by a horizontal force of magnitude 8 N. The box is at rest at A and takes 3 seconds to reach B. The ground is smooth between A and B. Between B and C the ground is rough and the resistance to motion is 28 N. The box comes to rest at C.

Determine the distance AC. [8]

**END OF QUESTION PAPER**

---

# OCR

Oxford Cambridge and RSA

## Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.