



GCE

Mathematics B MEI

H640/02: Pure Mathematics and Statistics

A Level

Mark Scheme for June 2022

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2022

Text Instructions

1. Annotations and abbreviations

Annotation in scoris	Meaning
✓ and ✕	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
E	Explanation mark 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank page
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only previous M mark.
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This indicates that the instruction In this question you must show detailed reasoning appears in the question.

2. Subject-specific Marking Instructions for AS Level Mathematics B (MEI)

- a Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. If you are in any doubt whatsoever you should contact your Team Leader.

- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words “Determine” or “Show that”, or some other indication that the method must be given explicitly.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation *isw*. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation ‘dep*’ is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case, please escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.
- Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be ‘follow through’. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

- f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.)
We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.
- When a value is **given** in the paper only accept an answer correct to at least as many significant figures as the given value.
 - When a value is **not given** in the paper accept any answer that agrees with the correct value to **2 s.f.** unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.
NB for Specification A the rubric specifies 3 s.f. as standard, so this statement reads “3 s.f”
- Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.
Candidates using a value of 9.80, 9.81 or 10 for g should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.
- g Rules for replaced work and multiple attempts:
- If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
 - If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
 - if a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.
- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate’s data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors. If a candidate corrects the misread in a later part, do not continue to follow through. E marks are lost unless, by chance, the given results are established by equivalent working. Note that a miscopy of the candidate’s own working is not a misread but an accuracy error.
- i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold “In this question you must show detailed reasoning”, or the command words “Show” and “Determine. Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.
- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
1		$R^2 = 1^2 + \sqrt{3}^2$	M1	1.1	may be implied by correct answer
		$\tan\alpha = \frac{\sqrt{3}}{1}$ or $\sin\alpha = \frac{\sqrt{3}}{2}$ or $\cos\alpha = \frac{1}{2}$ soi	M1	1.1	may see eg $\alpha = \tan^{-1}\left(\frac{\sqrt{3}}{1}\right)$ may be implied by correct answer
		$R = 2$ or $\alpha = \frac{\pi}{3}$ or $\alpha = 60^\circ$ seen	A1	1.1	
		$2\cos\left(\theta - \frac{\pi}{3}\right)$ or $2\cos(\theta - 60^\circ)$ isw	A1	1.1	
			[4]		
		<i>Alternatively</i> $\cos\theta + \sqrt{3}\sin\theta = R\cos\theta\cos\alpha + R\sin\theta\sin\alpha$ so $1 = R\cos\alpha$ and $\sqrt{3} = R\sin\alpha$ $\frac{1}{\cos\alpha} = \frac{\sqrt{3}}{\sin\alpha}$ $\alpha = \frac{\pi}{3}$ or $\alpha = 60^\circ$ seen $2\cos\left(\theta - \frac{\pi}{3}\right)$ or $2\cos(\theta - 60^\circ)$ isw	M1 M1 A1 A1		for equating coefficients for eliminating R
2		$\frac{50}{1-0.5}$ soi	M1	1.1	
		100	A1	1.1	
			[2]		

H640/02

Mark Scheme


June 2022

Question		Answer	Marks	AO	Guidance
3	(a)		<p>M1</p> <p>B1</p> <p>A1</p>	<p>1.1</p> <p>1.1</p> <p>1.1</p>	<p>decreasing concave up curve in 1st and 2nd quadrants which does not cut the x-axis; mark intent</p> <p>decreasing curve with intercept $(0,3)$; may be in one quadrant only</p> <p>smooth curve from $(-0.5, a)$ through $(2.5, b)$, where $4.5 \leq a \leq 5$ and $0 < b < 0.5$</p>
			[3]		
3	(b)	$\log(3 \times 0.4^x) = \log(0.8)$ oe $x \log 0.4 = \log 0.8 - \log 3$ oe 1.44 cao	<p>M1</p> <p>M1</p> <p>A1</p>	<p>3.1a</p> <p>1.1</p> <p>1.1</p>	<p>taking logarithms in any base</p> <p>3rd law of logs used correctly</p> <p>if M0M0 allow SC1 for 1.44 unsupported</p>
		<p><i>Alternatively</i></p> $0.4^x = \frac{0.8}{3}$ $x = \log_{0.4} \left(\frac{0.8}{3} \right)$ $x = 1.44$ cao	<p>M1</p> <p>M1</p> <p>A1</p>		<p>may see $x \log 0.4 = \log \left(\frac{0.8}{3} \right)$ oe</p>
			[3]		

H640/02

Mark Scheme


June 2022

Question		Answer	Marks	AO	Guidance
4	(a)		M1	1.1	Venn diagram with 2 overlapping regions and 0.22 correctly placed; condone incorrect or no labelling
			A1	1.1	all probabilities or percentages correctly placed and correctly labelled; ignore values in intersection allow if no box drawn if labels are eg A and B , A and B need to be defined
			[2]		
4	(b)	$0.31 + 0.23 + p(J \cap S) = 1 - 0.22$ oe	M1	1.1a	may be implied by correct answer or by 24%
		0.24 oe isw	A1	1.1	do not allow 24%
			[2]		
5		$2^n - 1$ correctly evaluated for any odd positive integer $2^n - 1$ correctly evaluated for any odd positive integer for which Tom's conjecture is false eg 511 is divisible by 7 with 9 seen [so not prime]	B1	1.1	$n \geq 3$ B0 if only rounded number in standard form seen
			B1	2.1	eg $2^9 - 1 = 511$, eg $2^{15} - 1 = 32767$ eg $2^{21} - 1 = 2097151$
			B1	2.2a	NB 32767 and 2097151 both divisible by 7; 2047 divisible by 23 correct value of n may be embedded in formula NB B0 if answer spoiled by eg so 511 is prime
			[3]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
6			M1	3.1a	area identified which is symmetrical about the mean
			A1	1.1	$x = \mu \pm \sigma$ at points of inflection
			[2]		
7		$0.4 \times 20 + 1.3 \times 10 + 3.6 \times 5 + 2 \times 10 + 0.8 \times 15$ 71	M1	1.1	allow one incorrect frequency density and/or one incorrect class width NB $8 + 13 + 18 + 20 + 12$ soi with four of five correct implies M1 may be implied by 71
			A1	1.1	if M0 allow SC1 for 8, 13, 18, 20, 12 and no others seen
			[2]		
8	(a)	population since all distances of at least 120 km are used oe	B1	1.2	
8	(b)	161-163 km	B1	1.1	
8	(c)	we need to see two elements: Ali's complaint is justified oe and correct numerical reasoning with reference to upper tail	B1	2.2a	eg 10 riders rode more than 160 km or 14 rode 156 km or more or the reserve should have ridden (approximately) 161 km so (Ali's complaint is) justified oe

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
9	(a)	mean 112.4 isw or 112 isw	B1	1.1	
		variance 8.8 or $\sqrt{8.8^2}$ cao isw	B1	1.1	B0 for 8.757 explicitly rounded to 8.8
			[2]		
9	(b)	N(their 112.4, their 8.8)	M1	3.3	allow M1 for 8.8^2 or $\sqrt{8.8}$
		$N(a, b)$	A1	1.1	$a = 112.4$ or 112 and $b = 8.8$ or 2.97^2
			[2]		
9	(c)	P(mark < 104.5) or P(mark < 105) found from their distribution in part (b)	M1	3.4	may see $N(-\infty, 104.5, 112.4, \sqrt{8.8})$ NB 0.00387 or 0.0063(06) implies M1 NB 0.00573 or 0.00914 implies M1 NB 0.00379(69..) or 0.00619(81...) may imply M1 FT use of variance = 8.757 NB 0.200(199...) and 0.184(665...) may imply M1 FT use of sd = 8.8 if probability is correctly found to be 0 eg from use of $N(112.4, \frac{8.8}{205})$ allow M1 only – no further marks available
		$205 \times \text{their non-zero } 0.00387$	M1	3.1a	or compare $\frac{1}{205}$ (≈ 0.00488) with <i>their</i> non-zero 0.00387
		0.79 to 0.794 or 1.17 to 1.175 so consistent oe	A1	3.5a	or probabilities similar so consistent oe
			[3]		

H640/02

Mark Scheme

June 2022

Question			Answer	Marks	AO	Guidance
			<p><i>Alternatively</i></p> <p>$\text{InvNorm}\left(\frac{1}{205}, 112.4, \sqrt{8.8}\right)$ or</p> <p>$\text{InvNorm}\left(\frac{1}{205}, 112, \sqrt{8.8}\right)$ used to find their mark</p> <p>compares their mark with 105</p> <p>104.7 or 104.3 is close to 105 so good fit</p>	<p>M1</p> <p>M1</p> <p>A1</p>		FT their distribution
9	(d)		<p>P(mark between 114.5 and 115.5) found</p> <p>18.75 to 18.77 so allow 18 or 19</p> <p>or 16.5 to 16.534 so allow 16 or 17</p>	<p>M1</p> <p>A1</p>	<p>3.4</p> <p>3.5a</p>	<p>NB awrt 0.0915 or awrt 0.0807 implies M1</p> <p>unsupported answers score M0</p>
				[2]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
10	(a)	$(x - 2) = 5\cos\theta$ and $(y - 1) = 5\sin\theta$ $(x - 2)^2 + (y - 1)^2 = (5\cos\theta)^2 + (5\sin\theta)^2$ oe $(x - 2)^2 + (y - 1)^2 = 5^2$ oe isw or $\frac{(x-2)^2}{5^2} + \frac{(y-1)^2}{5^2} = 1$ oe isw	M1 M1 A1	3.1a 1.1 1.1	allow sign errors or $\left(\frac{x-2}{5}\right)^2 + \left(\frac{y-1}{5}\right)^2 = \cos^2\theta + \sin^2\theta$ oe may see eg $x^2 - 4x + y^2 - 2y = 20$ if M0M0 allow SC1 for $y = 1 + 5\sin\left(\cos^{-1}\left(\frac{x-2}{5}\right)\right)$ or $x = 2 + 5\cos\left(\sin^{-1}\left(\frac{y-1}{5}\right)\right)$
			[3]		
		<i>Alternatively</i> $x^2 = (2 + 5\cos\theta)^2$ and $y^2 = (1 + 5\sin\theta)^2$ $x^2 + y^2 = 5 + 20\cos\theta + 10\sin\theta + 25\sin^2\theta + 25\cos^2\theta$ $x^2 + y^2 = 20 + 4x + 2y$ oe isw	M1 M1 A1		if only seen in expanded form, allow one coefficient error; allow sign errors must have terms in $\cos\theta$ and $\sin\theta$
		<i>Alternatively</i> radius = 5 and centre is (2, 1) $(x - 2)^2 + (y - 1)^2 = 5^2$	M1 M1 A1		allow sign error in coordinates of centre FT their centre all correct

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
10	(b)	gradient of radius is $\frac{-4}{3}$ gradient of tangent is $\frac{3}{4}$ $(y - -3) = \frac{3}{4}(x - 5)$ oe $3x - 4y - 27 = 0$ <i>or</i> $-3x + 4y + 27 = 0$	B1 M1 M1 A1	3.1a 2.1 2.4 1.1	FT 1 \div <i>their</i> $-\frac{4}{3}$ allow one sign error; FT <i>their</i> $\frac{3}{4}$ may see $-3 = \frac{3}{4} \times 5 + c$
			[4]		
		<i>Alternatively</i> $\frac{dy}{dx} = \frac{5\cos\theta}{-5\sin\theta}$ oe substitution of $\cos\theta = \frac{3}{5}$ and $\sin\theta = -\frac{4}{5}$ oe or (5,-3) in their $\frac{dy}{dx}$ $(y - -3) = \frac{3}{4}(x - 5)$ oe $3x - 4y - 27 = 0$ <i>or</i> $-3x + 4y + 27 = 0$	B1 M1 M1 A1	or $\frac{dy}{dx} = \frac{2-x}{y-1}$ oe eg $2(x - 2) + 2(y - 1)\frac{dy}{dx} = 0$ $\frac{dy}{dx} = \frac{3/5}{-(-4/5)}$ or $\frac{2-5}{-3-1}$ oe ; allow one sign error; allow one sign error; FT <i>their</i> $\frac{3}{4}$ may see $-3 = \frac{3}{4} \times 5 + c$	
			[4]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
11	(a)	Nina's, because hers is the largest sample size oe	B1	2.2a	allow eg Nina's, because with a larger sample size the probabilities get closer to the true probabilities oe
			[1]		
11	(b)	$11p + kp = 1$ $p = \frac{1}{11+k}$	M1 A1	3.1a 1.1	
			[2]		
11	(c)	<i>their</i> $\frac{1}{11+k} \times k$ or <i>their</i> $\frac{1}{11+k} \times 120$ $120 \times \textit{their} \frac{k}{11+k}$ $\frac{120k}{11+k}$ oe	M1 M1 A1	2.1 1.2 1.1	multiply by k or by 120; may be embedded multiplying by both k and 120
			[3]		
11	(d)	$32 = \textit{their} \frac{120k}{11+k}$ oe $k = 4$	M1 A1	1.1 1.1	
			[2]		

H640/02

Mark Scheme

June 2022

Question			Answer	Marks	AO	Guidance
			<i>Alternatively</i> $11p = 1 - \frac{32}{120}$ may be implied by $p = \frac{1}{15}$ (from $(P(X \neq 12))$) $k = 4$	M1 A1		or $\frac{kf}{120} = \frac{32}{120}$ (from $11f = 120 - 32 = 88$ so $f = 8$ and so $kp = \dots$) $k = 4$
11	(e)		$Y \sim B\left(30, \text{their } \frac{4}{11+4}\right)$ or $Y \sim B\left(30, \frac{32}{120}\right)$ used to find $P(Y = 8)$ $0.16 - 0.163$ BC	M1 A1	3.1a 1.1	Y is the number of 12s obtained in 30 rolls; allow B2 for $0.1628 - 0.163$ unsupported
				[2]		
12	(a)		$H_0 : \mu = 1.5$ $H_1 : \mu < 1.5$ μ is the population mean weight of flour in a bag	B1 B1	1.1 2.5	both hypotheses in terms of μ allow μ is the population mean weight of a bag of flour
				[2]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
12	(b)	$N(a, b)$ $a = 1.5$ or $b = \frac{0.24^2}{32}$ or 0.0018 $N(1.5, \frac{0.24^2}{32})$ isw or $N(1.5, 0.0018)$ isw	M1 A1 A1	3.3 2.2a 3.1a	a and b are numerical values allow 0.0424 ² for variance A0 if answer spoiled by wrong variable quoted eg $\mu \sim N(1.5, \frac{0.24^2}{32})$ or $X \sim N(1.5, \frac{0.24^2}{32})$; allow only \bar{X} oe if variable included
			[3]		
12	(c)	$0.0786 > 0.05$ or $-1.4142 > -1.645$ do not reject H_0 there is insufficient evidence at the 5% level to suggest that the mean weight of the flour in the bags is less than 1.5 kg	M1 A1 A1	3.4 1.1 2.2b	or $1.44 > 1.43(02586 \dots)$ NB 1.43(02586...) is from InvNorm(0.05, 1.5, 0.0424) allow accept H_0 or not significant or reject H_1 do not allow eg conclude / prove / indicate or other assertive statement instead of suggest if calculated values are used full marks may be awarded for awrt 0.07865 or 0.0786, or $-1.415 \leq z \leq -1.414$; otherwise award a maximum of M1A1 for 0.07...or -1.4... other calculated values score M0
			[3]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
13	(a)	<p>it can't be fully justified because</p> <p>eg different samples may lead to different conclusions oe</p> <p>eg the proportion could be 0.35 and 61/140 may have arisen by chance oe</p> <p>eg the sample may not be representative oe</p> <p>eg the researcher used a sample not a population oe</p>	B1	2.4	<p>do not allow</p> <p>eg the sample is too small</p> <p>eg the sample is too small to be representative</p>
			[1]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
13	(b)	$H_0 : p = 0.35$	B1	1.1	allow equivalent in words; do not allow percentages
		$H_1 : p > 0.35$			
		p is the probability that a baby (selected at random) is born without wisdom teeth	B1	2.5	or p is the proportion of babies that are born without wisdom teeth B1B1 if other symbol instead of p used if correctly defined
		$P(X \geq k)$ found using $B(140, 0.35)$, where $k = 60, 61$ or 62	M1*	3.3	or critical region is $X \geq k$ found from calculation of probability; allow $k = 58, 59$ or 60
		NB $P(X \geq 60) = 0.03272 - 0.033$ or $P(X \geq 62) = 0.01438 - 0.015$ NB $0.967\dots, 0.978\dots$ and $0.985\dots$ imply M1			
		$P(X \geq 61) = 0.02197 - 0.022$ their 0.022 correctly compared with 0.05	A1 M1dep*	1.1 3.4	or critical region is $X \geq 59$ from $0.0475\dots$ or 0.048 or 61 correctly compared with their 59 ; allow their 0.978 correctly compared with 0.95
		do not accept H_0 or reject H_0 or accept H_1 or significant	A1FT	1.1	A0 if their $0.022 > 0.05$ or $61 < 59$
sufficient evidence at the 5% level to suggest that the probability that a baby is born without wisdom teeth is more than 0.35	A1	1.1	dependent on award of all other marks apart from second B1 do not allow eg conclude / prove / indicate or other assertive statement instead of suggest		
			[7]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
14	(a)	$0.2 \times \{0.96154 + 0.86207 + 0.73529 + 0.60976 + 0.5\}$ soi	M1	2.1	allow M1A1 for calculation of exact values using formula in parts (a) and (b)
		$0.73373... \approx 0.7337$ AG	A1	2.4	need to see 0.73373... as well as 0.7337 for A1
			[2]		
14	(b)	$0.2 \times \{1 + 0.96154 + 0.86207 + 0.73529 + 0.60976\}$	M1	1.1	or $(3.66866 - 0.5 + 1) \times 0.2$
		0.8337 correct to 4 dp	A1	1.1	
			[2]		
14	(c)	0.1	B1	1.1	FT their $0.8337(32) - 0.7337(32)$, dependent on award of M1 in part (b)
			[1]		
14	(d)	$0.79162 - 0.77912$	M1	3.1a	if M0 allow SC1 for correct interval identified eg 0.77912 to 0.79162
		0.0125	A1	2.4	
			[2]		
14	(e)	increase n oe use rectangles of smaller width oe	B1	2.2a	do not allow eg reduce interval eg just 'smaller' rectangles – need to specify width reduction
			[1]		
15	(a)	51.635 or 51.64 or 51.6	B1	3.4	
			[1]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance
15	(b)	1995 estimate (probably) reliable since it is interpolation	B1	2.2b	allow eg the first estimate..
		2025 estimate (probably) not reliable since it is extrapolation	B1	2.2b	allow eg the second estimate...
			[2]		
15	(c)	No, because trends in life expectancy at birth may vary considerably between nations	B1	2.4	LDS advantage
			[1]		
15	(d)	series 2 (the top one) is Italy – life expectancy (generally) higher in Europe (than Africa)	B1	2.4	LDS advantage
		the values are decreasing (from 1990) in South Africa (– unusual since most show an upward trend) or little (or no) overall increase in South Africa (since 1970) or South Africa has lower life expectancy (than most developed countries)	B1	2.4	LDS advantage
			[2]		

H640/02

Mark Scheme

June 2022


Question		Answer	Marks	AO	Guidance
15	(e)	<p>Scatter diagram of life expectancy at birth in 2010 against GDP per capita in US \$</p> <p>The scatter diagram plots life expectancy at birth in 2010 on the y-axis (ranging from 40.00 to 90.00) against GDP per capita in US \$ on the x-axis (ranging from 0 to 150,000). A cluster of blue dots shows a positive correlation between GDP per capita and life expectancy, with most points between 20,000 and 100,000 US \$ GDP per capita and 70.00 to 85.00 life expectancy. One point at approximately (700, 47.56) is circled in orange, representing a country with very low GDP per capita and low life expectancy.</p>	B1	1.1	<p>Point at (700, 47.56) ringed</p> <p>LDS advantage</p>
			[1]		
15	(f)	<p>the diagram supports this statement for values of GDP per capita from k to n where $0 < k \leq 20\,000$ and $40\,000 \leq n \leq 60\,000$ since there appears to be positive correlation oe</p> <p>for values of GDP per capita $\geq K$ where $40\,000 \leq K \leq 60\,000$ there appears to be no association between GDP per capita and life expectancy at birth so the diagram does not support Sundip's statement for these values</p>	B1	2.3	<p>must give specific range of values ; must say supports statement oe</p>
			B1	2.2b	<p>the range may be implied by reference to a specific range identified for the first mark; must say does not support statement oe</p>
			[2]		

H640/02

Mark Scheme

June 2022

Question		Answer	Marks	AO	Guidance							
16	(a)	$\frac{dy}{dx} = 24x^3 + 24x^2 - 42x + 12$	M1	3.1a	allow one sign or coefficient error; must be four terms							
		their $\frac{dy}{dx} = 0$	M1	1.1	at least two terms correct							
		$f(k)$ evaluated, where k is a factor of ± 12 or $\pm \frac{a}{12}$, where $a = 1, 2, 3, 4$ or 6	M1	2.1	may be implied by $x = -2$ seen unsupported or $(x + 2)$ identified as factor							
		$(x + 2)(4x^2 - 4x + 1)$ or $(2x - 1)(2x^2 + 3x - 2)$	M1	3.1a	by inspection or long division; allow one sign error or one coefficient error in trinomial may be implied by $x = \frac{1}{2}$ seen unsupported or $(2x - 1)$ oe identified as factor							
		$x = -2$ and $x = \frac{1}{2}$ and no others	A1	1.1	may see $x = \frac{1}{2}$ (repeated) A0 for $x = -2$ (repeated)							
		$(\frac{1}{2}, -\frac{31}{8})$ and $(-2, -82)$ and no others	A1	1.1								
		$\frac{d^2y}{dx^2} = 72x^2 + 48x - 42$	M1*	1.1	allow one sign or one coefficient error, FT their $\frac{dy}{dx}$; allow M1 for $12x^2 + 8x - 7$							
		$\frac{d^2y}{dx^2} = 150$ when $x = -2$ so minimum value	A1	1.1	NB test indecisive at $x = \frac{1}{2}$ A0 for just eg $\frac{d^2y}{dx^2} > 0$ so minimum							
or eg				award M1A1 for consideration of gradient either side of -2 , values must be correct to at least 2sf for A1								
		<table border="1"> <tbody> <tr> <td>x</td> <td>-2.1</td> <td>(-2)</td> <td>-1.9</td> </tr> <tr> <td>$\frac{dy}{dx}$</td> <td>-16.224</td> <td>(0)</td> <td>13.824</td> </tr> </tbody> </table>	x	-2.1	(-2)	-1.9	$\frac{dy}{dx}$	-16.224	(0)	13.824		
x	-2.1	(-2)	-1.9									
$\frac{dy}{dx}$	-16.224	(0)	13.824									

Question		Answer	Marks	AO	Guidance																								
		eg <table border="1" style="display: inline-table; margin-left: 10px;"> <tr> <td>x</td> <td>0</td> <td>$(\frac{1}{2})$</td> <td>1</td> </tr> <tr> <td>$\frac{dy}{dx}$</td> <td>12</td> <td>(0)</td> <td>18</td> </tr> </table> <p>dependent on at least two terms correct in derivative; must see values</p> <p>inflection at $(\frac{1}{2}, -\frac{31}{8})$ their $72x^2 + 48x - 42 = 0$ $x = -\frac{7}{6}$ isw</p>	x	0	$(\frac{1}{2})$	1	$\frac{dy}{dx}$	12	(0)	18	<p>M1</p> <p>A1</p> <p>M1dep*</p> <p>A1</p>	<p>3.1a</p> <p>3.2a</p> <p>1.1</p> <p>1.1</p>	or eg <table border="1" style="display: inline-table; margin-left: 10px;"> <tr> <td>x</td> <td>0</td> <td>$(\frac{1}{2})$</td> <td>1</td> </tr> <tr> <td>y</td> <td>-6</td> <td>(-3.875)</td> <td>-1</td> </tr> </table> <p>or eg</p> <table border="1" style="display: inline-table; margin-left: 10px;"> <tr> <td>x</td> <td>0</td> <td>$(\frac{1}{2})$</td> <td>1</td> </tr> <tr> <td>$\frac{d^2y}{dx^2}$</td> <td>-42</td> <td>(0)</td> <td>78</td> </tr> </table> <p>values in table must be correct</p> <p>ignore calculation of associated y-value</p> <p>allow any correct decimals to 3 sf or more</p>	x	0	$(\frac{1}{2})$	1	y	-6	(-3.875)	-1	x	0	$(\frac{1}{2})$	1	$\frac{d^2y}{dx^2}$	-42	(0)	78
x	0	$(\frac{1}{2})$	1																										
$\frac{dy}{dx}$	12	(0)	18																										
x	0	$(\frac{1}{2})$	1																										
y	-6	(-3.875)	-1																										
x	0	$(\frac{1}{2})$	1																										
$\frac{d^2y}{dx^2}$	-42	(0)	78																										
			[12]																										
16	(b)		<p>M1</p> <p>B1</p> <p>A1</p>	<p>1.1</p> <p>1.1</p> <p>1.1</p>	<p>curve with a minimum in 3rd quadrant and stationary point of inflection in 4th quadrant and no other stationary points</p> <p>(0, -6) identified as y-intercept (intercept must be below the x-axis and above -20)</p> <p>correct curve with intercepts at $(-a,0)$ and $(b,0)$, where $-3 < a < -2.6$ and $0.8 < b < 1.2$; minimum at $(-2, y)$ where $-90 < y < -80$ and inflection for $0 < x < 1$ and y is between the x-axis and the y-intercept</p>																								
			[3]																										

Need to get in touch?

If you ever have any questions about OCR qualifications or services (including administration, logistics and teaching) please feel free to get in touch with our customer support centre.

Call us on

01223 553998

Alternatively, you can email us on

support@ocr.org.uk

For more information visit

 ocr.org.uk/qualifications/resource-finder

 ocr.org.uk

 [Twitter/ocrexams](https://twitter.com/ocrexams)

 [/ocrexams](https://twitter.com/ocrexams)

 [/company/ocr](https://www.linkedin.com/company/ocr)

 [/ocrexams](https://www.youtube.com/ocrexams)



OCR is part of Cambridge University Press & Assessment, a department of the University of Cambridge.

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored. © OCR 2022 Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee. Registered in England. Registered office The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA.

Registered company number 3484466. OCR is an exempt charity.

OCR operates academic and vocational qualifications regulated by Ofqual, Qualifications Wales and CCEA as listed in their qualifications registers including A Levels, GCSEs, Cambridge Technicals and Cambridge Nationals.

OCR provides resources to help you deliver our qualifications. These resources do not represent any particular teaching method we expect you to use. We update our resources regularly and aim to make sure content is accurate but please check the OCR website so that you have the most up-to-date version. OCR cannot be held responsible for any errors or omissions in these resources.

Though we make every effort to check our resources, there may be contradictions between published support and the specification, so it is important that you always use information in the latest specification. We indicate any specification changes within the document itself, change the version number and provide a summary of the changes. If you do notice a discrepancy between the specification and a resource, please [contact us](#).

Whether you already offer OCR qualifications, are new to OCR or are thinking about switching, you can request more information using our [Expression of Interest form](#).

Please [get in touch](#) if you want to discuss the accessibility of resources we offer to support you in delivering our qualifications.