

OCR

Oxford Cambridge and RSA

Accredited

A Level Mathematics B (MEI)

H640/03 Pure Mathematics and Comprehension

Sample Insert

Version 2

Date – Morning/Afternoon

Time allowed: 2 hours

INFORMATION FOR CANDIDATES

- This insert contains the article for Section B
- This document consists of **4** pages. Any blank pages are indicated.

INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Insert for marking, it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

Archimedes's approximation of π

The constant π is defined to be the circumference of a circle divided by its diameter.

The value of π has been determined to an accuracy of more than twelve trillion decimal places. To the non-mathematician this may appear strange since it is not possible to measure the circumference and diameter of a circle to that degree of accuracy; this article explains how one of the greatest mathematicians of all time found the value of π to a high degree of accuracy without requiring any physical measurement.

Archimedes (287-212 BC) lived in Syracuse, Sicily. He developed many branches of mathematics, including calculus, in which he devised methods for finding areas under parabolas nearly 2000 years before Newton and Leibniz, and mechanics, in which he found the centres of gravity of various plane figures and solids and devised a method for calculating the weight of a body immersed in a liquid.

Whilst absorbed in a mathematical problem, Archimedes was killed by a soldier during the capture of Syracuse by the Romans.

Archimedes's method for determining the value of π is described below.

Fig. C1 shows a circle with unit radius and two regular hexagons.

The smaller regular hexagon has its vertices on the circle; it is called an *inscribed* polygon. Its perimeter is 6.

The larger regular hexagon has the midpoints of its edges on the circle; it is called an *escribed* polygon. Its perimeter is $4\sqrt{3}$.




Fig. C1

The circumference of the circle is greater than the perimeter, ABCDEF, of the smaller hexagon but less than the perimeter, A'B'C'D'E'F', of the larger hexagon.

Dividing the perimeters by the diameter of the circle gives lower and upper bounds for π of 3 and $2\sqrt{3}$, so that $3 < \pi < 2\sqrt{3}$.

- 25 To find tighter bounds, Archimedes repeatedly doubled the number of edges in the two regular polygons, from 6 to 12, 24, 48 and finally 96. The process of doubling the number of edges is described below.

- 30 **Fig. C2** shows two adjacent vertices, P and Q, of a regular polygon inscribed in a circle with unit radius and centre O. PQ has length a . M is the midpoint of PQ. OM is extended to meet the circle at R. MR has length h . PR and RQ are adjacent edges of a regular polygon which has twice as many edges as the polygon which has PQ as an edge. PR has length b .




Fig. C2

Applying Pythagoras' Theorem

- to triangle OMP gives $1 = \frac{a^2}{4} + (1-h)^2$,
- to triangle PMR gives $b^2 = \frac{a^2}{4} + h^2$.

- 35 For the inscribed regular hexagon, $a=1$. Substituting $a=1$ in the equations above gives $h = \frac{2-\sqrt{3}}{2}$ and $b = \sqrt{2-\sqrt{3}}$. This can be written in the equivalent form $b = \frac{\sqrt{6}-\sqrt{2}}{2}$.

Therefore a regular polygon with 12 edges inscribed in a unit circle has edge length $\frac{\sqrt{6}-\sqrt{2}}{2}$.

Archimedes repeated this process to find the edge lengths of inscribed regular polygons with 24, 48 and 96 edges. He then used a similar technique for escribed regular polygons.

- 40 The inscribed and escribed regular polygons with 96 edges provide bounds for π which we now write, using decimal notation, as $3.14103... < \pi < 3.14271...$.

Summary of Updates

Date	Version	Change
October 2018	2	We've reviewed the look and feel of our papers through text, tone, language, images and formatting. For more information please see our assessment principles in our "Exploring our question papers" brochures on our website.

Specimen

Copyright Information:

OCR is committed to seeking permission to reproduce all third-party content that it uses in the assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge, CB2 9EA.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.