



GCE

Mathematics B (MEI)

H640/03: Pure Mathematics and Comprehension

Advanced GCE

Mark Scheme for Autumn 2021

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021

H640/03

Mark Scheme

November 2021

1. Annotations and abbreviations

Annotation in scoris	Meaning
✓ and *	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
E	Explanation mark 1
SC	Special case
^	Omission sign
MR	Misread
BP	Blank page
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *. The * may be omitted if only previous M mark.
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This indicates that the instruction In this question you must show detailed reasoning appears in the question.

H640/03

Mark Scheme

November 2021

2. Subject-specific Marking Instructions for AS/A Level Further Mathematics B (MEI)

- a a Annotations must be used during your marking. For a response awarded zero (or full) marks a single appropriate annotation (cross, tick, M0 or ^) is sufficient, but not required.

For responses that are not awarded either 0 or full marks, you must make it clear how you have arrived at the mark you have awarded and all responses must have enough annotation for a reviewer to decide if the mark awarded is correct without having to mark it independently.

It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

Award NR (No Response)

- if there is nothing written at all in the answer space and no attempt elsewhere in the script
- OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
- OR if there is a mark (e.g. a dash, a question mark, a picture) which isn't an attempt at the question.

Note: Award 0 marks only for an attempt that earns no credit (including copying out the question).

If a candidate uses the answer space for one question to answer another, for example using the space for 8(b) to answer 8(a), then give benefit of doubt unless it is ambiguous for which part it is intended.

- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not always be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. If you are in any doubt whatsoever you should contact your Team Leader.

H640/03

Mark Scheme

November 2021

c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A method mark may usually be implied by a correct answer unless the question includes the DR statement, the command words “Determine” or “Show that”, or some other indication that the method must be given explicitly.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation ‘dep*’ is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case, please escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.
- Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be ‘follow through’. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

H640/03

Mark Scheme

November 2021

- f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.)

We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so.

- When a value is **given** in the paper only accept an answer correct to at least as many significant figures as the given value.
- When a value is **not given** in the paper accept any answer that agrees with the correct value to **2 s.f.** unless a different level of accuracy has been asked for in the question, or the mark scheme specifies an acceptable range.

NB for Specification A the rubric specifies 3 s.f. as standard, so this statement reads “3 s.f”

Follow through should be used so that only one mark in any question is lost for each distinct accuracy error.

Candidates using a value of 9.80, 9.81 or 10 for g should usually be penalised for any final accuracy marks which do not agree to the value found with 9.8 which is given in the rubric.

- g Rules for replaced work and multiple attempts:

- If one attempt is clearly indicated as the one to mark, or only one is left uncrossed out, then mark that attempt and ignore the others.
- If more than one attempt is left not crossed out, then mark the last attempt unless it only repeats part of the first attempt or is substantially less complete.
- if a candidate crosses out all of their attempts, the assessor should attempt to mark the crossed out answer(s) as above and award marks appropriately.

- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate’s data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A or B mark in the question. Marks designated as cao may be awarded as long as there are no other errors. If a candidate corrects the misread in a later part, do not continue to follow through. E marks are lost unless, by chance, the given results are established by equivalent working. Note that a miscopy of the candidate’s own working is not a misread but an accuracy error.

- i If a calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers provided that there is nothing in the wording of the question specifying that analytical methods are required such as the bold “In this question you must show detailed reasoning”, or the command words “Show” and “Determine. Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.

- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

H640/03

Mark Scheme


November 2021

Question		Answer	Marks	AOs	Guidance
1	(a)	$a = 4$ $b = -14$	B1 B1 [2]	1.1a 1.1	May be seen as $(x + 4)^2 - 14$
1	(b)	$(-4, -14)$	B1 [1]	2.2a	FT <i>their</i> (a)
1	(c)	Translation $\begin{pmatrix} -4 \\ -14 \end{pmatrix}$ oe	B1 B1 [2]	1.2 2.2a	If other transformations as well (eg stretch) then B0 'Shift' does not score FT <i>their</i> (a)
2		8.7° 81.3°	B1 B1 [2]	1.1a 1.1	If two solution given scoring B0 B0 allow B1 for awrt either answer
3	(a)	$3x = k + 1$ or $2k - 3y = 1$ $x = \frac{k+1}{3}$ oe isw $y = \frac{2k-1}{3}$ oe isw	M1 A1 A1 [3]	1.1a 1.1 2.2a	Elimination of one of x, y May be as part of a pair of coordinates May be as part of a pair of coordinates
					Allow correct unsimplified answers

H640/03

Mark Scheme


November 2021

3	(b)	$k - x = x^2 + k$ $x^2 + x = 0 \Rightarrow x = 0, x = -1$ $(0, k)$ $(-1, k + 1)$	M1 A1 A1 A1 [4]	3.1a 1.1 1.1 2.1	$(0, k)$ unsupported earns B1 SC Need not be as coordinates but must be clear which y goes with which x	Substituting for x gets M0 unless it leads to correct values of y
4	(a)	$\frac{4.13746... - 4}{1.1 - 1}$ 1.37	M1 A1 [2]	1.1a 1.1	Attempt to find gradient Do not penalise more accurate answer 1.37(46 ...)	Condone 4.13 or 4.14 for M1
4	(b)	Suitable value	B1 [1]	1.1	Anything between 1 and 1.1	
5	(a)	(i) 	B1	1.2		

H640/03

Mark Scheme

November 2021

5	(a)	(ii)		M1 A1	2.2a 2.2a	Straight line (+ve gradient) through origin (may stop short of origin) First quadrant only	
				[3]			
5	(b)	(i)	<p>Suitable reason e.g.</p> <ul style="list-style-type: none"> Reasonable to assume population growth is proportional to population Populations are often modelled by exponential growth 	E1	3.3	Allow e.g. wolves give birth to more wolves than they started with Do not allow e.g. population is proportional to time	
				[1]			
5	(b)	(ii)	$A = 21$ $51 = 21e^k$ $k = \ln\left(\frac{51}{21}\right) = \ln\left(\frac{17}{7}\right) \approx 0.887$ or better	B1 M1 A1	3.4 3.4 1.1	Allow 0.89	
				[3]			
5	(b)	(iii)	<p>Suitable reason, e.g.</p> <ul style="list-style-type: none"> Population cannot keep growing The wolves will run out of food if the population gets too big 	E1	3.5b	Allow e.g. lack of resources or deforestation	
				[1]			

H640/03

Mark Scheme


November 2021

6		<p>DR</p> $\frac{1}{\sqrt{2} + 1} + \frac{1}{\sqrt{3} + \sqrt{2}} + \frac{1}{\sqrt{4} + \sqrt{3}}$ $\frac{\sqrt{2} - 1}{2 - 1} + \frac{\sqrt{3} - \sqrt{2}}{1} + \frac{\sqrt{4} - \sqrt{3}}{1}$ $\sqrt{2} - 1 + \sqrt{3} - \sqrt{2} + 2 - \sqrt{3} = 1$	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>[4]</p>	<p>1.1a</p> <p>3.1a</p> <p>1.1</p> <p>2.1</p>	<p>Substituting values</p> <p>Attempt to rationalise denominator for one term</p> <p>All correct</p> <p>Convincing completion (AG)</p>	<p>Either $\times \frac{\sqrt{2}-1}{\sqrt{2}-1}$ or $\frac{\sqrt{2}-1}{2-1}$ at least once for M1</p>
---	--	--	---	---	---	---

H640/03

Mark Scheme

November 2021

7		<p>Let $u = x$, $\frac{dv}{dx} = \cos 2x$</p> $v = \frac{1}{2} \sin 2x$ $\int x \cos 2x \, dx = \frac{1}{2} x \sin 2x - \int \frac{1}{2} \sin 2x \, dx$ $\frac{1}{2} x \sin 2x + \frac{1}{4} \cos 2x + c$	<p>M1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>3.1a</p> <p>1.1</p> <p>2.5</p>	<p>Parts with $u = x$, $\frac{dv}{dx} = \cos 2x$</p> <p>Allow if $-\left[\frac{1}{2} x \sin 2x - \int \frac{1}{2} \sin 2x \, dx\right]$ or if 1 error</p> <p>+c needed for A1</p>	
8		<p>$a = 9$</p> <p>(-3, 1)</p> <p>(1, 9)</p> <p>(-1, 9)</p>	<p>M1</p> <p>A2</p> <p>[3]</p>	<p>3.1a</p> <p>1.1</p> <p>2.2a</p>	<p>Implied by any of the points</p> <p>A2 for all three points. A1 for one correct point</p>	<p>Ignore (3, 1) If other incorrect points A1 max</p>
9	(a)		<p>B1</p> <p>[1]</p>	<p>1.1</p>	<p>Correct line through ((0, -1) and (1, 4)</p>	<p>If x-intercept marked and (1, 4) slightly out may award B1 BOD</p>

H640/03

Mark Scheme

November 2021

9	(b)	DR $3 - \sqrt{x} = 5x - 1$ or $y = 5(3 - y)^2 - 1$ $4 - 5x = \sqrt{x} \Rightarrow (4 - 5x)^2 = x$	M1	3.1a		
		$25x^2 - 41x + 16 = 0$ or $5y^2 - 31y + 44 = 0$	M1	1.1	Getting into suitable form for solution ie =0	Or as a quadratic in \sqrt{x} $5x + \sqrt{x} - 4 = 0$
		$(x - 1)(25x - 16) = 0$ or $(5y - 11)(y - 4) = 0$	M1	1.1	Attempt to solve quadratic by formula or factorising (oe, via \sqrt{x} quadratic)	$(5\sqrt{x} - 4)(\sqrt{x} + 1) = 0$
		$x = \frac{16}{25} = 0.64$ or $y = 2.2$	A1	2.2a	Correct root chosen Or replace M1A1 with SC1 if $x = 0.64$ or $y = 2.2$ seen with no method for solving	$\sqrt{x} = \frac{4}{5}$ so $x = 0.64$
		$y = 5 \times 0.64 - 1 = 2.2$ or $x = (3 - 2.2)^2 = 0.64$	M1	1.1	FT <i>their</i> positive root	
		$\frac{1}{2}(0.64 - 0.2)(2.2)$ [= 0.484 = $\frac{121}{250}$]	M1	3.1a	Attempt to find area of triangle	$\frac{1}{2} \times 0.64 \times 2.2$ is M0
		Alternative method for area of triangle $\int_{0.2}^{0.64} (5x - 1) dx$	M1 M1		Correct integral Correct limits (FT <i>their</i> positive root)	
		$\int_{0.64}^4 (3 - \sqrt{x}) dx$	M1*	2.1	Allow any limits $0 \leq x \leq 4$	Allow if clearly embedded eg $\int_{0.64}^4 (5x - 4 + \sqrt{x}) dx$
		$\left[3x - \frac{2}{3}x^{\frac{3}{2}}\right]_{0.64}$	M1**	1.1	Integration of M1* integral \sqrt{x} term correct (ignore limits)	Dep on M1* only
		$\left(12 - \frac{16}{3}\right) - \left(1.92 - \frac{128}{375}\right) = \frac{636}{125} = 5.088$	M1	1.1	Evaluation of M1* integral substitution seen	Dep on M1**
Total area = $0.484 + 5.088 = 5.572 = \frac{1393}{250} = 5\frac{143}{250}$	A1	1.1	All correct, other partitions possible	Dep on all 9 previous marks		
			[10]			

H640/03

Mark Scheme

November 2021

10	(a)	$\frac{1}{(4x+1)(x+1)} = \frac{A}{4x+1} + \frac{B}{x+1}$ $1 = A(x+1) + B(4x+1)$ $x = -1 \Rightarrow 1 = -3B \Rightarrow B = -\frac{1}{3}$ $x = -\frac{1}{4} \Rightarrow 1 = \frac{3}{4}A \Rightarrow A = \frac{4}{3}$ <p>So $\frac{1}{(4x+1)(x+1)} = \frac{4}{3(4x+1)} - \frac{1}{3(x+1)}$</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>[3]</p>	<p>1.1a</p> <p>1.1</p> <p>1.1</p>	<p>Method mark is implied by correct answer.</p> <p>Final solution needed for A1. Can be recovered in 10(b)</p>	
10	(b)	$\int \frac{1}{y} dy = \int \frac{1}{(4x+1)(x+1)} dx$ $\int \frac{1}{y} dy = \int \left\{ \frac{4}{3(4x+1)} - \frac{1}{3(x+1)} \right\} dx$ $\ln y = \frac{1}{3} \ln 4x+1 - \frac{1}{3} \ln x+1 + c$ <p>When $x = 0$, $y = 2 \Rightarrow c = \ln 2$</p> $\ln y = \ln 2 \left(\frac{ 4x+1 }{ x+1 } \right)^{\frac{1}{3}}$ $y = 2 \left(\frac{4x+1}{x+1} \right)^{\frac{1}{3}}$	<p>M1</p> <p>M1</p> <p>M1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>3.1a</p> <p>2.2a</p> <p>1.1</p> <p>2.2a</p> <p>1.1</p> <p>2.1</p>	<p>Separation of variables - both sides seen</p> <p>Use of <i>their</i> (a). RHS only needed</p> <p>Integration. One correct x term (ft their partial fractions) Condone missing modulus signs</p> <p>Finding constant (FT <i>their</i> (a))</p> <p>For use of $\ln m = \ln m^a$ and $\ln m - \ln n = \ln \frac{m}{n}$ Not dep on c Condone missing modulus signs throughout</p> <p>Answer with correct values of A and B.</p>	<p>Integral signs, dy and dx needed</p> <p>Condone no dx</p>

H640/03

Mark Scheme


November 2021

11		<p>DR</p> <p>In triangle BDC, $\cos D = \frac{7^2+5^2-8^2}{2 \times 7 \times 5}$</p> $\cos D = \frac{1}{7}$ $\sin D = \sqrt{1 - \frac{1}{49}}$ $\sin D = \frac{\sqrt{48}}{7}$ $\frac{AB}{\sin D} = \frac{7}{\sin 45^\circ} \Rightarrow \frac{7AB}{\sqrt{48}} = \frac{7 \times 2}{\sqrt{2}}$ <p>$AB = 4\sqrt{6}$ [cm] oe</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>[5]</p>	<p>3.1a</p> <p>1.1</p> <p>1.1</p> <p>3.1a</p> <p>2.2a</p>	<p>Use of cosine rule in triangle BDC (for any angle)</p> <p>Approximate values must not be seen to earn M1 i.e. must be exact</p> <p>Use of sin rule in triangle ABD Exact values must be seen to earn M1</p> <p>Must be exact answer</p>	<p>Or</p> $\cos C = \frac{5^2 + 8^2 - 7^2}{2 \times 5 \times 8}$ <p>Or $\cos C = \frac{1}{2}$</p> <p>Or $C = 60^\circ$</p> <p>Or $\sin C = \frac{\sqrt{3}}{2}$</p> <p>Or use of sin rule in triangle ABC</p> $\frac{2AB}{\sqrt{3}} = \frac{8 \times 2}{\sqrt{2}}$
12		$\frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} = 1 \Rightarrow \frac{\frac{1}{2} + \tan \beta}{1 - \frac{1}{2} \tan \beta} = 1$ $\frac{1}{2} + \tan \beta = 1 - \frac{1}{2} \tan \beta$ $1.5 \tan \beta = 0.5 \Rightarrow \tan \beta = \frac{0.5}{1.5} = \frac{1}{3}$ $\Rightarrow \beta = \arctan\left(\frac{1}{3}\right)$	<p>M1</p> <p>M1</p> <p>E1</p> <p>[3]</p>	<p>1.1a</p> <p>1.1</p> <p>2.1</p>	<p>Use of $\tan(\alpha + \beta)$</p> <p>Rearranging</p> <p>Convincing completion. AG</p>	<p>Accept valid alternative solutions</p> <p>Approximate solutions with decimal angles do not score.</p>

H640/03

Mark Scheme

November 2021

13	(a)	$\arctan\left(\frac{1}{x}\right) = \text{angle BEA} = \frac{\pi}{2} - \theta$ <p>So $\arctan x + \arctan\left(\frac{1}{x}\right) = \frac{\pi}{2}$</p>	E1 [1]	2.4	Convincing explanation of given result	Must relate to triangle. Do not need to mention angle BEA.
13	(b)		B1 [1]	1.2		Ignore any values on axes
13	(c)	Increasing [function]	E1 [1]	2.4		

H640/03

Mark Scheme

November 2021

14	(a)		If $n = 1, n + 1 = 2$ and $n^2 + n + 1 = 3$	E1 [1]	2.2a	
14	(b)		For n a positive integer, $\frac{1}{n} \left(\frac{1}{n^2+n+1} \right) < 1$ $\arctan\left(\frac{1}{n+1}\right) + \arctan\left(\frac{1}{n^2+n+1}\right)$ $= \arctan\left(\frac{\left(\frac{1}{n+1}\right) + \left(\frac{1}{n^2+n+1}\right)}{1 - \left(\frac{1}{n+1}\right)\left(\frac{1}{n^2+n+1}\right)}\right)$ $\arctan\left(\frac{n^2+n+1+n+1}{(n^2+n+1)(n+1)-1}\right)$ $\arctan\left(\frac{n^2+2n+2}{n^3+2n^2+2n}\right) = \arctan\left(\frac{n^2+2n+2}{n(n^2+2n+2)}\right)$ $= \arctan\left(\frac{1}{n}\right)$	E1 M1 M1 A1 [4]	2.3 3.1a 1.1 2.1	Use of arctan addition formula Clearing fractions within the fraction Condone omission of arctan for this mark and condone 1 other error Convincing completion (AG)

H640/03

Mark Scheme

November 2021

15	Use of $\arctan x + \arctan \left(\frac{1}{x}\right) = \frac{\pi}{2}$	M1	3.1a	eg $\arctan 2 + \arctan \left(\frac{1}{2}\right) = \frac{\pi}{2}$	Allow valid alternative methods
	$\arctan \left(\frac{1}{2}\right) + \arctan 2 + \arctan \left(\frac{1}{3}\right) + \arctan 3 = \pi$	M1	1.1	Oe	
	Use of $\arctan \left(\frac{1}{2}\right) + \arctan \left(\frac{1}{3}\right) = \arctan 1$	M1	1.1	Oe	
	$\Rightarrow \arctan 1 + \arctan 2 + \arctan 3 = \pi$	E1	2.4	Convincing completion possibly using the three results above (AG)	
	Alternative method				
	$\arctan 2 + \arctan 3 = \arctan \left(\frac{2+3}{1-6}\right) + \pi$	M1		Use of $\arctan x + \arctan y = \arctan \left(\frac{x+y}{1-xy}\right) + \pi$	
$\arctan 2 + \arctan 3 = \arctan(-1) + \pi$	M1				
$\arctan 1 = -\arctan(-1)$	M1				
$\arctan 2 + \arctan 3 = -\arctan 1 + \pi$	E1				
$\Rightarrow \arctan 1 + \arctan 2 + \arctan 3 = \pi$					
		[4]			

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored