Surname		Centre Number	Candidate Number
First name(s)			2

wjec cbac

GCE A LEVEL

1305U40-1

S24-1305U40-1

MONDAY, 3 JUNE 2024 – AFTERNOON

FURTHER MATHEMATICS – A2 unit 4
FURTHER PURE MATHEMATICS B

2 hours 30 minutes

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1	11	
2	13	
3	9	
4	21	
5	14	
6	8	
7	12	
8	11	
9	9	
10	12	
Total	120	

ADDITIONAL MATERIALS

In addition to this examination paper, you will need:
- a Formula Booklet;
- a calculator.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

You may use a pencil for graphs and diagrams only.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The maximum mark for this paper is 120.

The number of marks is given in brackets at the end of each question or part-question.

Sufficient working must be shown to demonstrate the **mathematical** method employed.

Answers without working may not gain full credit.

Unless the degree of accuracy is stated in the question, answers should be rounded appropriately.

You are reminded of the necessity for good English and orderly presentation in your answers.

© WJEC CBAC Ltd.   SJJ/GR*(S24-1305U40-1)

2

Reminder: Sufficient working must be shown to demonstrate the **mathematical** method employed.

1.   (a)   Express the three cube roots of $5+10\mathrm{i}$ in the form $r\mathrm{e}^{\mathrm{i}\theta}$, where $0 \leqslant \theta < 2\pi$.   [6]

(1305U40-1)

(b) The three cube roots of $5+10i$ are plotted in an Argand diagram. The points are joined by straight lines to form a triangle. Find the area of this triangle, giving your answer correct to two significant figures. [5]

© WJEC CBAC Ltd. (1305U40-1) **Turn over.**

4

2. The function f is defined by $f(x) = \cosh\left(\dfrac{x}{2}\right)$.

(a) State the Maclaurin series expansion for $\cosh\left(\dfrac{x}{2}\right)$ up to and including the term in x^4.

[2]

....

....

....

....

....

....

....

....

Another function g is defined by $g(x) = x^2 - 2$. The diagram below shows parts of the graphs of $y = f(x)$ and $y = g(x)$.

© WJEC CBAC Ltd.

(1305U40-1)

(b)   The two graphs intersect at the point A, as shown in the diagram. Use your answer from part (a) to find an approximation for the x-coordinate of A, giving your answer correct to two decimal places.   [5]

© WJEC CBAC Ltd.   (1305U40-1)   **Turn over.**

6

(c)   Using your answer to part (b), find an approximation for the area of the shaded region enclosed by the two graphs, the x-axis and the y-axis.　　　　[6]

© WJEC CBAC Ltd.　　　　(1305U40-1)

BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

(1305U40-1)

Turn over.

3. Given the differential equation

$$\cos x \frac{dy}{dx} + y \sin x = 4\cos^3 x \sin x + 5$$

and $y = 3\sqrt{2}$ when $x = \frac{\pi}{4}$, find an equation for y in terms of x. [9]

© WJEC CBAC Ltd.   (1305U40-1)

1305U401
09

(1305U40-1) **Turn over.**

4. (a) Given that $z^n + \dfrac{1}{z^n} = 2\cos n\theta$, where $z = \cos\theta + i\sin\theta$, express $16\cos^4\theta$ in the form

$a\cos 4\theta + b\cos 2\theta + c$, where a, b, c are integers whose values are to be determined. [5]

© WJEC CBAC Ltd. (1305U40-1)

The diagram below shows a sketch of the curve C with polar equation

$$r = 3 - 4\cos^2\theta, \quad \text{where } \frac{\pi}{6} \leqslant \theta \leqslant \frac{5\pi}{6}.$$

Initial line

(b) Calculate the area of the region enclosed by the curve *C*.   [8]

© WJEC CBAC Ltd.   (1305U40-1)   **Turn over.**

(c)   Find the exact polar coordinates of the points on *C* at which the tangent is perpendicular to the initial line.   [8]

BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

(1305U40-1) **Turn over.**

5. Find each of the following integrals.

(a) $\displaystyle\int \frac{3-x}{x\left(x^2+1\right)}\,\mathrm{d}x$ [8]

© WJEC CBAC Ltd. (1305U40-1)

(b) $\displaystyle\int \frac{\sinh 2x}{\sqrt{\cosh^4 x - 9\cosh^2 x}}\,dx$ [6]

© WJEC CBAC Ltd.   (1305U40-1)   **Turn over.**

6. The matrix $\mathbf{M}$ is defined by

$$\mathbf{M} = \begin{pmatrix} 12 & 30 & 8 \\ 18 & 25 & 20 \\ 19 & 50 & 16 \end{pmatrix}.$$

(a) Given that $\det \mathbf{M} = -1040$, give a geometrical interpretation of the solution to the following equation. [2]

$$\begin{pmatrix} 12 & 30 & 8 \\ 18 & 25 & 20 \\ 19 & 50 & 16 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2668 \\ 3402 \\ 4581 \end{pmatrix}$$

....

....

....

....

....

....

....

(b) Three hotels A, B, C each have different types of room available to book: single, double and family rooms. For each type of room, the price per night is the same in each of the three hotels.

The table below gives, for each hotel, details of the number of each type of room and the total revenue per night when the hotel is full.

Hotel	Types of room			Total revenue
	Single	Double	Family	
A	12	30	8	£2,668
B	18	25	20	£3,402
C	19	50	16	£4,581

(1305U40-1)

Find the price per night of each type of room. [6]

© WJEC CBAC Ltd. (1305U40-1) **Turn over.**

7. (a) A curve C is defined by the equation $y = \dfrac{1}{\sqrt{16 - 6x - x^2}}$ for $-3 \leqslant x \leqslant 1$.

(i) Find the mean value of $y = \dfrac{1}{\sqrt{16 - 6x - x^2}}$ between $x = -3$ and $x = 1$. [4]

Examiner
only

(ii)   The region *R* is bounded by the curve *C*, the *x*-axis and the lines $x = -3$ and $x = 1$.
       Find the volume of the solid generated when *R* is rotated through four right-angles
       about the *x*-axis.   [5]

© WJEC CBAC Ltd.   (1305U40-1)   **Turn over.**

(b)   Evaluate the improper integral

$$\int_1^\infty \frac{-8e^{-2x}}{4e^{-2x}-5}\,dx,$$

giving your answer correct to three decimal places.   [3]

© WJEC CBAC Ltd.   (1305U40-1)

BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

(1305U40-1)   **Turn over.**

8. (a) By writing $y = \sinh^{-1}(4x+3)$ as $\sinh y = 4x+3$, show that $\dfrac{dy}{dx} = \dfrac{4}{\sqrt{16x^2 + 24x + 10}}$. [5]

© WJEC CBAC Ltd.   (1305U40-1)

(b)   Show that the graph of $e^{-3x}y = \sinh 2x$ has only one stationary point.   [6]

9. Find the general solution of the equation

$$\sin 6\theta + 2\cos^2\theta = 3\cos 2\theta - \sin 2\theta + 1.$$

[9]

© WJEC CBAC Ltd.   (1305U40-1)

Examiner
only

© WJEC CBAC Ltd.   (1305U40-1)   **Turn over.**

10. The following simultaneous equations are to be solved.

$$\frac{dx}{dt} = 4x + 2y + 6e^{3t}$$

$$\frac{dy}{dt} = 6x + 8y + 15e^{3t}$$

(a) Show that $\dfrac{d^2x}{dt^2} - 12\dfrac{dx}{dt} + 20x = 0$. [5]

Examiner
only

(b) Given that $\dfrac{dx}{dt} = 9$ and $\dfrac{d^2x}{dt^2} = 10$ when $t = 0$, find the particular solution for x in terms of t.

[7]

END OF PAPER

© WJEC CBAC Ltd.   (1305U40-1)   **Turn over.**

Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only

(1305U40-1)

Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examiner only

(1305U40-1)

Turn over.

BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

(1305U40-1)

BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

(1305U40-1) **Turn over.**

BLANK PAGE

**PLEASE DO NOT WRITE
ON THIS PAGE**

(1305U40-1)