



GCE A LEVEL MARKING SCHEME

SUMMER 2022

**A LEVEL (NEW)
MATHEMATICS
UNIT 3 PURE MATHEMATICS B
1300U30-1**

INTRODUCTION

This marking scheme was used by WJEC for the 2022 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

WJEC GCE A LEVEL MATHEMATICS

UNIT 3 PURE MATHEMATICS B

SUMMER 2022 MARK SCHEME

Q	Solution	Mark Notes
1	$6(1 + \tan^2 x) - 8 = \tan x$	M1 use of $\sec^2 x = 1 + \tan^2 x$ Must be seen for M1
	$a\tan^2 x + b\tan x + c = 0$	
	$6\tan^2 x - \tan x - 2 = 0$	
	$(A\tan x + B)(C\tan x + D) = 0$	m1 $AC = a$ and $BD = c, c \neq 0$ oe
	$(3\tan x - 2)(2\tan x + 1) = 0$	
	$\tan x = -\frac{1}{2}, \frac{2}{3}$	A1 cao
	$\tan x = \frac{2}{3}, x = 33.69^\circ, 213.69^\circ$	B1 first 2 correct solutions Condone $0.588^\circ, 3.730^\circ$
	$\tan x = -\frac{1}{2}, x = 153.43^\circ,$	B1 3 rd correct solution Condone 2.678°
	$x = 333.43^\circ$	B1 4th correct solution Condone 5.820°

Notes: If one or two roots obtained for $\tan x$, even if incorrectly obtained, full follow through from these values for B1 B1 B1, provided one +ve and one -ve root. If only one sign obtained, only B1 available for one pair of correct angles.

Do not follow through for sin, cos or anything else.

Ignore all roots outside range $0^\circ \leq x \leq 360^\circ$.

For 5th, 6th, 7th extra root within range, -1 mark each extra root.

If all answers in radians, but radians **not** specified, penalise -1.

Accept all answers correctly rounded to the nearest whole number or better.

Q Solution**Mark Notes**

2(a) $y = x^3 \ln(5x)$

$$\frac{dy}{dx} = 3x^2 \ln(5x) + x^3 \frac{5}{5x}$$

M1 $f(x) \ln(5x) + x^3 g(x)$

M0 if $f(x) = 0$ or 1 or $g(x) = 0$ or 1

A1 $3x^2 \ln(5x)$

A1 $x^3 \frac{5}{5x}$

ISW

$$\frac{dy}{dx} = 3x^2 \ln(5x) + x^2 = x^2(3 \ln(5x) + 1)$$

2(b) $y = (x + \cos 3x)^4$

$$\frac{dy}{dx} = 4(x + \cos 3x)^3(1 - 3\sin 3x)$$

M1 $4(x + \cos 3x)^3 f(x)$

M0 if $f(x) = 1$

A1 $f(x) = (1 - 3\sin 3x)$

Condone absence of brackets

for M1 A0, unless corrected for A1.

ISW

Q Solution**Mark Notes**

3 $OB \left(= \frac{4}{\cos \frac{\pi}{3}} \right) = 8$ or $OA \left(= \frac{4}{\tan 30^\circ} \right) = 4\sqrt{3}$ B1 si ($OA = 6.928\dots$)

Area $OAB = \frac{1}{2} \times 4 \times 8 \sin \frac{\pi}{3}$ M1 Use of $A = \frac{1}{2} \times AB \times OA$
 $= 8\sqrt{3} = 13.856\dots$

Area $BOC = \frac{1}{2} \times 8 \times 8 \times \frac{\pi}{3}$ M1 Use of $A = \frac{1}{2}r^2\theta$
 $= \frac{32\pi}{3} = 33.510\dots$ ft OB, OA

Required area $OABC = 47.37 \text{ (m}^2\text{)}$ A1 cao Must be to 2dp

Q	Solution	Mark Notes
4	$\frac{a}{1-r} = 120$	B1 si
	$\frac{a}{1-4r^2} = 112 \frac{1}{2}$	B1 si
	$120(1-r) = \frac{225}{2}(1-4r^2)$	M1 or elimination of r
	$900r^2 - 240r + 15 = 0$ or $a^2 - 208a + 10800 = 0$	m1 attempt to solve their quadratic equation Implied by correct answers
	$60r^2 - 16r + 1 = 0$	
	$(6r - 1)(10r - 1) = 0$	
	$r = \frac{1}{6}, r = \frac{1}{10}$	A1 One correct pair, cao
	$a = 100, a = 108$	A1 all correct, cao

Q	Solution	Mark Notes
5(a)	$\left(\frac{6x+4}{(x-1)(x+1)(2x+3)} = \right) \frac{A}{(x-1)} + \frac{B}{(x+1)} + \frac{C}{(2x+3)}$	M1 correct form
		Implied by equation below
	$6x + 4 = A(x+1)(2x+3) + B(x-1)(2x+3)$	
	$+ C(x+1)(x-1)$	M1 si correct equation
	Put $x = -1, -2 = B(-2)(1)$	
	$B = 1$	
	Put $x = -\frac{3}{2}, -9 + 4 = C(-\frac{1}{2})(-\frac{5}{2})$	
	$C = -4$	A1 two correct constants
	Put $x = 1, 10 = A(2)(5)$	
	$A = 1$	A1 third constant correct
	$f(x) = \frac{1}{(x-1)} + \frac{1}{(x+1)} - \frac{4}{(2x+3)}$	
5(b)	$\int \frac{3x+2}{(x-1)(x+1)(2x+3)} dx$	
	$= \int \frac{1}{2} \left[\frac{1}{(x-1)} + \frac{1}{(x+1)} - \frac{4}{(2x+3)} \right] dx$	
	$= \frac{1}{2} [\ln x-1 + \ln x+1 - 2\ln 2x+3 (+\ln C)]$	B3 B1 correct int of $\frac{1}{(x-1)}$
		B1 correct int of $\frac{1}{(x+1)}$
		B1 correct int of $\frac{K}{(2x+3)}$
		Condone no modulus signs for B3
		M1 attempt to tidy up into one ln term M0 if extra terms seen
	$= \frac{1}{2} \left[\ln \left \frac{C(x+1)(x-1)}{(2x+3)^2} \right \right] \text{ or } \left[\ln \left \frac{\sqrt{C(x+1)(x-1)}}{(2x+3)} \right \right]$	A1 cao accept $+C$
		A0 if no C . ISW

Q Solution**Mark Notes**

6(a) $T_{12} = 10 + (12 - 1) \times 0.2$

M1 use of $a + (n - 1)d$ Allow $d = 20$ for M1.

Implied by correct answer.

$T_{12} = \text{£}12.20$

A1

6(b) $(954 =) \frac{n}{2} [2 \times 10 + (n - 1) \times 0.2]$

M1 use of $\frac{n}{2} [2a + (n - 1)d]$ Allow $d = 20$ for M1.

$9540 = n[100 + n - 1]$

$n^2 + 99n - 9540 = 0$

m1 equating to 954 and
writing as quadratic
Implied by $n = 60$

$(n - 60)(n + 159) = 0$

$n = 60$

A1 cao Dependent on M1
A0 if $n = -159$ present in final
answer

60 (months)

Q	Solution	Mark Notes
7	$x^2 = 8\sqrt{x}$ or $y = \left(\frac{y^2}{64}\right)^2$ $x^4 = 64x$ or $y^4 = 4096y$	M1 equating y 's
	$x(x^3 - 64) = 0$ or $y(y^3 - 4096) = 0$	A1 oe e.g. $x^{\frac{3}{2}} = 8$
	$x = (0,)$ 4 or $y = (0,)$ 16	A1 si, 0 not required.
	$\text{Area} = \int_0^4 \left(8x^{\frac{1}{2}} - x^2\right) dx$	M1 oe allow $x^2 - 8x^{\frac{1}{2}}$ limits not required
	$\text{Area} = \left[\frac{16}{3}x^{\frac{3}{2}} - \frac{1}{3}x^3\right]_0^4$	A1 one correct term Must be seen
		A1 other correct term
	$\text{Area} = \frac{16}{3} \times 8 - \frac{1}{3} \times 64$	
	$\text{Area} = \frac{64}{3}$	A1 cso A0 if integral gives negative answer, unless corrected without any incorrect statements.

Alternative Solution for last 4 marks

$$\begin{aligned}
 A &= \int_0^4 8x^{\frac{1}{2}} dx && \text{(M1)} \\
 &= \left[\frac{16}{3}x^{\frac{3}{2}}\right]_0^4 && \text{(A1) Must be seen} \\
 &= \frac{16}{3} \times 8 \quad (= \frac{128}{3}) \\
 B &= \int_0^4 x^2 dx && \text{Award M1 if not awarded previously} \\
 &= \left[\frac{1}{3}x^3\right]_0^4 && \text{(A1) Must be seen} \\
 &= \left(\frac{64}{3}\right) \\
 \text{Required area} &= A - B = \frac{64}{3} && \text{(A1)}
 \end{aligned}$$

Note: Answer only, M1 A0 A0 A0

Q Solution

Mark Notes

8 $\frac{2-x}{\sqrt{1+3x}} = (2-x)(1+3x)^{-1/2}$

$$(1+3x)^{-1/2} = (1 + \left(-\frac{1}{2}\right)(3x) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2}(3x)^2 + \dots) \text{B1}$$

$$\text{B1} \quad 1 + \left(-\frac{1}{2}\right)(3x) + \frac{(-\frac{1}{2})(-\frac{3}{2})}{2}(3x)^2$$

$$\frac{2-x}{\sqrt{1+3x}} = (2-x)\left(1 - \frac{3}{2}x + \frac{27}{8}x^2 + \dots\right)$$

$$= 2 - 3x + \frac{27}{4}x^2 - x + \frac{3}{2}x^2 + \dots$$

$$= 2 - 4x + \frac{33}{4}x^2 + \dots$$

B3 B1 each term

Ignore further terms, ISW

Expansion valid for $|3x| < 1$

$$|x| < \frac{1}{3} \quad \text{or} \quad -\frac{1}{3} < x < \frac{1}{3}$$

B1 B1 for $x < \frac{1}{3}$ and $x > -\frac{1}{3}$

B0 anything else

When $x = \frac{1}{22}$,

$$\frac{2 - \frac{1}{22}}{\sqrt{1 + \frac{3}{22}}} \approx 2 - \frac{4}{22} + \frac{33}{4} \left(\frac{1}{22}\right)^2 \quad \text{M1 sub into LHS and RHS}$$

$$\frac{\frac{43}{22}}{\frac{5\sqrt{22}}{22}} = \frac{43}{5\sqrt{22}} \approx \frac{323}{176} \quad \text{or} \quad \frac{43\sqrt{22}}{110} \approx \frac{323}{176}$$

$$\sqrt{22} \approx \frac{7568}{1615} \quad \text{or} \quad \frac{1615}{344} \quad \text{A1 cao}$$

(= 4.686068111..., or 4.694767442..., actual value is 4.69041576...)

Special case for $(1 + 3x)^{1/2}$ used

$$(1 + 3x)^{1/2} = (1 + \left(\frac{1}{2}\right)(3x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2}(3x)^2 + \dots) \quad (B0)$$

(B0)

$$\begin{aligned} \frac{2-x}{\sqrt{1+3x}} &= (2-x)\left(1 + \frac{3}{2}x - \frac{9}{8}x^2 + \dots\right) \\ &= 2 + 3x - \frac{9}{4}x^2 - x - \frac{3}{2}x^2 + \dots \\ &= 2 + 2x - \frac{15}{4}x^2 + \dots \end{aligned} \quad (B3) \quad \text{B1 each term}$$

Ignore further terms, ISW

Expansion valid for $|3x| < 1$

$$|x| < \frac{1}{3} \text{ or } -\frac{1}{3} < x < \frac{1}{3} \quad (B1) \quad \text{B1 for } x < \frac{1}{3} \text{ and } x > -\frac{1}{3}$$

B0 anything else

Correct substitution (M1)

(A0)

Q Solution**Mark Notes**

9(a) $u_1 = \sin\left(\frac{\pi}{2}\right) = 1$

$u_2 = \sin\left(\frac{2\pi}{2}\right) = 0$

$u_3 = \sin\left(\frac{3\pi}{2}\right) = -1$

$u_4 = \sin\left(\frac{4\pi}{2}\right) = 0$

$u_5 = \sin\left(\frac{5\pi}{2}\right) = \sin\left(\frac{\pi}{2}\right) = 1$

Sequence is periodic (with period 4)

B1 All 5 terms

B1 Condone 'Repeats every 4 terms'
or 'Oscillates'

9(b) $u_5 = 17$

B1

$(u_5 = 17), u_4 = 9, u_3 = 5, u_2 = 3, u_1 = 2$

B1

Sequence is increasing.

B1 Accept 'Divergent'

Q	Solution	Mark Notes
10	$\frac{6x^5 - 17x^4 - 5x^3 + 6x^2}{(3x+2)} = \frac{(x^2)(6x^3 - 17x^2 - 5x + 6)}{(3x+2)}$ $= \frac{(x^2)(3x+2)(2x^2 - 7x + 3)}{(3x+2)}$ $= x^2(2x - 1)(x - 3) = 0$ $x = 0(\text{twice}), \frac{1}{2}, 3.$	M1 or removing x^2 from pentic M1 divide by $(3x + 2)$, or realising $(3x + 2)$ is a factor of the cubic and cancelling A1 Sight of $(2x^2 - 7x + 3)$ A1 Must be seen A1 cao A0 if $-\frac{2}{3}$ present
	<u>Note:</u> $(6x^3 - 17x^2 - 5x + 6) = (x - 3)(6x^2 + x - 2)$	
	$(6x^3 - 17x^2 - 5x + 6) = (2x - 1)(3x^2 - 7x - 6)$	

Alternative Solution

10	$\frac{6x^5 - 17x^4 - 5x^3 + 6x^2}{(3x+2)} = \frac{(x^2)(6x^3 - 17x^2 - 5x + 6)}{(3x+2)}$ $= \frac{(x^2)(3x+2)(2x^2 - 7x + 3)}{(3x+2)}$ $= x^2(2x - 1)(x - 3) = 0$ $x = 0 \text{ (twice)}, \frac{1}{2}, 3.$	(M1) or removing x^2 from pentic (M1) any linear factor or divide by $(3x + 2)$ (A1) Sight of $(2x^2 - 7x + 3)$ oe or second factor from factor theorem (A1) $(3x + 2)$ must be cancelled or solution discarded (A1) cao A0 if $-\frac{2}{3}$ present
	<u>Note:</u> $(6x^3 - 17x^2 - 5x + 6) = (x - 3)(6x^2 + x - 2)$	
	$(6x^3 - 17x^2 - 5x + 6) = (2x - 1)(3x^2 - 7x - 6)$	

Q Solution**Mark Notes**

11(a) $9\cos x + 40\sin x = R\cos x \cos \alpha + R\sin x \sin \alpha$

$R\cos \alpha = 9$ and $R\sin \alpha = 40$

M1 implied by correct α if
nothing seen.

M0 for incorrect equations

$R = \sqrt{9^2 + 40^2} = 41$

B1

$\alpha = \tan^{-1}\left(\frac{40}{9}\right) = 77.32^\circ$

A1 accept 1.349 rad, not 1.349
ft R if $\alpha = \sin^{-1}\left(\frac{40}{R}\right) = \cos^{-1}\left(\frac{9}{R}\right)$

$9\cos x + 40\sin x \equiv 41\cos(x - 77.32^\circ)$

11(b) $y = \frac{12}{9\cos x + 40\sin x + 47}$

Maximum y when denominator is minimum,

i.e. when $\cos(x - 77.32^\circ) = -1$

M1 implied by correct max

Max $y\left(= \frac{12}{-41+47}\right) = 2$

A1 ft R

Q Solution**Mark Notes**

12(a) $f(f(p)) = f(0) = 10$

B1

12(b) $2x^2 + 12x + 10 = 0$
 $2(x^2 + 6x + 5) = 0$
 $2(x + 5)(x + 1) = 0$

M1 may be implied by solution

$p = -5, q = -1$

A1 both

12(c) $f(x) = 2[x^2 + 6x + 5]$
 $= 2[(x + 3)^2 - 4]$
 $= 2(x + 3)^2 - 8$
Min point at $(-3, -8)$

M1 condone absence of '2'
A1 cao
B112(d) $f(x)$ is not a one-to-one function
(on its domain).

B1

12(e)(i) Let $y = 2(x + 3)^2 - 8$

$$(x + 3)^2 = \frac{y+8}{2}$$

$$x = -3 \pm \sqrt{\frac{y+8}{2}}$$

since $x \geq -3, x = -3 + \sqrt{\frac{y+8}{2}}$

M1 ft similar form from (c)


A1 Condone $x = -3 + \sqrt{\frac{y+8}{2}}$

A1 Must discard negative root

$$g^{-1}(x) = -3 + \sqrt{\frac{x+8}{2}}$$

A1 interchange x and y , could be done earlier

12(e)(ii)



B1 Correct shape

B1 (10, 0) (0, -1), cao

Q Solution**Mark Notes**

13(a) $f'(x) = 6x^2 + 3$

B1

Hence $f'(x) > 0$ for all x ,i.e. $f(x)$ does not have a stationary point.

E1 oe

e.g. $f'(x) = 0$ has no real rootsdiscriminant = $0^2 - 4(6)(3) < 0$, no real roots

13(b) $f''(x) = 12x$

M1

At point of inflection $f''(x) = 0, x = 0$

m1

 $f'(x) > 0$ when $x < 0$ and when $x > 0$.Therefore, when $x = 0$,

there is a point of inflection.


A1 oe cubic curve no max/min

must have a point of inflection.

OR $x > 0, f''(x) > 0; x < 0, f''(x) < 0$ The point of inflection is $(0, -5)$

B1

13(c)



G1 cubic curve no max/min

ft point in (b) coords not required.

(1,0) not required.

Q Solution**Mark Notes**

14	$I = [\pm \cos x \cdot x^2]_0^\pi - \int_0^\pi \pm \cos x \cdot 2x \, dx$	M1	attempt at parts, 2 terms, at least one term correct. Limits not required
	$I = [-\cos x \cdot x^2]_0^\pi - \int_0^\pi -\cos x \cdot 2x \, dx$	A1	
	$I = [-\cos x \cdot x^2]_0^\pi + [\sin x \cdot 2x]_0^\pi$		
	$- \int_0^\pi 2\sin x \, dx$	A1	correct integration of
			$\int_0^\pi \pm \cos x \cdot 2x \, dx$
	$I = [-\cos x \cdot x^2]_0^\pi + [2\sin x \cdot x]_0^\pi + [2\cos x]_0^\pi$	A1	correct integration of
			$\int_0^\pi \pm \sin x \, dx$
	$I = [2x\sin x + (2 - x^2)\cos x]_0^\pi$		
	$I = \pi^2 + 0 + 2(-1 - 1)$	m1	correct use of correct limits
			Implied by correct answer
	$I = \pi^2 - 4 (= 5.87)$	A1	cao

Note

No marks for answer unsupported by workings.

If integration is incorrect and answer of 5.87 seen with **no working**, m0 A0. If substitution seen m1 is available.

Be careful of use of calculators to obtain correct answer after incorrect integration.

Condone missing dx .

M1A0 only for $I = \left[\sin x \cdot \frac{x^3}{3} \right]_0^\pi - \int_0^\pi \frac{x^3}{3} \cos x \, dx$

Q Solution**Mark Notes**

15(a) $y = \sqrt{16 - x^2}$ OR $A = 2xy$ B1

$A = 2x\sqrt{16 - x^2}$ B1

15(b) $\frac{dA}{dx} = \frac{d}{dx}[2x(16 - x^2)^{1/2}]$ M1 $f(x) (16 - x^2)^{1/2} + 2xg(x)$

M0 if $f(x) = 0$ or 1 or $g(x) = 0$ or 1

Only ft if product with $Bx\sqrt{K - x^2}$

$\frac{dA}{dx} = 2(16 - x^2)^{1/2} + 2x \times \frac{1}{2}(16 - x^2)^{-1/2}(-2x)$ A1A1 one each term, ft (a)

$\frac{dA}{dx} = \frac{4}{(16 - x^2)^{1/2}}[8 - x^2]$

At max, $\frac{dA}{dx} = 0$ m1

$x^2 = 8$ A1 cao

$x = 2\sqrt{2}$ (-ve value inadmissible)

$y = \sqrt{16 - x^2} = \sqrt{16 - 8} = 2\sqrt{2}$ A1 cao accept $y^2 = 8$

therefore $y = x$.

Justification of maximum B1 $\frac{d^2A}{dx^2} = -22$ when $x = 2\sqrt{2}$

OR

$A^2 = 4x^2(16 - x^2) = 64x^2 - 4x^4$

$\frac{dA^2}{dx} = 128x - 16x^3$ (M1A1A1)

At max, $\frac{dA^2}{dx} = 0$ (m1)

$x^2 = 8, x = 2\sqrt{2}$ (-ve value inadmissible) (A1) cao

$y = \sqrt{16 - x^2} = \sqrt{16 - 8} = 2\sqrt{2}$ (A1) cao accept $y^2 = 8$

therefore $y = x$.

Justification of maximum (B1) $\frac{d^2A^2}{dx^2} = -256$ when $x = 2\sqrt{2}$

Q Solution**Mark Notes**

16(a) Where C meets the y -axis,

$$3 - 4t + t^2 = 0 \quad \text{M1}$$

$$(t - 1)(t - 3) = 0$$

$$t = 1, \text{ point is } (0, 9) \quad \text{A1} \quad \text{or } t = 1, 3$$

$$t = 3, \text{ point is } (0, 1) \quad \text{A1} \quad \text{all correct}$$

$$16(b) \quad \frac{dy}{dt} = -2(4 - t) \quad \text{B1}$$

$$\frac{dx}{dt} = -4 + 2t \quad \text{B1}$$

$$\frac{dy}{dx} = \frac{-2(4-t)}{-4+2t} \quad \text{B1} \quad \text{ft their } \frac{dy}{dt} \text{ and } \frac{dx}{dt}$$


Note: May be seen in (a)

$$\text{At stationary point, } \frac{-2(4-t)}{-4+2t} = 0 \quad \text{M1}$$

$$t = 4$$

$$\text{At stationary point, } y = (4 - 4)^2 = 0.$$

Hence the x -axis is a tangent to the curve C . A1



Q Solution**Mark Notes**

17(a) $\cos(\alpha - \beta) + \sin(\alpha + \beta)$

$$\begin{aligned}
 &= \cos\alpha\cos\beta + \sin\alpha\sin\beta + \sin\alpha\cos\beta + \cos\alpha\sin\beta \quad \text{B1} \quad \text{expand } \cos(\alpha - \beta), \sin(\alpha + \beta) \\
 &= \cos\alpha(\cos\beta + \sin\beta) + \sin\alpha(\cos\beta + \sin\beta) \\
 &= (\cos\alpha + \sin\alpha)(\cos\beta + \sin\beta) \quad \text{B1} \quad \text{convincing}
 \end{aligned}$$

OR

$$\begin{aligned}
 &(\cos\alpha + \sin\alpha)(\cos\beta + \sin\beta) \\
 &= \cos\alpha\cos\beta + \cos\alpha\sin\beta + \sin\alpha\cos\beta + \sin\alpha\sin\beta \quad (\text{B1}) \quad \text{remove brackets} \\
 &= \cos\alpha\cos\beta + \sin\alpha\sin\beta + \sin\alpha\cos\beta + \cos\alpha\sin\beta \\
 &= \cos(\alpha - \beta) + \sin(\alpha + \beta) \quad (\text{B1}) \quad \text{convincing}
 \end{aligned}$$

OR

$$\begin{aligned}
 &\cos(\alpha - \beta) + \sin(\alpha + \beta) \\
 &= \cos\alpha\cos\beta + \sin\alpha\sin\beta + \sin\alpha\cos\beta + \cos\alpha\sin\beta \quad (\text{B1}) \quad \text{expand } \cos(\alpha - \beta), \sin(\alpha + \beta) \\
 &(\cos\alpha + \sin\alpha)(\cos\beta + \sin\beta) \\
 &= \cos\alpha\cos\beta + \cos\alpha\sin\beta + \sin\alpha\cos\beta + \sin\alpha\sin\beta \quad (\text{B1}) \quad \text{remove brackets}
 \end{aligned}$$

Hence $\cos(\alpha - \beta) + \sin(\alpha + \beta)$

$$= (\cos\alpha + \sin\alpha)(\cos\beta + \sin\beta)$$

Q Solution**Mark Notes**17(b)(i) Put $\alpha = 4\theta$, $\beta = \theta$

M1

$$\begin{aligned} & \cos(4\theta - \theta) + \sin(4\theta + \theta) \\ &= (\cos 4\theta + \sin 4\theta)(\cos \theta + \sin \theta) \\ & \frac{\cos 3\theta + \sin 5\theta}{\cos 4\theta + \sin 4\theta} = \cos \theta + \sin \theta \end{aligned}$$

A1 convincing

17(b)(ii) When $\theta = \frac{3\pi}{16}$,

$$\cos 4\theta + \sin 4\theta = \cos \frac{3\pi}{4} + \sin \frac{3\pi}{4} = 0$$

So $\frac{\cos 3\theta + \sin 5\theta}{\cos 4\theta + \sin 4\theta}$ is undefined. B1 oe

OR

$$\cos 4\theta + \sin 4\theta \neq 0$$

$$\tan 4\theta \neq -1$$

$$4\theta \neq \frac{3\pi}{4}$$

$$\theta \neq \frac{3\pi}{16} \quad \text{B1}$$

Q Solution**Mark Notes**18(a) Put $u = x + 3$

B1

$$\begin{aligned}
 \int \frac{x^2}{(x+3)^4} dx &= \int \frac{(u-3)^2}{u^4} du & M1 & \text{Allow one slip} \\
 &= \int \frac{u^2 - 6u + 9}{u^4} du & & \\
 &= \int (u^{-2} - 6u^{-3} + 9u^{-4}) du & A1 & \text{integrable form} \\
 & & & \text{ft } (u + 3) \text{ only} \\
 &= \frac{u^{-1}}{-1} - \frac{6u^{-2}}{-2} + \frac{9u^{-3}}{-3} (+C) & A1 & \text{correct integration} \\
 & & & \text{ft } (u + 3) \text{ only} \\
 &= -\frac{1}{u} + \frac{3}{u^2} - \frac{3}{u^3} (+C) & & \\
 &= -\frac{1}{x+3} + \frac{3}{(x+3)^2} - \frac{3}{(x+3)^3} + C & A1 & \text{cao} \text{ Correct expression in terms} \\
 & & & \text{of } x \\
 & & & \text{Must include } + C
 \end{aligned}$$

18(b) $\int_0^1 \frac{x^2}{(x+3)^4} dx = \left[-\frac{1}{x+3} + \frac{3}{(x+3)^2} - \frac{3}{(x+3)^3} \right]_0^1$

M1 correct use of correct limits
ft for equivalent difficulty
for M1 only

$$\begin{aligned}
 &= \left(-\frac{1}{4} + \frac{3}{16} - \frac{3}{64} \right) - \left(-\frac{1}{3} + \frac{1}{3} - \frac{1}{9} \right) & A1 & \text{cao} \\
 &= \frac{1}{576} (= 0.001736) & & \text{No workings, 0 marks}
 \end{aligned}$$

OR

$$\begin{aligned}
 \int_0^1 \frac{x^2}{(x+3)^4} dx &= \left[-\frac{1}{u} + \frac{3}{u^2} - \frac{3}{u^3} \right]_3^4 & & \\
 &= \left(-\frac{1}{4} + \frac{3}{16} - \frac{3}{64} \right) - \left(-\frac{1}{3} + \frac{1}{3} - \frac{1}{9} \right) & (M1) & \text{correct use of correct limits} \\
 & & & \text{ft for equivalent difficulty} \\
 & & & \text{for M1 only} \\
 &= \frac{1}{576} (= 0.001736) & (A1) & \text{cao}
 \end{aligned}$$

No workings, 0 marks