



---

# **GCE A LEVEL MARKING SCHEME**

---

**SUMMER 2018**

**A LEVEL (NEW)  
MATHEMATICS – UNIT 3 PURE MATHEMATICS B  
1300U30-1**

## **INTRODUCTION**

This marking scheme was used by WJEC for the 2018 examination. It was finalised after detailed discussion at examiners' conferences by all the examiners involved in the assessment. The conference was held shortly after the paper was taken so that reference could be made to the full range of candidates' responses, with photocopied scripts forming the basis of discussion. The aim of the conference was to ensure that the marking scheme was interpreted and applied in the same way by all examiners.

It is hoped that this information will be of assistance to centres but it is recognised at the same time that, without the benefit of participation in the examiners' conference, teachers may have different views on certain matters of detail or interpretation.

WJEC regrets that it cannot enter into any discussion or correspondence about this marking scheme.

## GCE Mathematics – A2 Unit 3 Pure Mathematics B

## SUMMER 2018 MARK SCHEME

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
1	Either $2x + 1 = 3(x - 2)$	M1	Attempt to equate both sides +ve
	$2x + 1 = 3x - 6$		
	$x = 7$	A1	
	OR $2x + 1 = -3(x - 2)$	m1	
	$2x + 1 = -3x + 6$		
	$x = 1$	A1	
Or			
	$(2x + 1)^2 = 9(x - 2)^2$	(M1)	
	$5x^2 - 40x + 35 = 0$	(A1)	any correct equation
	$x^2 - 8x + 7 = 0$		
	$(x - 7)(x - 1) = 0$	(m1)	oe
	$x = 1, 7$	(A1)	both solutions

If considering:

$x < -1/2$  (both sides negative),

$-1/2 \leq x < 2$  (LHS negative, RHS positive),

$x \geq 2$  (both sides positive),


give M1, then m1 if all values considered, A1 for 1 and A1 for 7, extra solution/s -1 however many.

Q	Solution	Mark	Notes
2(a)	$s = r\theta$	M1	used
	$5 = 4\theta$		
	$\theta = 1.25^\circ$	A1	condone $71.62^\circ$ , $5.033$
2(b)	Area of sector $OAB = \frac{1}{2} \times r^2\theta$	M1	used
	Area of sector $OAB = \frac{1}{2} \times 4^2 \times 1.25$		
	Area of sector $OAB = 10 \text{ (cm}^2\text{)}$	A1	ft $\theta$ , accept $40.27$

**Q Solution**

**Mark Notes**

3(a)




B1 correct shape (hill) axes required

B1  $(-2, 18)$  as max

B1  $(-5, 0), (1, 0)$

3(b)



B1 correct shape (cup) axes required

B1  $(1, -4)$  as min

B1  $(0, -3)$

Q	Solution	Mark	Notes
4	$2\tan^2\theta + 2\tan\theta - (1 + \tan^2\theta) = 2$ $\tan^2\theta + 2\tan\theta - 3 = 0$ $(\tan\theta - 1)(\tan\theta + 3) = 0$ $\tan\theta = 1, -3$ $\theta = 45^\circ, 225^\circ$ $\theta = 108.43^\circ, 288.43^\circ$	M1	oe si
		m1	$(\tan\theta + 1)(\tan\theta - 3)$
		A1	cao
		B1	ft $\tan\theta$
		B1	

Ignore all roots outside range. For each branch, award B0 if extra root/s present.

2+ve roots ft for B1

2-ve roots ft for B1

Q	Solution	Mark	Notes
5(a)	$\frac{3x}{(x-1)(x-4)^2} = \frac{A}{(x-1)} + \frac{B}{(x-4)} + \frac{C}{(x-4)^2}$ $3x = A(x-4)^2 + B(x-1)(x-4) + C(x-1)$ $x=4, 12 = 3C, C = 4$ $x=1, 3 = 9A, A = \frac{1}{3}$ $\text{coefficient } x^2, 0 = A + B, B = -\frac{1}{3}$	M1 m1	RHS over common denominator compare coefficients or substitute values.
		A1	all 3 values correct
5(b)	$I = \int_5^7 \frac{1}{3(x-1)} - \frac{1}{3(x-4)} + \frac{4}{(x-4)^2} dx$ $I = \left[ \frac{1}{3} \ln x-1  - \frac{1}{3} \ln x-4  - \frac{4}{(x-4)} \right]_5^7$ $I = \left( \frac{1}{3} \ln 6 - \frac{1}{3} \ln 3 - \frac{4}{3} \right) - \left( \frac{1}{3} \ln 4 - 4 \right)$ $I = \frac{1}{3} (8 - \ln 2) = 2.436 \text{ (3d.p. required)}$	M1	attempt to integrate partial  Fractions
		A1	ft any one term correct
		A1	ft all correct integration
		m1	correct use of correct limits
		A1	cao

Q	Solution	Mark	Notes
6	$(1 - 4x)^{-\frac{1}{2}} = 1 + \left(-\frac{1}{2}\right)(-4) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2!}(-4x)^2 + \dots$	B1	2 correct unsimplified terms
	$= 1 + 2x + 6x^2 + \dots$	B1	all simplified terms correct
	Expansion is valid when $ 4x  < 1$		
	Expansion is valid when $ x  < \frac{1}{4}$	B1	oe
	When $x = \frac{1}{13}$ ,		
	$\frac{1}{\sqrt{1 - \frac{4}{13}}} \cong 1 + 2 \times \frac{1}{13} + 6 \times \left(\frac{1}{13}\right)^2$	M1	attempt to substitute both sides.
	$\frac{\sqrt{13}}{3} \cong \frac{201}{169}$		
	$\sqrt{13} \cong \frac{603}{169} \quad \text{or} \quad \frac{2197}{603}$	A1	



Q	Solution	Mark	Notes
7	$\sin x \cong x, \cos x \cong 1 - \frac{1}{2}x^2$	M1	used
	$x + 1 - \frac{1}{2}x^2 = \frac{1}{2}$		
	$x^2 - 2x - 1 = 0$	A1	oe
	$x = \frac{2 \pm \sqrt{2^2 + 4}}{2}$		
	$x = 1 - \sqrt{2} (= -0.4142)$	A1	cao

Q	Solution	Mark	Notes
8	$a + 6d = 71$	B1	
	$\frac{7}{2}(2a + 6d) = 329$	B1	
	$a + 3d = 47$		
	$a + 6d = 71$		
	$3d = 24$		
	$d = 8$	B1	
	$a = 23$	B1	
	The numbers are 23, 31, 39, 47, 55, 63, 71.	B1	

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
9(a)	<p>The sum to <math>n</math> terms of a series is <math>S_n = \frac{a(1-r^n)}{(1-r)}</math></p> <p>The sum to infinity is <math>\lim_{n \rightarrow \infty} S_n</math>.</p> <p>This only converges if <math>\lim_{n \rightarrow \infty} r^n</math> converges.</p> <p>Hence the sum to infinity of a GP</p> <p>only converges if <math> r  &lt; 1</math></p>	B1	oe eg. terms increasing
9(b)	<p>For <math>W</math>, the <math>k^{\text{th}}</math> term <math>T_k</math> is <math>(2r^{k-1})^2 = 4r^{2k-2}</math>.</p> <p>The <math>(k+1)^{\text{th}}</math> term is <math>(2r^k)^2 = 4r^{2k}</math>.</p> $\frac{T_{k+1}}{T_k} = \frac{4r^{2k}}{4r^{2k-2}} = r^2 \text{ for all values of } k.$ <p>Therefore <math>W</math> is a GP.</p> <p>For <math>V</math>, 1<sup>st</sup> term is 2, common ratio <math>r</math></p> <p>For <math>W</math>, 1<sup>st</sup> term is 4, common ratio <math>r^2</math></p> $S_V = \frac{2}{1-r}, S_W = \frac{4}{(1-r^2)}$ $S_W = 3S_V$ $\frac{4}{(1-r^2)} = 3\left(\frac{2}{1-r}\right)$ $\frac{4}{(1+r)(1-r)} = 3\left(\frac{2}{1-r}\right)$ $\frac{2}{(1+r)} = 3 \quad (r \neq 1)$ $2 = 3 + 3r$ $r = -\frac{1}{3}$	B1 B1 B1 M1	common ratio $r^2$ si either correct used
		A1	oe. ft $W$ eg quadratic equation
		A1	cao

Q	Solution	Mark Notes
9(c)	Total savings $T = 5000[(1.03) + (1.03)^2 + (1.03)^3 + \dots + (1.03)^{20}]$	M1 si
	$T = \frac{5000(1.03)(1 - 1.03^{20})}{1 - 1.03}$	m1
	$T = 138382 \text{ (£)}$	A1

<b>Q</b>	<b>Solution</b>	<b>Mark</b>	<b>Notes</b>
10(a)	$x = 2\cos^2\theta - 1$	M1	$\cos 2\theta = 2\cos^2\theta - 1$
	$x = 2y^2 - 1$	A1	isw
	$2y^2 = x + 1$		
10(b)	$\cos 2\theta - \cos\theta + 1 = 0$	M1	
	$2\cos^2\theta - 1 - \cos\theta + 1 = 0$	m1	
	$\cos\theta(2\cos\theta - 1) = 0$	A1	si
	$\cos\theta = \frac{1}{2}, 0$	A1	
	$\theta = \frac{\pi}{3}, \frac{\pi}{2}$		answer given
	Co-ordinates are $P(-\frac{1}{2}, \frac{1}{2})$ and $Q(-1, 0)$	B1	

Alternative solution

Using $x - y + 1 = 0$ and $2y^2 = x + 1$	(M1)	attempt to solve simultaneously
$2(x + 1)^2 = x + 1, 2x^2 + 3x + 1 = 0$	(m1)	eliminate one variable
$(2x + 1)(x + 1) = 0, \quad x = -\frac{1}{2}, -1$		
$y = \frac{1}{2}, y = 0,$	(A1)	
$\cos\theta = \frac{1}{2}, \cos\theta = 0$	(A1)	si both
$\theta = \frac{\pi}{3}, \theta = \frac{\pi}{2},$		answer given
$P(-\frac{1}{2}, \frac{1}{2}), Q(-1, 0)$	(B1)	

Accept  $P(-\frac{1}{2}, \frac{1}{2}), Q(-1, 0)$  (B1), verification for  $P$  M1 A1, verification for  $Q$  m1 A1.

Q	Solution	Mark	Notes
10(c)	$\frac{dy}{dx} = \frac{dy}{d\theta} / \frac{dx}{d\theta}$	M1	used
	$\frac{dx}{d\theta} = -2\sin 2\theta$	A1	
	$\frac{dy}{d\theta} = -\sin\theta$	A1	
	$\frac{dy}{dx} = \frac{-\sin\theta}{-2\sin 2\theta} = \left(\frac{1}{4\cos\theta}\right)$		
	Grad of tgt at $P = \frac{1}{2}$	A1	
	Equ of tgt at $P$ is $y - \frac{1}{2} = \frac{1}{2}\left(x + \frac{1}{2}\right)$	A1	
	Equ of tgt at $P$ is $4y = 2x + 3$		
	Grad of tgt at $Q$ is undefined.		
	Equ of tgt at $Q$ is $x = -1$	A1	
	Point of intersection is $\left(-1, \frac{1}{4}\right)$	A1	
OR (first 3 marks)			
	$2 \times 2y \frac{dy}{dx} = 1$	(M1)	attempt implicit differentiation
		(A1)	$2y \frac{dy}{dx}$
		(A1)	all correct.

Q	Solution	Mark	Notes
11	<p>Suppose that <math>\sin x + \cos x \geq 1</math> is not true. Then there exists an <math>x</math> in the given domain</p> <p>for which <math>\sin x + \cos x &lt; 1</math></p> <p><math>(\sin x + \cos x)^2 &lt; 1^2</math></p> <p><math>\sin^2 x + 2\sin x \cos x + \cos^2 x &lt; 1^2</math></p> <p><math>1 + 2\sin x \cos x &lt; 1</math></p> <p><math>\sin x \cos x &lt; 0</math></p> <p>As <math>\sin x \geq 0</math> and <math>\cos x \geq 0</math>, this is impossible.</p> <p>Hence <math>\sin x + \cos x &lt; 1</math> cannot be true,</p> <p>hence <math>\sin x + \cos x \geq 1</math> for <math>0 \leq x \leq \frac{\pi}{2}</math>.</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>A1</p> <p>A1</p>	<p>or <math>\sin 2x &lt; 0</math></p> <p>cso</p>

**Q Solution****Mark Notes**

12(a)(i)  $f$  has an inverse function if and only if

$f$  is both one-to-one (and onto).

B1

12(a)(ii)  $ff^{-1}(x) = x$

B1

12(b)(i)  $g^{-1}$  exists if the domain of  $g$  is  $[0, \infty)$

B1

or  $(-\infty, 0]$  or subset of one of these

12(b)(ii) Let  $y = e^x + 1$

M1

$$e^x = y - 1$$

$$x = \ln(y - 1)$$

$$h^{-1}(x) = \ln(x - 1)$$

A1

or

$$h(x) = e^x + 1$$


$$x = e^{h^{-1}(x)} + 1$$

(M1)

$$e^{h^{-1}(x)} = x - 1$$

$$h^{-1}(x) = \ln(x - 1)$$

(A1)



G1  $h(x)$  with  $y=1$  as asymptote

G1  $h^{-1}(x)$  with  $x=1$  as asymptote

G1  $(0, 2), (2, 0)$



Q	Solution	Mark Notes
12(b)(iii)	$gh(x) = g(e^x + 1)$	M1 accept $(h(x))^2 - 1$
	$gh(x) = (e^x + 1)^2 - 1$	
	$gh(x) = e^{2x} + 2e^x$ or $e^x(e^x + 2)$	A1

Q	Solution	Mark	Notes
13(a)	$R\sin(\theta - \alpha) \equiv 8\sin\theta - 15\cos\theta$ $R\sin\theta\cos\alpha - R\cos\theta\sin\alpha \equiv 8\sin\theta - 15\cos\theta$ $R\cos\alpha = 8$ $R\sin\alpha = 15$ $R = \sqrt{8^2 + 15^2} = 17$ $\alpha = \tan^{-1}\left(\frac{15}{8}\right) = 61.93^\circ$	M1	oe si    B1  A1
13(b)	$17\sin(\theta - 61.93^\circ) = 7$ $\theta - 61.93^\circ = \sin^{-1}\left(\frac{7}{17}\right)$ $\theta - 61.93^\circ = 24.32^\circ, 155.68^\circ$ $\theta = 86.24^\circ$ $\theta = 217.61^\circ$	M1	   A1    cao accept 86.25 A1    cao
13(c)	$\frac{1}{8\sin\theta - 15\cos\theta + 23} = \frac{1}{17\sin(\theta - 61.93^\circ) + 23}$ <p>Greatest value = <math>\frac{1}{6}</math></p> <p>Least value = <math>\frac{1}{40}</math></p>	B1	B1

Q	Solution	Mark	Notes
14(a)	Use integration by parts	M1	
	$I = \left[ \frac{x^4}{4} \ln x \right]_1^2 - \int_1^2 \frac{x^4}{4} \times \frac{1}{x} dx$	A1	1 <sup>st</sup> term
		A1	2 <sup>nd</sup> term
	$I = \left[ \frac{x^4}{4} \ln x \right]_1^2 - \left[ \frac{x^4}{16} \right]_1^2$	A1	2 <sup>nd</sup> bracket
	$I = (4 \ln 2) - \left( 1 - \frac{1}{16} \right)$	m1	correct use of limits
	$I = 4 \ln 2 - \frac{15}{16} = 1.835$	A1	cao
14(b)	Let $x = 2 \sin \theta$	M1	
	$dx = 2 \cos \theta d\theta$		
	$\sqrt{4 - x^2} = \sqrt{4 - 4 \sin^2 \theta} = 2 \cos \theta$		
	$x = 0, \theta = 0; x = 1, \theta = \frac{\pi}{6}$	A1	or 0, 1 if $x$ used.
	$I = \int_0^{\frac{\pi}{6}} \frac{2 + 2 \sin \theta}{2 \cos \theta} 2 \cos \theta dx$	A1	correct integrand
	$I = 2 \int_0^{\frac{\pi}{6}} 1 + \sin \theta d\theta$		
	$I = 2 \left[ \theta - \cos \theta \right]_0^{\frac{\pi}{6}}$	A1	correct integration
	$I = 2 \left( \frac{\pi}{6} - \frac{\sqrt{3}}{2} + 1 \right)$	m1	correct use of limits
	$I = \frac{\pi}{3} + 2 - \sqrt{3} = 1.315$	A1	cao

**Q Solution****Mark Notes**

14(b)

Alternative solution

Let  $x = 2\cos\theta$

(M1)

$$dx = -2\sin\theta d\theta$$

$$\sqrt{4-x^2} = \sqrt{4-4\cos^2\theta}$$

$$= 2\sqrt{1-\cos^2\theta} = 2\sin\theta$$

$$x = 0, \theta = \frac{\pi}{2}; x = 1, \theta = \frac{\pi}{3}$$

(A1) or 0, 1 if x used.

$$I = \int_{\frac{\pi}{2}}^{\frac{\pi}{3}} \frac{2+2\cos\theta}{2\sin\theta} (-2\sin\theta) d\theta$$

(A1) correct integrand

$$I = -2 \int_{\frac{\pi}{2}}^{\frac{\pi}{3}} 1 + \cos\theta d\theta$$

$$I = -2[\theta + \sin\theta]_{\frac{\pi}{2}}^{\frac{\pi}{3}}$$

(A1) correct integration

$$I = -2\left[\left(\frac{\pi}{3} + \frac{\sqrt{3}}{2}\right) - \left(\frac{\pi}{2} + 1\right)\right]$$

(m1) correct use of limits

$$I = -2\left(-\frac{\pi}{6} + \frac{\sqrt{3}}{2} - 1\right)$$

$$I = \frac{\pi}{3} + 2 - \sqrt{3} = 1.315$$

(A1) cao

Q	Solution	Mark	Notes
15	$\int \frac{2dy}{5-2y} = \int dx$	M1	separate variable
	$-\ln 5-2y  = x (+ C)$	A1	5-2y not separated. correct integration
	When $x = 0, y = 1$		
	$-\ln 3  = C$	m1	use of boundary conditions
	$\ln 5-2y  - \ln 3 = -x$		
	$\frac{5-2y}{3} = e^{-x}$	m1	inversion
	$y = \frac{1}{2}(5 - 3e^{-x})$	A1	cao any correct expression

**Q Solution****Mark Notes**

16(a)(i)

$$\frac{dy}{dx} = e^{3\tan x} \times 3\sec^2 x$$

M1 chain rule  $e^{3\tan x} f(x)$ 

$$\frac{dy}{dx} = 3\sec^2 x e^{3\tan x}$$

A1  $f(x) = 3\sec^2 x$ 

16(a)(ii)

$$\frac{dy}{dx} = \frac{x^2(\cos 2x \times 2) - \sin 2x(2x)}{x^4}$$

M1 use of quotient rule oe

$$\frac{x^2 f(x) - \sin 2x g(x)}{x^4}$$

$$\frac{dy}{dx} = \frac{2x \cos 2x - 2 \sin 2x}{x^3}$$

A1  $f(x) = 2\cos 2x$  or  $g(x) = 2x$ 

A1 cao

Alternative solution

$$y = x^{-2} \sin 2x$$

$$\frac{dy}{dx} = -2x^{-3} \sin 2x + 2x^{-2} \cos 2x$$

(M1) use of product rule

$$f(x) \sin 2x + x^{-2} g(x)$$


(A1)  $f(x) = -2x^{-3}$  or  $g(x) = 2\cos 2x$ 

(A1) cao

Q	Solution	Mark	Notes
16(b)	$3x^2 \frac{dy}{dx} + 6xy + 2y \frac{dy}{dx} - 5 = 0$	B1	$3x^2 \frac{dy}{dx} + 6xy$
		B1	$2y \frac{dy}{dx}$
		B1	- 5
	$(3x^2 + 2y) \frac{dy}{dx} = 5 - 6xy$		
	$\frac{dy}{dx} = \frac{5 - 12}{3 + 4} = -1$	B1	cao
	Use of gradient = $-1 / \frac{dy}{dx}$	M1	
	Equation of normal is $y - 2 = 1(x - 1)$		
	Equation of normal is $y = x + 1$	A1	correct equation any form

**Q Solution****Mark Notes**

17



G1 both graphs

The two graphs intersect only once.

B1

(Root is between 0 and  $\pi/2$ .)

Using Newton-Raphson Method

$$f(x) = x - 1 - \cos x$$

$$f'(x) = 1 + \sin x$$

B1 or  $-1 - \sin x$ 

$$x_{n+1} = x_n - \frac{x_n - 1 - \cos x_n}{1 + \sin x_n}$$

M1

$$x_0 = 1$$

$$x_1 = 1.293408$$

A1 si

$$x_2 = 1.283436$$

$$x_3 = 1.283429$$

$$x_4 = 1.283429$$

Root is 1.28 (correct to 2 d. p.)

A1