

Cambridge
International
AS & A Level

Cambridge International Examinations
Cambridge International Advanced Subsidiary and Advanced Level

MATHEMATICS

9709/63

Paper 6

May/June 2016

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.

This document consists of **6** printed pages.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9709	63

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9709	63

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only – often written by a ‘fortuitous’ answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR –2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9709	63

Qu	Answer	Marks	Guidance																
1 (i)	<table border="1"> <thead> <tr> <th></th> <th>Wears specs</th> <th>Not wears specs</th> <th>Total</th> </tr> </thead> <tbody> <tr> <th>RH</th> <td>6</td> <td>19</td> <td>25</td> </tr> <tr> <th>Not RH</th> <td>2</td> <td>3</td> <td>5</td> </tr> <tr> <th>Total</th> <td>8</td> <td>22</td> <td></td> </tr> </tbody> </table>		Wears specs	Not wears specs	Total	RH	6	19	25	Not RH	2	3	5	Total	8	22		B1	One correct row or col including total other than the Total row/column
	Wears specs	Not wears specs	Total																
RH	6	19	25																
Not RH	2	3	5																
Total	8	22																	
(ii)	$P(X) = 25/30, P(Y) = 8/30$ $P(X) \times P(Y) = 25/30 \times 8/30 = 200/900 = 2/9$ $P(X \cap Y) = 6/30 = 1/5 \neq P(X) \times P(Y)$ Not independent	B1 [2] M1 M1 A1 [3]	All correct $P(X)$ or $P(Y)$ from their table or correct from question (denom 30) oe Comparing their $P(X) \times P(Y)$ (values substituted) with their evaluated $P(X \cap Y)$ – not $P(X) \times P(Y)$																
2 (i)		B1 B1 B1 [3]	Labels 'time' and 'seconds', 'boys' and 'girls' on correct plots and scaled line One box and whisker all correct on graph paper – ignore boy or girl label Second box and whisker all correct (on graph paper and ignore boy/girl label) on SAME scaled line.																
(ii)	girls smaller range or IQ range than boys /girls less spread out oe girls generally quicker than boys or girls median < boys median (not mean) oe boys almost symmetrical, girls +vely skewed oe	B1 B1 [2]	Any 2 comments – MUST be a comparison																
3 (i)	$P(0) = 6/36, P(1) = 10/36, P(2) = 8/36$ $P(3) = 6/36, P(4) = 4/36, P(5) = 2/36$	B1 B1 M1 A1 [4]	Table oe seen with 0, 1, 2, 3, 4, 5 (6 if $P(6) = 0$) Any three probs correct $\Sigma p = 1$ and at least 3 outcomes All probs correct																
(ii)	mean score = $(0 \times 6 + 1 \times 10 + 16 + 18 + 16 + 10) / 36$ $= 70/36$ (35/18, 1.94)	M1 A1 [2]	Using Σxp (unsimplified) on its own – condone Σp not = 1																

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9709	63

Qu	Answer	Marks	Guidance
4	(i) 1845/9 (= 205) $c = 2205 - 205 = 2000$ OR $\Sigma x = 2205 \times 9 (= 19845)$ $\Sigma x - \Sigma c = 1845$ $\Sigma c = 19845 - 1845 = 18000$ $c = 2000$	M1 A1 M1 A1 [2]	Accept (1845± anything)/ 9 For 2205 × 9 seen
	(ii) $\text{var} = \frac{477450}{9} - 205^2$ $= 11025$ OR $\text{var} = \frac{43857450}{9} - 2205^2$ $= 11025$	M1 A1 M1 A1 [2]	For $\frac{477450}{9} - (\text{their coded mean})^2$ For their $\Sigma x^2/9 - 2205^2$ where Σx^2 is obtained from expanding $\Sigma(x - c)^2$ with $2c\Sigma x$ seen
	(iii) new total = 2120.5 × 10 = 21205 new price = 21205 – 19845 $= 1360$	M1 A1 [2]	Attempt at new total
5	(i) $z = 1.015$ $1.015 = \frac{70 - 69}{\sigma}$ $\sigma = 0.985 (200/203)$	B1 M1 A1 [3]	Accept z between ±1.01 and 1.02 Standardising
	(ii) $58 + 9 = 67$ $P(> 67) = P\left(z > \frac{67 - 69}{0.9852}\right)$ $= P(z > -2.03)$ $= 0.9788$ 300×0.9788 $= 293.6$ so 293	M1 M1 M1 M1 A1 [5]	58 + 9 seen or implied (or 69-58 or 69-9) Standardising ± z no cc allow their sd (must be +ve) Alt. 1 69-58 = 11, $P(>9) = P\left(z > \frac{9 - 11}{0.9852}\right)$ Alt. 2 69-9 = 60, $P(>58) = P\left(z > \frac{58 - 60}{0.9852}\right)$ Correct prob area Multiply their prob (from use of tables) by 300 – accept 293 or 294 from fully correct working

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – May/June 2016	9709	63

Qu	Answer	Marks	Guidance
6 (i)	7560 ways	B1 [1]	
(ii)	RxxxxxxxG in $\frac{7!}{4!}$	B1	7! alone seen in num or 4! alone in denom Must be in a fraction. $\frac{7 \times 2}{4 \times 2}$ gets full marks
	= 210 ways	B1 [2]	
(iii)	eg EEEEExxxx in $\frac{6!}{2!}$	B1	6! or 5! $\times 6$ seen in numerator or on own Can be 6! $\times k$ but not 6! $\pm k$
	= 360 ways	B1 [2]	
(iv)	1 R eg RVG or RVN or RGN = 3	B1 [1]	
(v)	no Rs eg VGN or 3C3 ways = 1 2 Rs eg RRV or 3C1 ways = 3	M1	Summing at least 2 options for R
	Total = 7	A1 A1 [3]	Correct outcome for no Rs or 2 Rs – evaluated
7 (i)	${}^{12}C_8 (0.65)^8(0.35)^4 + {}^{12}C_9 (0.65)^9(0.35)^3 + {}^{12}C_{10} (0.65)^{10}(0.35)^2$	M1	Bin term with ${}^{12}C_r p^r (1-p)^{12-r}$ seen $r \neq 0$ any $p < 1$
	= 0.541	M1 A1 [3]	Summing 2 or 3 bin probs $p = 0.65$ or 0.35 , $n = 12$
(ii)	$P(\overline{RRRR}) = 0.35 \times 0.35 \times 0.35 \times 0.65$	M1	Mult 4 probs either $(0.35)^3(0.65)$ or $(0.65)^3(0.35)$
	= 0.0279	A1 [2]	
(iii)	$P(7) = 0.2039$ (unsimplified)	B1	${}^{12}C_7 (0.65)^7(0.35)^5$
	Mean = 250×0.2039 (= 50.9798) Var = $250 \times 0.2039 \times (1 - 0.2039)$ (= 40.5851)	B1	Correct unsimplified np and npq using ‘their 0.2039’ but not 0.65 or 0.35
	$P(> 54) = P\left(\frac{54.5 - 50.9798}{\sqrt{40.5851}}\right)$	M1	Standardising need sq rt – must be from working with 54
	= $P(z > 0.5526)$	M1	cc either 53.5 or 54.5
	= $1 - \Phi(0.5526) = 1 - 0.7098$	M1	correct area < 0.5 i.e. $1 - \Phi$ - must be from working with 54
	= 0.290	A1 [6]	