

Question	Scheme	Marks	AOs
1(a)	$V = \pi r^2 h = 355 \Rightarrow h = \frac{355}{\pi r^2}$ $\left( \text{or } rh = \frac{355}{\pi r} \text{ or } \pi rh = \frac{355}{r} \right)$	B1	1.1b
	$C = 0.04(\pi r^2 + 2\pi rh) + 0.09(\pi r^2)$	M1	3.4
	$C = 0.13\pi r^2 + 0.08\pi rh = 0.13\pi r^2 + 0.08\pi r \left( \frac{355}{\pi r^2} \right)$	dM1	2.1
	$C = 0.13\pi r^2 + \frac{28.4}{r} *$	A1*	1.1b
		(4)	
(b)	$\frac{dC}{dr} = 0.26\pi r - \frac{28.4}{r^2}$	M1 A1	3.4 1.1b
	$\frac{dC}{dr} = 0 \Rightarrow r^3 = \frac{28.4}{0.26\pi} \Rightarrow r = \dots$	M1	1.1b
	$r = \sqrt[3]{\frac{1420}{13\pi}} = 3.26\dots$	A1	1.1b
		(4)	
(c)	$\left( \frac{d^2C}{dr^2} = \right) 0.26\pi + \frac{56.8}{r^3} = 0.26\pi + \frac{56.8}{"3.26" ^3}$	M1	1.1b
	$\left( \frac{d^2C}{dr^2} = \right) (2.45\dots) > 0 \text{ Hence minimum (cost)}$	A1	2.4
		(2)	
(d)	$C = 0.13\pi ("3.26")^2 + \frac{28.4}{"3.26"}$	M1	3.4
	$(C =) 13$	A1	1.1b
		(2)	

(12 marks)

## Notes

(a)

B1: Correct expression for  $h$  or  $rh$  or  $\pi rh$  in terms of  $r$ . This may be implied by their later substitution.

M1: Scored for the sum of the three terms of the form  $0.04\dots r^2$ ,  $0.09\dots r^2$  and  $0.04 \times \dots rh$   
 The  $0.04 \times \dots rh$  may be implied by eg  $0.04 \times \dots r \times \frac{355}{\pi r^2}$  if  $h$  has already been replaced

dM1: Substitutes  $h$  or  $rh$  or  $\pi rh$  into their equation for  $C$  which must be of an allowable form (see above) to obtain an equation connecting  $C$  and  $r$ .  
 It is dependent on a correct expression for  $h$  or  $rh$  or  $\pi rh$  in terms of  $r$

A1\*: Achieves given answer with no errors. Allow Cost instead of  $C$  but they cannot just have an expression.

As a minimum you must see

- the separate equation for volume
- the two costs for the top and bottom separate before combining
- a substitution before seeing the  $\frac{28.4}{r}$  term

$$\text{Eg } 355 = \pi r^2 h \text{ and } C = 0.04\pi r^2 + 0.09\pi r^2 + 0.04 \times 2\pi r h = 0.13\pi r^2 + 0.08\pi \times \left(\frac{355}{\pi r}\right)$$

(b)

M1: Differentiates to obtain at least  $r^{-1} \rightarrow r^{-2}$

A1: Correct derivative.

M1: Sets  $\frac{dC}{dr} = 0$  and solves for  $r$ . There must have been some attempt at differentiation of the equation for  $C$  ( $\dots r^2 \rightarrow \dots r$  or  $\dots r^{-1} \rightarrow \dots r^{-2}$ ) Do not be concerned with the mechanics of their rearrangement and do not withhold this mark if their solution for  $r$  is negative

A1: Correct value for  $r$ . Allow exact value or awrt 3.26

(c)

M1: Finds  $\frac{d^2C}{dr^2}$  at their (positive)  $r$  or considers the sign of  $\frac{d^2C}{dr^2}$ .

This mark can be scored as long as their second derivative is of the form  $A + \frac{B}{r^3}$  where  $A$  and  $B$  are non zero

A1: Requires

- A correct  $\frac{d^2C}{dr^2}$
- Either
  - deduces  $\frac{d^2C}{dr^2} > 0$  for  $r > 0$  (without evaluating). There must be some minimal explanation as to why it is positive.
  - substitute their positive  $r$  into  $\frac{d^2C}{dr^2}$  without evaluating and deduces  $\frac{d^2C}{dr^2} > 0$  for  $r > 0$
  - evaluate  $\frac{d^2C}{dr^2}$  (which must be awrt 2.5) and deduces  $\frac{d^2C}{dr^2} > 0$  for  $r > 0$

(d)

M1: Uses the model and their positive  $r$  found in (b) to find the minimum cost. Their  $r$  embedded in the expression is sufficient. May be seen in (b) but must be used in (d).

A1: ( $C =$ ) 13 ignore units

Question	Scheme	Marks	AOs
<b>2 (a)</b>	$\left\{ \frac{dy}{dx} = \right\} 2x^2 - 7x - 4$	M1 A1	1.1b 1.1b
		<b>(2)</b>	
<b>(b)</b>	Attempts to solve $\left\{ \frac{dy}{dx} = \right\} 2x^2 - 7x - 4 \dots 0$ e.g., $(2x+1)(x-4) = 0$ leading to $x = \dots$ and $x = \dots$	M1	1.1b
	Correct critical values $x = -\frac{1}{2}, 4$	A1	1.1b
	Chooses inside region for their critical values	dM1	1.1b
	Accept either $-\frac{1}{2} < x < 4$ <b>or</b> $-\frac{1}{2} \leq x \leq 4$	A1	1.1b
		<b>(4)</b>	

**(6 marks)****Notes:****(a)**

**M1:** Decreases the power of  $x$  by one for at least one of their terms. Look for  $x^n \rightarrow \dots x^{n-1}$   
Allow for  $5 \rightarrow 0$

**A1:**  $\left\{ \frac{dy}{dx} = \right\} 2x^2 - 7x - 4$

**(b)**

**M1:** Sets their  $\frac{dy}{dx} \dots 0$  where  $\dots$  may be an equality or an inequality and proceeds to find two values for  $x$  from a 3TQ using the usual rules. This may be implied by their critical values.

**A1:** Correct critical values  $x \dots -\frac{1}{2}, 4$

These may come directly from a calculator and might only be seen on a sketch.

**dM1:** Chooses the inside region for their critical values.

**A1:** Accept either  $-\frac{1}{2} < x < 4$  **or**  $-\frac{1}{2} \leq x \leq 4$  but not, e.g.,  $-\frac{1}{2} < x \leq 4$

Condone, e.g.,  $x > -\frac{1}{2}, x < 4$  **or**  $x > -\frac{1}{2}$  and  $x < 4$  **or**  $\left\{ x : x > -\frac{1}{2} \right\} \cap \left\{ x : x < 4 \right\}$

**or**  $x \in \left( -\frac{1}{2}, 4 \right)$  **or**  $x \in \left[ -\frac{1}{2}, 4 \right]$

**Note:** You may see  $x < -\frac{1}{2}, x < 4$  in their initial work before  $-\frac{1}{2} < x < 4$ . Condone this so long as it is clear that the  $-\frac{1}{2} < x < 4$  is their final answer.

Question	Scheme	Marks	AOs
3	$y = \frac{x-4}{2+\sqrt{x}} \Rightarrow \frac{dy}{dx} = \frac{2+\sqrt{x} - (x-4)\frac{1}{2}x^{-\frac{1}{2}}}{(2+\sqrt{x})^2}$	M1 A1	2.1 1.1b
	$= \frac{2+\sqrt{x} - (x-4)\frac{1}{2}x^{-\frac{1}{2}}}{(2+\sqrt{x})^2} = \frac{2+\sqrt{x} - \frac{1}{2}\sqrt{x} + 2x^{-\frac{1}{2}}}{(2+\sqrt{x})^2} = \frac{2\sqrt{x} + \frac{1}{2}x + 2}{\sqrt{x}(2+\sqrt{x})^2}$	M1	1.1b
	$= \frac{x+4\sqrt{x}+4}{2\sqrt{x}(2+\sqrt{x})^2} = \frac{(2+\sqrt{x})^2}{2\sqrt{x}(2+\sqrt{x})^2} = \frac{1}{2\sqrt{x}}$	A1	2.1
		(4)	
<b>(4 marks)</b>			
<b>Notes</b>			

M1: Attempts to use a correct rule e.g. quotient or product (& chain) rule to achieve the following forms

Quotient :  $\frac{\alpha(2+\sqrt{x}) - \beta(x-4)x^{-\frac{1}{2}}}{(2+\sqrt{x})^2}$  but be tolerant of attempts where the  $(2+\sqrt{x})^2$  has been

incorrectly expanded

Product:  $\alpha(2+\sqrt{x})^{-1} + \beta x^{-\frac{1}{2}}(x-4)(2+\sqrt{x})^{-2}$

Alternatively with  $t = \sqrt{x}$ ,  $y = \frac{t^2-4}{2+t} \Rightarrow \frac{dy}{dx} = \frac{dy}{dt} \times \frac{dt}{dx} = \frac{2t(2+t) - (t^2-4)}{(2+t)^2} \times \frac{1}{2}x^{-\frac{1}{2}}$  with same rules

A1: Correct derivative in any form. Must be in terms of a single variable (which could be  $t$ )

M1: Following a correct attempt at differentiation, it is scored for multiplying both numerator and denominator by  $\sqrt{x}$  and collecting terms to form a single fraction. It can also be scored from  $\frac{uv' - vu'}{v^2}$

For the  $t = \sqrt{x}$ , look for an attempt to simplify  $\frac{t^2 + 4t + 4}{(2+t)^2} \times \frac{1}{2t}$

A1: Correct expression showing all key steps with no errors or omissions.  $\frac{dy}{dx}$  must be seen at least once

Question	Scheme	Marks	AOs
3	$y = \frac{x-4}{2+\sqrt{x}} \Rightarrow y = \frac{(\sqrt{x}+2)(\sqrt{x}-2)}{2+\sqrt{x}} = \sqrt{x}-2$	M1 A1	2.1 1.1b
	$\frac{dy}{dx} = \frac{1}{2\sqrt{x}}$	M1 A1	1.1b 2.1
		(4)	
	<b>(4 marks)</b>		
<b>Notes</b>			

M1: Attempts to use difference of two squares. Can also be scored using

$$t = \sqrt{x} \Rightarrow y = \frac{t^2-4}{t+2} \Rightarrow y = \frac{(t+2)(t-2)}{t+2}$$

A1:  $y = \sqrt{x} - 2$  or  $y = t - 2$

M1: Attempts to differentiate an expression of the form  $y = \sqrt{x} + b$

A1: Correct expression showing all key steps with no errors or omissions.  $\frac{dy}{dx}$  must be seen at least once

Question	Scheme	Marks	AOs
4(a)(i)	$\frac{dy}{dx} = 20x^3 - 72x^2 + 84x - 32$	M1 A1	1.1b 1.1b
(ii)	$\frac{d^2y}{dx^2} = 60x^2 - 144x + 84$	A1ft	1.1b
		(3)	
(b)(i)	$x = 1 \Rightarrow \frac{dy}{dx} = 20 - 72 + 84 - 32$	M1	1.1b
	$\frac{dy}{dx} = 0$ so there is a stationary point at $x = 1$	A1	2.1
<b>Alternative for (b)(i)</b>			
	$20x^3 - 72x^2 + 84x - 32 = 4(x-1)^2(5x-8) = 0 \Rightarrow x = \dots$	M1	1.1b
	When $x = 1$ , $\frac{dy}{dx} = 0$ so there is a stationary point	A1	2.1
(b)(ii)	Note that in (b)(ii) there are no marks for <u>just</u> evaluating $\left(\frac{d^2y}{dx^2}\right)_{x=1}$		
	E.g. $\left(\frac{d^2y}{dx^2}\right)_{x=0.8} = \dots \left(\frac{d^2y}{dx^2}\right)_{x=1.2} = \dots$	M1	2.1
	$\left(\frac{d^2y}{dx^2}\right)_{x=0.8} > 0, \left(\frac{d^2y}{dx^2}\right)_{x=1.2} < 0$ Hence point of inflection	A1	2.2a
		(4)	
<b>Alternative 1 for (b)(ii)</b>			
	$\left(\frac{d^2y}{dx^2}\right)_{x=1} = 60x^2 - 144x + 84 = 0$ (is inconclusive) $\left(\frac{d^3y}{dx^3}\right) = 120x - 144 \Rightarrow \left(\frac{d^3y}{dx^3}\right)_{x=1} = \dots$	M1	2.1
	$\left(\frac{d^2y}{dx^2}\right)_{x=1} = 0$ and $\left(\frac{d^3y}{dx^3}\right)_{x=1} \neq 0$ Hence point of inflection	A1	2.2a
<b>Alternative 2 for (b)(ii)</b>			
	E.g. $\left(\frac{dy}{dx}\right)_{x=0.8} = \dots \left(\frac{dy}{dx}\right)_{x=1.2} = \dots$	M1	2.1
	$\left(\frac{dy}{dx}\right)_{x=0.8} < 0, \left(\frac{dy}{dx}\right)_{x=1.2} < 0$ Hence point of inflection	A1	2.2a
<b>(7 marks)</b>			
<b>Notes</b>			
(a)(i) M1: $x^n \rightarrow x^{n-1}$ for at least one power of $x$ A1: $\frac{dy}{dx} = 20x^3 - 72x^2 + 84x - 32$			
(a)(ii)			

A1ft: Achieves a correct  $\frac{d^2y}{dx^2}$  for their  $\frac{dy}{dx} = 20x^3 - 72x^2 + 84x - 32$

(b)(i)

M1: Substitutes  $x = 1$  into their  $\frac{dy}{dx}$

A1: Obtains  $\frac{dy}{dx} = 0$  following a correct derivative and makes a conclusion which can be minimal

e.g. tick, QED etc. which may be in a preamble e.g. stationary point when  $\frac{dy}{dx} = 0$  and then

shows  $\frac{dy}{dx} = 0$

**Alternative:**

M1: Attempts to solve  $\frac{dy}{dx} = 0$  by factorisation. This may be by using the factor of  $(x - 1)$  or possibly using a calculator to find the roots and showing the factorisation. Note that they may divide by 4 before factorising which is acceptable. Need to either see either  $4(x - 1)^2(5x - 8)$  or  $(x - 1)^2(5x - 8)$  for the factorisation or  $x = \frac{8}{5}$  and  $x = 1$  seen as the roots.

A1: Obtains  $x = 1$  and makes a conclusion as above

(b)(ii)

M1: Considers the value of the second derivative either side of  $x = 1$ . Do not be too concerned with the interval for the method mark.

(NB  $\frac{d^2y}{dx^2} = (x - 1)(60x - 84)$  so may use this factorised form when considering  $x < 1$ ,  $x > 1$  for sign change of second derivative)

A1: Fully correct work including a correct  $\frac{d^2y}{dx^2}$  with a reasoned conclusion indicating that the stationary point is a point of inflection. Sufficient reason is e.g. "sign change"/ "> 0, < 0". If values are given they should be correct (but be generous with accuracy) but also just allow "> 0" and "< 0" provided they are correctly paired. The interval must be where  $x < 1.4$

**Alternative 1 for (b)(ii)**

M1: Shows that second derivative at  $x = 1$  is zero and **then finds the third derivative at  $x = 1$**

A1: Fully correct work including a correct  $\frac{d^2y}{dx^2}$  with a reasoned conclusion indicating that stationary point is a point of inflection. Sufficient reason is " $\neq 0$ " but must follow a correct third derivative and a correct value if evaluated. For reference  $\left(\frac{d^3y}{dx^3}\right)_{x=1} = -24$

**Alternative 2 for (b)(ii)**

M1: Considers the value of the first derivative either side of  $x = 1$ . Do not be too concerned with the interval for the method mark.

A1: Fully correct work with a reasoned conclusion indicating that stationary point is a point of inflection. Sufficient reason is e.g. "same sign"/"both negative"/"< 0, < 0". If values are given they should be correct (but be generous with accuracy). The interval must be where  $x < 1.4$

$x$	0	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1
$f'(x)$	-32	-24.3	-17.92	-12.74	-8.64	-5.5	-3.2	-1.62	-0.64	-0.14	0
$f''(x)$	84	70.2	57.6	46.2	36	27	19.2	12.6	7.2	3	0

$x$	1.1	1.2	1.3	1.4	1.5	1.6	1.7
$f'(x)$	-0.1	-0.32	-0.54	-0.64	-0.5	0	0.98
$f''(x)$	-1.8	-2.4	-1.8	0	3	7.2	12.6

Question	Scheme	Marks	AOs
5 (a)	$2 < x < 6$	B1	1.1b
		(1)	
(b)	States either $k > 8$ or $k < 0$	M1	3.1a
	States e.g. $\{k : k > 8\} \cup \{k : k < 0\}$	A1	2.5
		(2)	
(c)	<b>Please see notes for alternatives</b>		
	States $y = ax(x-6)^2$ or $f(x) = ax(x-6)^2$	M1	1.1b
	Substitutes (2,8) into $y = ax(x-6)^2$ and attempts to find $a$	dM1	3.1a
	$y = \frac{1}{4}x(x-6)^2$ or $f(x) = \frac{1}{4}x(x-6)^2$ o.e	A1	2.1
		(3)	
<b>(6 marks)</b>			
<b>Notes: Watch for answers written by the question. If they are beside the question and in the answer space, the one in the answer space takes precedence</b>			

(a)

B1: Deduces  $2 < x < 6$  o.e. such as  $x > 2, x < 6$   $x > 2$  and  $x < 6$   $\{x : x > 2\} \cap \{x : x < 6\}$   $x \in (2, 6)$

Condone attempts in which set notation is incorrectly attempted but correct values can be seen or implied E.g.  $\{x > 2\} \cap \{x < 6\}$   $\{x > 2, x < 6\}$ . Allow just the open interval  $(2, 6)$

Do not allow for incorrect inequalities such as e.g.  $x > 2$  or  $x < 6$ ,  $\{x : x > 2\} \cup \{x : x < 6\}$   $x \in [2, 6]$

(b)

M1: Establishes a correct method by finding one of the (correct) inequalities

States either  $k > 8$  (condone  $k \geq 8$ ) or  $k < 0$  (condone  $k \leq 0$ )

Condone for this mark  $y \leftrightarrow k$  or  $f(x) \leftrightarrow k$  and  $8 < k < 0$

A1: Fully correct solution in the form  $\{k : k > 8\} \cup \{k : k < 0\}$  or  $\{k | k > 8\} \cup \{k | k < 0\}$  either way around

but condone  $\{k < 0\} \cup \{k > 8\}$ ,  $\{k : k < 0 \cup k > 8\}$ ,  $\{k < 0 \cup k > 8\}$ . It is not necessary to mention  $\mathbb{R}$ , e.g.  $\{k : k \in \mathbb{R}, k > 8\} \cup \{k : k \in \mathbb{R}, k < 0\}$  Look for  $\{ \}$  and  $\cup$

Do not allow solutions not in set notation such as  $k < 0$  or  $k > 8$ .

(c)

M1: Realises that the equation of  $C$  is of the form  $y = ax(x-6)^2$ . Condone with  $a = 1$  for this mark.

So award for sight of  $ax(x-6)^2$  even with  $a = 1$

dM1: Substitutes (2,8) into the form  $y = ax(x-6)^2$  and attempts to find the value for  $a$ .

It is dependent upon having an equation, which the ( $y = \dots$ ) may be implied, of the correct form.

A1: Uses all of the information to form a correct **equation** for  $C$   $y = \frac{1}{4}x(x-6)^2$  o.e.

ISW after a correct answer. Condone  $f(x) = \frac{1}{4}x(x-6)^2$  but not  $C = \frac{1}{4}x(x-6)^2$ .

Allow this to be written down for all 3 marks

## Examples of alternative methods

.....  
**Alternative I part (c):****Using the form  $y = ax^3 + bx^2 + cx$  and setting up then solving simultaneous equations.****There are various versions of this but can be marked similarly**M1: Realises that the equation of  $C$  is of the form  $y = ax^3 + bx^2 + cx$  and forms two equations in  $a, b$  and  $c$ . Condone with  $a = 1$  for this mark.Note that the form  $y = ax^3 + bx^2 + cx + d$  is M0 until  $d$  is set equal to 0.

There are four equations that could be formed, only two are necessary for this mark.

Condone slips

Using  $(6, 0) \Rightarrow 216a + 36b + 6c = 0$

Using  $(2, 8) \Rightarrow 8a + 4b + 2c = 8$

Using  $\frac{dy}{dx} = 0$  at  $x = 2 \Rightarrow 12a + 4b + c = 0$

Using  $\frac{dy}{dx} = 0$  at  $x = 6 \Rightarrow 108a + 12b + c = 0$

dM1: Forms and solves three different equations, one of which must be using  $(2, 8)$  to find values for  $a, b$  and  $c$ . A calculator can be used to solve the equationsA1: Uses all of the information to form a correct equation for  $C$   $y = \frac{1}{4}x^3 - 3x^2 + 9x$  o.e.

ISW after a correct answer. Condone  $f(x) = \frac{1}{4}x^3 - 3x^2 + 9x$

.....  
**Alternative II part (c)****Using the gradient and integrating**M1: Realises that the gradient of  $C$  is zero at 2 and 6 so sets  $f'(x) = k(x-2)(x-6)$  or **and** attempts to integrate. Condone with  $k = 1$ dM1: Substitutes  $x = 2, y = 8$  into  $f(x) = k(\dots x^3 + \dots x + \dots)$  and finds a value for  $k$ A1: Uses all of the information to form a correct equation for  $C$   $y = \frac{3}{4}\left(\frac{1}{3}x^3 - 4x^2 + 12x\right)$  o.e.


ISW after a correct answer. Condone  $f(x) = \frac{3}{4}\left(\frac{1}{3}x^3 - 4x^2 + 12x\right)$

.....



Question	Scheme	Marks	AOs
<b>6 (a)</b>	Sets up an allowable equation using volume = 240 E.g. $\frac{1}{2}r^2 \times 0.8h = 240 \Rightarrow h = \frac{600}{r^2}$ o.e.	M1 A1	3.4 1.1b
	Attempts to substitute their $h = \frac{600}{r^2}$ into $(S =) \frac{1}{2}r^2 \times 0.8 + \frac{1}{2}r^2 \times 0.8 + 2rh + 0.8rh$	dM1	3.4
	$S = 0.8r^2 + 2.8rh = 0.8r^2 + 2.8 \times \frac{600}{r} = 0.8r^2 + \frac{1680}{r} *$	A1*	2.1
		(4)	
<b>(b)</b>	$\left(\frac{dS}{dr}\right) = 1.6r - \frac{1680}{r^2}$	M1 A1	3.1a 1.1b
	Sets $\frac{dS}{dr} = 0 \Rightarrow r^3 = 1050$ $r = \text{awrt } 10.2$	dM1 A1	2.1 1.1b
		(4)	
<b>(c)</b>	Attempts to substitute their positive $r$ into $\left(\frac{d^2S}{dr^2}\right) = 1.6 + \frac{3360}{r^3}$ and considers its value or sign	M1	1.1b
	E.g. Correct $\frac{d^2S}{dr^2} = 1.6 + \frac{3360}{r^3}$ with $\frac{d^2S}{dr^2} \Big _{r=10.2} = 5 > 0$ proving a minimum value of $S$	A1	1.1b
		(2)	
<b>(10 marks)</b>			
<b>Notes:</b>			

$$\text{Volume} = 0.4r^2h$$



$$\text{Total surface area} = 2rh + 0.8r^2 + 0.8rh$$

**(a)**

**M1:** Attempts to use the fact that the volume of the toy is  $240 \text{ cm}^3$

Sight of  $\frac{1}{2}r^2 \times 0.8 \times h = 240$  leading to  $h = \dots$  or  $rh = \dots$  scores this mark

But condone an equation of the correct form so allow for  $kr^2h = 240 \Rightarrow h = \dots$  or  $rh = \dots$

**A1:** A correct expression for  $h = \frac{600}{r^2}$  or  $rh = \frac{600}{r}$  which may be left unsimplified.

This may be implied when you see an expression for  $S$  or part of  $S$  E.g.  $2rh = 2r \times \frac{600}{r^2}$

**dM1:** Attempts to substitute their  $h = \frac{a}{r^2}$  o.e. such as  $hr = \frac{a}{r}$  into a **correct** expression for  $S$

Sight of  $\frac{1}{2}r^2 \times 0.8 + \frac{1}{2}r^2 \times 0.8 + rh + rh + 0.8rh$  with an appropriate substitution

Simplified versions such as  $0.8r^2 + 2rh + 0.8rh$  used with an appropriate substitution is fine.

**A1\*:** Correct work leading to the given result.

$S =$ ,  $SA =$  or surface area = must be seen at least once in the correct place

The method must be made clear so expect to see evidence. For example

$S = 0.8r^2 + 2rh + 0.8rh \Rightarrow S = 0.8r^2 + 2r \times \frac{600}{r^2} + 0.8r \times \frac{600}{r^2} \Rightarrow S = 0.8r^2 + \frac{1680}{r}$  would be fine.

**(b)** There is no requirement to see  $\frac{dS}{dr}$  in part (b). It may even be called  $\frac{dy}{dx}$ .

**M1:** Achieves a derivative of the form  $pr \pm \frac{q}{r^2}$  where  $p$  and  $q$  are non- zero constants

**A1:** Achieves  $\left(\frac{dS}{dr}\right) = 1.6r - \frac{1680}{r^2}$

**dM1:** Sets or implies that their  $\frac{dS}{dr} = 0$  and proceeds to  $mr^3 = n$ ,  $m \times n > 0$ . It is dependent upon a

correct attempt at differentiation. This mark may be implied by a correct answer to their  $pr - \frac{q}{r^2} = 0$

**A1:**  $r = \text{awrt } 10.2$  or  $\sqrt[3]{1050}$

**(c)**

**M1:** Attempts to substitute their positive  $r$  (found in (b)) into  $\left(\frac{d^2S}{dr^2}\right) = e \pm \frac{f}{r^3}$  where  $e$  and  $f$  are non zero

and finds its value or sign.

Alternatively considers the sign of  $\left(\frac{d^2S}{dr^2}\right) = e \pm \frac{f}{r^3}$  (at their positive  $r$  found in (b))

Condone the  $\frac{d^2S}{dr^2}$  to be  $\frac{d^2y}{dx^2}$  or being absent, but only for this mark.

**A1:** States that  $\frac{d^2S}{dr^2}$  or  $S'' = 1.6 + \frac{3360}{r^3} = \text{awrt } 5 > 0$  proving a minimum value of  $S$

This is dependent upon having achieved  $r = \text{awrt } 10$  and a correct  $\frac{d^2S}{dr^2} = 1.6 + \frac{3360}{r^3}$

It can be argued without finding the value of  $\frac{d^2S}{dr^2}$ . E.g.  $\frac{d^2S}{dr^2} = 1.6 + \frac{3360}{r^3} > 0$  as  $r > 0$ , so

minimum value of  $S$ . For consistency it is also dependent upon having achieved  $r = \text{awrt } 10$

Do **NOT** allow  $\frac{d^2y}{dx^2}$  for this mark

Question	Scheme	Marks	AOs
7(a)	$\{f'(x) = \dots x^2 + \dots x + \dots \Rightarrow \{f''(x) = \dots x + \dots$	<b>M1</b>	1.1b
	$\{f'(x) = \} 3x^2 + 4x - 8 \Rightarrow \{f''(x) = \} 6x + 4$	<b>A1cso</b>	1.1b
		<b>(2)</b>	
(b)(i)	$"6x + 4" = 0 \Rightarrow x = "-\frac{2}{3}"$	<b>B1ft</b>	1.1b
(ii)	$x ,, "-\frac{2}{3}" \text{ or } x < "-\frac{2}{3}"$	<b>B1ft</b>	2.2a
		<b>(2)</b>	

**(4 marks)****Notes****(a)****M1:** For attempting to differentiate twice.

It can be scored for any of:  $x^3 \rightarrow \dots x^2 \rightarrow \dots x$  or  $2x^2 \rightarrow \dots x \rightarrow k$  or  $-8x \rightarrow k \rightarrow 0$  where ... are constants.

You can ignore the lhs so do not be concerned what they call the first and/or second derivative, just look for their expressions.

The indices do not need to be processed for this mark so allow for e.g.  $x^3 \rightarrow \dots x^{3-1} \rightarrow \dots x^{3-1-1}$

**A1cso:** ( $f''(x) =$ )  $6x + 4$  Correct second derivative from fully correct work. The " $f''(x) =$ " is not required.

Allow  $6x^1$  for  $6x$  but not  $4x^0$  for 4 unless the  $4x^0$  becomes 4 later, e.g. in part (b).

Do **not** apply isw so mark their final answer. E.g. if  $6x + 4$  becomes  $3x + 2$  score A0

**(b)****(i)**

**B1ft:**  $ax + b = 0 \Rightarrow (x =) -\frac{b}{a}$ . This mark is for obtaining  $x = -\frac{2}{3}$  **or**  $x = -\frac{b}{a}$  which has come from solving an equation of the form  $ax + b$ ,  $a, b \neq 0$  where  $ax + b$  is their attempt to differentiate twice in part (a)

Allow equivalent fractions e.g.  $x = -\frac{4}{6}$  or equivalents for their  $x = -\frac{b}{a}$  or an exact decimal and isw.

**(ii)**

**B1ft:** Deduces  $x ,, -\frac{2}{3}$  **or** follow through their single value of  $x$  from part (i) obtained from their attempt to solve an equation of the form  $ax + b = 0$ ,  $a, b \neq 0$  where  $ax + b$  was their attempt to differentiate twice in part (a). Do not isw and mark their final answer.

If 2 inequalities are given e.g.  $x < "-\frac{2}{3}"$ ,  $x > "-\frac{2}{3}"$  without indicating which is their answer score B0

Condone  $<$  for  $,,$  and allow equivalent inequalities e.g.  $-\frac{2}{3} > x$

Allow equivalent fractions e.g.  $x = -\frac{4}{6}$  or equivalents for their  $x = -\frac{b}{a}$

Allow equivalent notation so these are all acceptable:

$$x ,, "-\frac{2}{3}", x < "-\frac{2}{3}", \left(-\infty, "-\frac{2}{3}"\right], \left(-\infty, "-\frac{2}{3}"\right), \left\{x : x ,, "-\frac{2}{3}"\right\}, \left\{x : x < "-\frac{2}{3}"\right\}$$

Ignore any reference to values of  $y$ .

Allow ft decimal answers from (i) which may be inexact.

Correct answers in part (b) with no working in (a) can score 0011.