

# DIFFERENTIATION

1 Differentiate with respect to  $x$

- |                      |                       |                                    |   |
|----------------------|-----------------------|------------------------------------|---|
| <b>a</b> $\cos x$    | <b>b</b> $5 \sin x$   | <b>c</b> $\cos 3x$                 | <b>d</b> $\sin \frac{1}{4}x$                  |
| <b>e</b> $\sin(x+1)$ | <b>f</b> $\cos(3x-2)$ | <b>g</b> $4 \sin(\frac{\pi}{3}-x)$ | <b>h</b> $\cos(\frac{1}{2}x + \frac{\pi}{6})$ |
| <b>i</b> $\sin^2 x$  | <b>j</b> $2 \cos^3 x$ | <b>k</b> $\cos^2(x-1)$             | <b>l</b> $\sin^4 2x$                          |

2 Use the derivatives of  $\sin x$  and  $\cos x$  to show that

- |  |   |
|--|---|
| <b>a</b> $\frac{d}{dx}(\tan x) = \sec^2 x$                                       | <b>b</b> $\frac{d}{dx}(\sec x) = \sec x \tan x$             |
| <b>c</b> $\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$ | <b>d</b> $\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$ |

3 Differentiate with respect to  $t$

- |                       |  |                                     |   |
|-----------------------|--|-------------------------------------|---|
| <b>a</b> $\cot 2t$    | <b>b</b> $\sec(t+2)$                                 | <b>c</b> $\tan(4t-3)$               | <b>d</b> $\operatorname{cosec} 3t$      |
| <b>e</b> $\tan^2 t$   | <b>f</b> $3 \operatorname{cosec}(t + \frac{\pi}{6})$ | <b>g</b> $\cot^3 t$                 | <b>h</b> $4 \sec \frac{1}{2}t$          |
| <b>i</b> $\cot(2t-3)$ | <b>j</b> $\sec^2 2t$                                 | <b>k</b> $\frac{1}{2} \tan(\pi-4t)$ | <b>l</b> $\operatorname{cosec}^2(3t+1)$ |

4 Differentiate with respect to  $x$

- |                        |                          |  |                         |
|------------------------|--------------------------|--|-------------------------|
| <b>a</b> $\ln(\sin x)$ | <b>b</b> $6e^{\tan x}$   | <b>c</b> $\sqrt{\cos 2x}$                | <b>d</b> $e^{\sin 3x}$  |
| <b>e</b> $2 \cot x^2$  | <b>f</b> $\sqrt{\sec x}$ | <b>g</b> $3e^{-\operatorname{cosec} 2x}$ | <b>h</b> $\ln(\tan 4x)$ |

5 Find the coordinates of any stationary points on each curve in the interval  $0 \leq x \leq 2\pi$ .

- |                             |                                  |                                 |
|-----------------------------|----------------------------------|---------------------------------|
| <b>a</b> $y = x + 2 \sin x$ | <b>b</b> $y = 2 \sec x - \tan x$ | <b>c</b> $y = \sin x + \cos 2x$ |
|-----------------------------|----------------------------------|---------------------------------|

6 Find an equation for the tangent to each curve at the point on the curve with the given  $x$ -coordinate.

- |                             |                     |   |                     |
|-----------------------------|---------------------|---|---------------------|
| <b>a</b> $y = 1 + \sin 2x,$ | $x = 0$             | <b>b</b> $y = \cos x,$                            | $x = \frac{\pi}{3}$ |
| <b>c</b> $y = \tan 3x,$     | $x = \frac{\pi}{4}$ | <b>d</b> $y = \operatorname{cosec} x - 2 \sin x,$ | $x = \frac{\pi}{6}$ |

7 Differentiate with respect to  $x$

- |                                       |                                |                             |                                   |
|---------------------------------------|--------------------------------|-----------------------------|-----------------------------------|
| <b>a</b> $x \sin x$                   | <b>b</b> $\frac{\cos 2x}{x}$   | <b>c</b> $e^x \cos x$       | <b>d</b> $\sin x \cos x$          |
| <b>e</b> $x^2 \operatorname{cosec} x$ | <b>f</b> $\sec x \tan x$       | <b>g</b> $\frac{x}{\tan x}$ | <b>h</b> $\frac{\sin 2x}{e^{3x}}$ |
| <b>i</b> $\cos^2 x \cot x$            | <b>j</b> $\frac{\sec 2x}{x^2}$ | <b>k</b> $x \tan^2 4x$      | <b>l</b> $\frac{\sin x}{\cos 2x}$ |

8 Find the value of  $f'(x)$  at the value of  $x$  indicated in each case.

- |   |                     |                                      |                     |
|---|---------------------|--------------------------------------|---------------------|
| <b>a</b> $f(x) = \sin 3x \cos 5x,$              | $x = \frac{\pi}{4}$ | <b>b</b> $f(x) = \tan 2x \sin x,$    | $x = \frac{\pi}{3}$ |
| <b>c</b> $f(x) = \frac{\ln(2 \cos x)}{\sin x},$ | $x = \frac{\pi}{3}$ | <b>d</b> $f(x) = \sin^2 x \cos^3 x,$ | $x = \frac{\pi}{6}$ |

## DIFFERENTIATION

continued

- 9 Find an equation for the normal to the curve  $y = 3 + x \cos 2x$  at the point where it crosses the  $y$ -axis.

- 10 A curve has the equation  $y = \frac{2 + \sin x}{1 - \sin x}$ ,  $0 \leq x \leq 2\pi$ ,  $x \neq \frac{\pi}{2}$ .

- a Find and simplify an expression for  $\frac{dy}{dx}$ .
- b Find the coordinates of the turning point of the curve.
- c Show that the tangent to the curve at the point  $P$ , with  $x$ -coordinate  $\frac{\pi}{6}$ , has equation

$$y = 6\sqrt{3}x + 5 - \sqrt{3}\pi.$$

- 11 A curve has the equation  $y = e^{-x} \sin x$ .

- a Find  $\frac{dy}{dx}$  and  $\frac{d^2y}{dx^2}$ .
- b Find the exact coordinates of the stationary points of the curve in the interval  $-\pi \leq x \leq \pi$  and determine their nature.


- 12 The curve  $C$  has the equation  $y = x \sec x$ .

- a Show that the  $x$ -coordinate of any stationary point of  $C$  must satisfy the equation

$$1 + x \tan x = 0.$$

- b By sketching two suitable graphs on the same set of axes, deduce the number of stationary points  $C$  has in the interval  $0 \leq x \leq 2\pi$ .

13



The diagram shows the curve  $y = f(x)$  in the interval  $0 \leq x \leq 2\pi$ , where

$$f(x) \equiv \cos x \sin 2x.$$

- a Show that  $f'(x) = 2 \cos x (1 - 3 \sin^2 x)$ .
- b Find the  $x$ -coordinates of the stationary points of the curve in the interval  $0 \leq x \leq 2\pi$ .
- c Show that the maximum value of  $f(x)$  in the interval  $0 \leq x \leq 2\pi$  is  $\frac{4}{9}\sqrt{3}$ .
- d Explain why this is the maximum value of  $f(x)$  for all real values of  $x$ .
- 14 A curve has the equation  $y = \operatorname{cosec} \left(x - \frac{\pi}{6}\right)$  and crosses the  $y$ -axis at the point  $P$ .
- a Find an equation for the normal to the curve at  $P$ .
- The point  $Q$  on the curve has  $x$ -coordinate  $\frac{\pi}{3}$ .
- b Find an equation for the tangent to the curve at  $Q$ .
- The normal to the curve at  $P$  and the tangent to the curve at  $Q$  intersect at the point  $R$ .
- c Show that the  $x$ -coordinate of  $R$  is given by  $\frac{8\sqrt{3} + 4\pi}{13}$ .