

Question	Scheme	Marks	AOs
1(a)	$f'(x) = 2x + \frac{4x-4}{2x^2-4x+5}$	M1 A1	1.1b 1.1b
	$2x + \frac{4x-4}{2x^2-4x+5} = 0 \Rightarrow 2x(2x^2-4x+5) + 4x-4 = 0$	dM1	1.1b
	$2x^3 - 4x^2 + 7x - 2 = 0^*$	A1*	2.1
		(4)	
(b)	(i) $x_2 = \frac{1}{7}(2 + 4(0.3)^2 - 2(0.3)^3)$	M1	1.1b
	$x_2 = 0.3294$	A1	1.1b
	(ii) $x_4 = 0.3398$	A1	1.1b
		(3)	
(c)	$h(x) = 2x^3 - 4x^2 + 7x - 2$ $h(0.3415) = 0.00366... \quad h(0.3405) = -0.00130...$	M1	3.1a
	States: <ul style="list-style-type: none"> • there is a change of sign • $f'(x)$ is continuous • $\alpha = 0.341$ to 3dp 	A1	2.4
		(2)	
(9 marks)			
Notes			

(a)

M1: Differentiates $\ln(2x^2 - 4x + 5)$ to obtain $\frac{g(x)}{2x^2 - 4x + 5}$ where $g(x)$ could be 1

A1: For $f'(x) = 2x + \frac{4x - 4}{2x^2 - 4x + 5}$

dM1: Sets their $f'(x) = ax + \frac{g(x)}{2x^2 - 4x + 5} = 0$ and uses "correct" algebra, condoning slips, to obtain a cubic equation. E.g Look for $ax(2x^2 - 4x + 5) \pm g(x) = 0$ o.e. , condoning slips, followed by some attempt to simplify

A1*: Achieves $2x^3 - 4x^2 + 7x - 2 = 0$ with no errors. (The dM1 mark must have been awarded)

(b)(i)

M1: Attempts to use the iterative formula with $x_1 = 0.3$. If no method is shown award for $x_2 = \text{awrt } 0.33$

A1: $x_2 = \text{awrt } 0.3294$ Note that $\frac{1153}{3500}$ is correct

Condone an incorrect suffix if it is clear that a correct value has been found

(b)(ii)

A1: $x_4 = \text{awrt } 0.3398$ Condone an incorrect suffix if it is clear that a correct value has been found

(c)

M1: Attempts to substitute $x = 0.3415$ and $x = 0.3405$ into a suitable function and gets one value correct (rounded or truncated to 1 sf). It is allowable to use a tighter interval that contains the root 0.340762654

Examples of suitable functions are $2x^3 - 4x^2 + 7x - 2$, $x - \frac{1}{7}(4x^2 - 2x^3 + 2)$ and $f'(x)$ as this has been

found in part (a) with $f'(0.3405) = -0.00067\dots$, $f'(0.3415) = (+)0.0018$

There must be sufficient evidence for the function, which would be for example, a statement such as $h(x) = 2x^3 - 4x^2 + 7x - 2$ or sight of embedded values that imply the function, not just a value or values

even if both are correct. Condone $h(x)$ being mislabelled as f

$h(0.3415) = 2 \times 0.3415^3 - 4 \times 0.3415^2 + 7 \times 0.3415 - 2$

A1: Requires

- both calculations correct (rounded or truncated to 1sf)
- a statement that there is a change in sign and that the function is continuous
- a minimal conclusion e.g. \checkmark , proven, $\alpha = 0.341$, root

Question	Scheme	Marks	AOs
2(a)	$\dots xe^x + \dots e^x$	M1	1.1b
	$k(xe^x + e^x)$	A1	1.1b
	$\frac{d}{dx}(\sqrt{e^{3x}-2}) = \frac{1}{2} \times 3e^{3x}(e^{3x}-2)^{-\frac{1}{2}}$	B1	1.1b
	$(f'(x) =) \frac{(e^{3x}-2)^{\frac{1}{2}}(7xe^x + 7e^x) - \frac{3}{2}e^{3x}(e^{3x}-2)^{-\frac{1}{2}} \times 7xe^x}{e^{3x}-2}$	dM1	2.1
	$f'(x) = \frac{7e^x(e^{3x}(2-x) - 4x - 4)}{2(e^{3x}-2)^{\frac{3}{2}}}$	A1	1.1b
		(5)	
(b)	$e^{3x}(2-x) - 4x - 4 = 0 \Rightarrow x(\dots e^{3x} \pm \dots) = \dots e^{3x} \pm \dots$	M1	1.1b
	$\Rightarrow x = \frac{2e^{3x}-4}{e^{3x}+4} *$	A1*	2.1
		(2)	
(c)	Draws a vertical line $x=1$ up to the curve then across to the line $y=x$ then up to the curve finishing at the root (need to see a minimum of 2 vertical and horizontal lines tending to the root)	B1	2.1
		(1)	
(d)(i)	$x_2 = \frac{2e^3-4}{e^3+4} = 1.5017756\dots$	M1	1.1b
	$x_2 = \text{awrt } 1.502$	A1	1.1b
(ii)	$\beta = 1.968$	dB1	2.2b
		(3)	
(e)	$h(x) = \frac{2e^{3x}-4}{e^{3x}+4} - x$ $h(0.4315) = -0.000297\dots$ $h(0.4325) = 0.000947\dots$	M1	3.1a
	Both calculations correct and e.g. states: <ul style="list-style-type: none"> • There is a change of sign • e.g $f'(x)$ is continuous • $\alpha = 0.432$ (to 3dp) 	A1cao	2.4
		(2)	

(13 marks)**Notes****(a)**

M1: Attempts the product rule on xe^x (or may be $7xe^x$) achieving an expression of the form $\dots xe^x \pm \dots e^x$. If it is clear that the quotient rule has been applied instead which may be quoted then M0.

A1: $k(xe^x + e^x)$ (e.g. $7(xe^x + e^x)$) or equivalent which may be unsimplified (may be implied by further work)

B1: $\left(\frac{d}{dx}(\sqrt{e^{3x}-2})\right) = \frac{1}{2} \times 3e^{3x}(e^{3x}-2)^{-\frac{1}{2}}$ (simplified or unsimplified)

dM1: Attempts to use the quotient rule. It is dependent on the previous method mark.

Score for achieving an expression of the form

$$(f'(x) =) \frac{(e^{3x} - 2)^{\frac{1}{2}} ("7" xe^x + "7" e^x) - " \frac{3}{2} " e^{3x} (e^{3x} - 2)^{-\frac{1}{2}} \times "7" xe^x}{e^{3x} - 2} \text{ or equivalent (do not be}$$

concerned by the constants for their "7" or their " $\frac{3}{2}$ " which may be both 1)

If it is clear that the quotient rule has been applied the wrong way round then score M0.

Alternatively, applies the product rule. Score for achieving an expression of the form

$$(f'(x) =) (e^{3x} - 2)^{\frac{1}{2}} ("7" xe^x + "7" e^x) - " \frac{3}{2} " e^{3x} (e^{3x} - 2)^{-\frac{3}{2}} \times "7" xe^x \text{ or equivalent (do not be}$$

concerned by the constants for their "7" or their " $\frac{3}{2}$ " which may be both 1)

Do not condone invisible brackets.

A1: $(f'(x) =) \frac{7e^x(e^{3x}(2-x) - 4x - 4)}{2(e^{3x} - 2)^{\frac{3}{2}}}$ following a fully correct differentiated expression.

You may need to check to see if (a) is continued after other parts for evidence of this.

Condone the lack of $f'(x) =$ on the left hand side or allow the use of $\frac{dy}{dx}$ or y' instead.

Alternative (a) attempt using the triple product rule

$$\text{e.g. } \frac{d}{dx} \left(7xe^x(e^{3x} - 2)^{\frac{1}{2}} \right) = 7e^x(e^{3x} - 2)^{\frac{1}{2}} + 7xe^x(e^{3x} - 2)^{\frac{1}{2}} + 7xe^x \times \left(-\frac{1}{2} \right) \times 3e^{3x}(e^{3x} - 2)^{-\frac{3}{2}}$$

$$\Rightarrow \frac{(7e^x + 7xe^x)(e^{3x} - 2) + 7xe^x \times \left(-\frac{1}{2} \right) \times 3e^{3x}}{(e^{3x} - 2)^{\frac{3}{2}}} = \frac{7e^x \left(e^{3x} - 2 + xe^{3x} - 2x - \frac{3}{2} xe^{3x} \right)}{(e^{3x} - 2)^{\frac{3}{2}}} \Rightarrow \frac{7e^x(e^{3x}(2-x) - 4x - 4)}{2(e^{3x} - 2)^{\frac{3}{2}}}$$

M1: Attempts the product rule on $xe^x \rightarrow \dots xe^x \pm \dots e^x$ which may be seen within the expression

$\dots e^x(e^{3x} - 2)^{\frac{1}{2}} \pm \dots xe^x(e^{3x} - 2)^{\frac{1}{2}} + \dots$ simplified or unsimplified.

A1: $k(xe^x + e^x)$ which may be seen within the expression $k \left(e^x(e^{3x} - 2)^{\frac{1}{2}} + xe^x(e^{3x} - 2)^{\frac{1}{2}} \right) + \dots$

simplified or unsimplified.

B1: $\left(-\frac{1}{2} \right) \times 3e^{3x}(e^{3x} - 2)^{-\frac{3}{2}}$ which may be seen within the expression $\dots + k \left(xe^x \times \left(-\frac{1}{2} \right) \times 3e^{3x}(e^{3x} - 2)^{-\frac{3}{2}} \right)$

simplified or unsimplified.

dM1: A complete method using all three products (which may appear all on one line). Do not condone invisible brackets.

A1: As above in main scheme notes.

(b) Note that if they do not have values $A = -4$, $B = -4$ in (a) (which may be seen later) then maximum score is M1A0*


M1: Sets their $e^{3x}(2-x) - 4x - 4$ equal to zero, collects terms in x on one side of the equation and non x terms on the other and attempts to factorise the side with x as a common factor. Condone sign slips only for this mark. Allow A and B to be used instead of "-4" and "-4"

A1*: Achieves the given answer with no errors including invisible brackets. If they do not reach the printed answer then it is A0. If they subsequently write $x_{n+1} = \frac{2e^{3x_n} - 4}{e^{3x_n} + 4}$ then isw


(c)

B1: Starting at $x_1 = 1$ look for at least 2 sets of vertical and horizontal lines drawn (may be dashes) tending to β . Condone a lack of arrows on the lines but the sequence of lines should finish at the point of intersection where the root is. Condone the initial vertical line not starting from the x -axis. Mark the intention to draw horizontal and vertical lines. If they have any lines to the left of $x = 1$ this is B0. If they use both diagrams and do not indicate which one they want marking, then the “copy of Diagram 1” should be marked.

Examples scoring B1:



Examples scoring B0:



(d)(i)

M1: Substitutes 1 into the iterative formula. The values embedded in the formula is sufficient for this mark. May be implied by awrt 1.50

A1: awrt 1.502 isw

(d)(ii)

dB1: 1.968 cao (which can only be scored if M1 is scored in (d)(i))

SC: If (d)(i) is rounded to 1.50 then allow 1.97 in (d)(ii) to score M1A0dB1 for (d)

(e)

M1: Attempts to substitute $x = 0.4315$ and 0.4325 into a suitable function and gets one value correct (rounded or truncated to 1sf). It is allowable to use a tighter interval that contains the root $0.4317388728\dots$

If no function is stated then may be implied by their answers to e.g. $f'(0.4315)$, $f'(0.4325)$

You will need to check their calculation is correct.

Other possible functions include:

- $h(x) = x - \frac{2e^{3x} - 4}{e^{3x} + 4}$ (other way round to MS) $h(0.4315) = 0.0002974\dots$, $h(0.4325) = -0.0009479\dots$

- their $f'(x) = \pm \left(\frac{7e^x (e^{3x}(2-x) - 4x - 4)}{2(e^{3x} - 2)^{\frac{3}{2}}} \right)$

(If correct A and B then $f'(0.4315) = \mp 0.005789\dots$, $f'(0.4325) = \pm 0.01831\dots$)

- their $g(x) = \pm (e^{3x}(2-x) - 4x - 4)$

(If correct A and B then $g(0.4315) = \mp 0.002275\dots$, $g(0.4325) = \pm 0.007261\dots$)

A1: Requires

- Both calculations correct (rounded or truncated to 1sf)
- A statement that there is a change in sign and that their **function** is continuous (must refer to the function used for the substitution (which is not $f(x)$)

Accept equivalent statements for $f'(0.4315) < 0$, $f'(0.4325) > 0$ e.g.

$f'(0.4315) \times f'(0.4325) < 0$, “one negative one positive”. A minimum is “change of sign and continuous” but do **not** allow this mark if the comment about continuity is clearly incorrect e.g. “because x is continuous” or “because the interval is continuous”

- A minimal conclusion e.g. “hence $\alpha = 0.432$ ”, “so rounds to 0.432”. Do not allow “hence root”