


## Vectors Cheat Sheet

### Vectors

A vector has both **magnitude** and **direction**.

You can represent a vector using a **directed line segment**. Direction of arrow shows the direction of vector



### Triangle law for vector addition:


You can add two vector using the triangle law for vector addition.

The law states that


$$\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}$$

If  $\overrightarrow{AB} = \mathbf{a}$ ,  $\overrightarrow{BC} = \mathbf{b}$  and  $\overrightarrow{AC} = \mathbf{c}$  then,  $\mathbf{a} + \mathbf{b} = \mathbf{c}$

**Example 1:** The diagram shows vectors  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$ . Draw a diagram to illustrate the vector addition




Subtracting a vector is same as adding a negative vector  $\mathbf{a} - \mathbf{b} = \mathbf{a} + (-\mathbf{b})$



### Multiplying a vector:

You can multiply a vector by a scalar. If the number is  $+ve$  the resultant vector has same direction but different length. If the number is  $-ve$ , the new vector has different length but opposite direction




Any vector parallel to the vector  $\mathbf{a}$  may be written as  $\lambda \mathbf{a}$  where  $\lambda$  is a non-zero scalar

### Parallelogram Law for vector addition:

For the parallelogram  $ABCD$ ,

If  $\overrightarrow{AB} = \mathbf{a}$ ,  $\overrightarrow{AD} = \mathbf{b}$  then  $\overrightarrow{AC} = \mathbf{a} + \mathbf{b}$



**Example 2:** Show that the vectors  $6\mathbf{a} + 8\mathbf{b}$  and  $9\mathbf{a} + 12\mathbf{b}$  are parallel.


Try to write one of the vectors as  $\lambda \times$  other vector

$$9\mathbf{a} + 12\mathbf{b} = \frac{3}{2}(6\mathbf{a} + 8\mathbf{b})$$

Since,  $9\mathbf{a} + 12\mathbf{b}$  can be written as  $\lambda \times (6\mathbf{a} + 8\mathbf{b})$  where  $\lambda = \frac{3}{2}$ , this implies that  $6\mathbf{a} + 8\mathbf{b}$  and  $9\mathbf{a} + 12\mathbf{b}$  are parallel.

### Representing vector:

A vector represent change in position or displacement relative to the change in  $x$ - and  $y$ -axes



To multiply a column vector by a scalar multiply each component by the scalar:  $\lambda \begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} \lambda p \\ \lambda q \end{pmatrix}$

To add 2 column vector add the  $x$ - component and the  $y$ -components:  $\begin{pmatrix} p \\ q \end{pmatrix} + \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} p+r \\ q+s \end{pmatrix}$

**Example 3:** Given that  $\mathbf{a} = 2\mathbf{i} + 3\mathbf{j}$  and  $\mathbf{b} = 4\mathbf{i} - \mathbf{j}$ , find the vector  $4\mathbf{b} - \mathbf{a}$  in terms of  $\mathbf{i}$  and  $\mathbf{j}$

Start by substituting vectors  $\mathbf{a}$  and  $\mathbf{b}$  in  $4\mathbf{b} - \mathbf{a}$  and then simplify

$$4\mathbf{b} - \mathbf{a} = 4(4\mathbf{i} - \mathbf{j}) - (2\mathbf{i} + 3\mathbf{j})$$

$$= (16\mathbf{i} - 4\mathbf{j}) - (2\mathbf{i} + 3\mathbf{j}) = (16 - 2)\mathbf{i} + (-4 - 3)\mathbf{j} = 14\mathbf{i} - 7\mathbf{j}$$

Hence,  $4\mathbf{b} - \mathbf{a}$  in terms of  $\mathbf{i}$  and  $\mathbf{j}$  is  $14\mathbf{i} - 7\mathbf{j}$

**Example 4:** Given that  $\mathbf{a} = \begin{pmatrix} 9 \\ 7 \end{pmatrix}$ ,  $\mathbf{b} = \begin{pmatrix} 11 \\ -3 \end{pmatrix}$  and  $\mathbf{c} = \begin{pmatrix} -8 \\ -1 \end{pmatrix}$  find  $2\mathbf{a} + \mathbf{b} + \mathbf{c}$

Start by substituting vectors  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{c}$  in  $2\mathbf{a} + \mathbf{b} + \mathbf{c}$

$$2\mathbf{a} + \mathbf{b} + \mathbf{c} = 2\begin{pmatrix} 9 \\ 7 \end{pmatrix} + \begin{pmatrix} 11 \\ -3 \end{pmatrix} + \begin{pmatrix} -8 \\ -1 \end{pmatrix} = \begin{pmatrix} 18 \\ 14 \end{pmatrix} + \begin{pmatrix} 11 \\ -3 \end{pmatrix} + \begin{pmatrix} -8 \\ -1 \end{pmatrix} = \begin{pmatrix} 21 \\ 10 \end{pmatrix}$$

### Unit Vector:

A unit vector is a vector of length 1.

The unit vectors along the  $x$ - and  $y$ -axes are usually denoted by  $\mathbf{i}$  and  $\mathbf{j}$  respectively.


$$\mathbf{i} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \mathbf{j} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

You can write any two-dimensional vector in the form  $\begin{pmatrix} p \\ q \end{pmatrix} = p\mathbf{i} + q\mathbf{j}$

In this topic you will learn to find a unit vector in the direction of a given vector

A unit vector in the direction of  $\mathbf{a}$  is  $\frac{\mathbf{a}}{|\mathbf{a}|}$

so if  $|\mathbf{a}| = 5$ , then a unit vector in the direction of  $\mathbf{a}$  is  $\frac{\mathbf{a}}{5}$



### Magnitude and direction:

To calculate the magnitude of vector you can use Pythagoras theorem.

For the vector  $\mathbf{a} = x\mathbf{i} + y\mathbf{j} = \begin{pmatrix} x \\ y \end{pmatrix}$  the magnitude of the vector is given by

$$|\mathbf{a}| = \sqrt{x^2 + y^2}$$

The two straight lines on either side of the vector is the notation for magnitude of a vector

**Example 4:**  $\mathbf{a} = 2\mathbf{i} + 3\mathbf{j}$ ,  $\mathbf{b} = 3\mathbf{i} - 4\mathbf{j}$ . Find the exact value of the magnitude  $\mathbf{a} + \mathbf{b}$

Since we need to calculate exact value of  $\mathbf{a} + \mathbf{b}$ , first we will have to add the two vectors.

We can do that by representing the vectors in column vector form and then add

So,  $\mathbf{a} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$  and  $\mathbf{b} = \begin{pmatrix} 3 \\ -4 \end{pmatrix}$  implies  $\mathbf{a} + \mathbf{b} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} + \begin{pmatrix} 3 \\ -4 \end{pmatrix} = \begin{pmatrix} 2+3 \\ 3-4 \end{pmatrix} = \begin{pmatrix} 5 \\ -1 \end{pmatrix}$

Use the formula for magnitude which is  $|\mathbf{a}| = \sqrt{x^2 + y^2}$

$$\text{Hence, } |\mathbf{a} + \mathbf{b}| = \sqrt{5^2 + (-1)^2} = \sqrt{26}$$

Keep your answer in surd form as you are asked to find exact value.

**Example 5:** For vector  $\mathbf{a} = 5\mathbf{i} - 12\mathbf{j}$ , find the unit vector in the same direction.

A unit vector in the direction of  $\mathbf{a}$  is  $\frac{\mathbf{a}}{|\mathbf{a}|}$


$$\text{Let us first calculate } |\mathbf{a}|, \text{ so } |\mathbf{a}| = \sqrt{5^2 + (-12)^2} = \sqrt{25 + 144} = \sqrt{169} = 13$$

To find the unit vector in the direction of  $\mathbf{a}$  first write the vector in column vector form and multiply by  $\frac{1}{|\mathbf{a}|}$

$$\frac{\mathbf{a}}{|\mathbf{a}|} = \frac{\mathbf{a}}{13} = \frac{1}{13} \begin{pmatrix} 5 \\ -12 \end{pmatrix} = \begin{pmatrix} \frac{5}{13} \\ -\frac{12}{13} \end{pmatrix}$$

Hence, the unit vector in the direction of  $\mathbf{a}$  is  $\begin{pmatrix} \frac{5}{13} \\ -\frac{12}{13} \end{pmatrix}$

**Example 6:** Find the angle that the vector  $3\mathbf{i} - 5\mathbf{j}$  makes with  $\mathbf{j}$  axis



The angle the vector  $3\mathbf{i} - 5\mathbf{j}$  makes with  $\mathbf{j}$  is  $90^\circ + \text{angle } \theta$

Vector  $3\mathbf{i} - 5\mathbf{j}$  on axes will be 3 units to the right on  $\mathbf{i}$  axis and 5 units down on  $\mathbf{j}$  axis

We can find angle  $\theta$  by using trigonometric ratio  $\tan \theta = \frac{\text{opposite}}{\text{adjacent}} = \frac{5}{3}$

So,  $\tan \theta = \frac{5}{3}$  taking inverse we get  $\theta = \tan^{-1} \left( \frac{5}{3} \right) = 59^\circ$

So the angle  $3\mathbf{i} - 5\mathbf{j}$  makes with  $\mathbf{j}$  is  $90^\circ + 59^\circ = 149^\circ$

### Position Vector:

Position vector are the vectors that gives us the position of a point relative to fixed origin

In general, a point  $P$  with coordinates  $(p, q)$  has a position vector

$$\overrightarrow{OP} = p\mathbf{i} + q\mathbf{j} = \begin{pmatrix} p \\ q \end{pmatrix}$$

**Example 7:**  $\overrightarrow{OA} = 5\mathbf{i} - 2\mathbf{j}$  and  $\overrightarrow{AB} = 3\mathbf{i} + 4\mathbf{j}$ . Find the position vector of  $B$

Start by drawing the vectors on the axes to find position vector of  $B$ . From the diagram position vector of  $B$  is vector  $\overrightarrow{OB}$ . It is easier to work with column vectors, so write the vectors

in column vector form


$$\overrightarrow{OA} = \begin{pmatrix} 5 \\ -2 \end{pmatrix} \text{ and } \overrightarrow{AB} = \begin{pmatrix} 3 \\ 4 \end{pmatrix}$$

Using triangle law of vector addition vector

$$\overrightarrow{OB} = \overrightarrow{OA} + \overrightarrow{AB}$$

$$\overrightarrow{OB} = \begin{pmatrix} 5 \\ -2 \end{pmatrix} + \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 5+3 \\ -2+4 \end{pmatrix} = \begin{pmatrix} 8 \\ 2 \end{pmatrix}$$

$$\text{In } \mathbf{i}, \mathbf{j} \text{ form } \overrightarrow{OB} = 8\mathbf{i} + 2\mathbf{j}$$




### Solving geometric problem:

You can use vector to find the position vector of a point that divides a line segment in a given ratio

If the point  $P$  divides the line segment  $AB$  in the ratio  $\lambda : \mu$ , then

$$\begin{aligned} \overrightarrow{OP} &= \overrightarrow{OA} + \frac{\lambda}{\lambda + \mu} \overrightarrow{AB} \\ &= \overrightarrow{OA} + \frac{\lambda}{\lambda + \mu} \overrightarrow{OB} - \overrightarrow{OA} \end{aligned}$$



**Example 8:** In the diagram the points  $A$  and  $B$  have position vectors  $\mathbf{a}$  and  $\mathbf{b}$  respectively.

The point  $P$  divides  $AB$  in the ratio  $1:2$ . Find the position vector of  $P$

The position vector of  $P$  is vector  $\overrightarrow{OP}$ . You can refer to the figure above. To find  $\overrightarrow{OP}$  we can make use of the equation given above for point  $P$  dividing the line segment  $AB$  in the ratio  $\lambda : \mu$ . Here  $\lambda = 1$  and  $\mu = 2$  and substituting them in to the equation gives us,

$$\overrightarrow{OP} = \overrightarrow{OA} + \frac{1}{3} \overrightarrow{AB}$$

$$\overrightarrow{OP} = \overrightarrow{OA} + \frac{1}{3} (\overrightarrow{OB} - \overrightarrow{OA})$$

$$\overrightarrow{OP} = \overrightarrow{OA} + \frac{1}{3} \overrightarrow{OB} - \frac{1}{3} \overrightarrow{OA}$$

$$= \overrightarrow{OA} - \frac{1}{3} \overrightarrow{OA} + \frac{1}{3} \overrightarrow{OB} = \frac{2}{3} \overrightarrow{OA} + \frac{1}{3} \overrightarrow{OB} = \frac{2}{3} \mathbf{a} + \frac{1}{3} \mathbf{b}$$

Hence,  $\overrightarrow{OP}$  is  $\frac{2}{3} \mathbf{a} + \frac{1}{3} \mathbf{b}$

You will have to write  $\overrightarrow{AB}$  in terms of the position vectors of  $A$  and  $B$  in order to find  $\overrightarrow{OP}$

If  $\mathbf{a}$  and  $\mathbf{b}$  are two non-parallel vectors and  $p\mathbf{a} + q\mathbf{b} = r\mathbf{a} + s\mathbf{b}$  then  $p = r$  and  $q = s$

### Modelling with vectors

In mechanics, vectors have both magnitude and direction. For example velocity, displacement, and force.

Magnitude of a vector is a scalar quantity - it has size but no direction

| Vector quantity | Its magnitude |
|-----------------|---------------|
| velocity        | speed         |
| displacement    | distance      |

**Example 9:** A girl walks 2km due east from fixed point  $O$  to  $A$ , and then 3km due south from  $A$  to  $B$ . Find: i. The position vector of  $B$  relative to  $O$  ii. the bearing of  $B$  from  $O$

i) To find the position vector of  $B$  relative to  $O$ ,


We will have to represent the girl's journey on a diagram

$$\overrightarrow{OB} = p\mathbf{i} + q\mathbf{j}$$

2km due east implies  $p = 2$  and 3km due south

implies  $q = -3$ , you can refer to the diagram here

$$\text{So } \overrightarrow{OB} = (2\mathbf{i} - 3\mathbf{j}) \text{ km}$$



ii) To find bearing of  $B$  from  $O$ , you will have to calculate angle clockwise from North of  $O$  to

line segment  $OB$  which is  $90^\circ + \text{angle } \theta$

Using  $\tan \theta$  we can find angle  $\theta$

$$\tan \theta = \frac{3}{2}$$

$$\theta = \tan^{-1} \frac{3}{2} = 56.3^\circ$$

So the bearing of  $B$  from  $O$  is  $90^\circ + 56.3^\circ = 146.3^\circ = 146^\circ$

**Example 10:** Find the distance moved by particle which travels for 10 seconds at velocity  $(5\mathbf{i} - \mathbf{j}) \text{ ms}^{-1}$

You are familiar with the formula for distance, Distance = speed  $\times$  time

In order to calculate distance we need to find speed (i.e. magnitude of velocity)

and substitute speed and time = 10 seconds in the formula.

Use the magnitude formula to calculate speed

$$\text{Speed} = |5\mathbf{i} - \mathbf{j}| = \sqrt{5^2 + (-1)^2} = \sqrt{25 + 1} = \sqrt{26} = 5.099$$

$$\text{Hence, Distance} = 5.099 \times 10 = 51.0\text{m}$$