

STATISTICS (C) UNIT 1**TEST PAPER 9**

1. Briefly describe what is meant by
- (i) a statistical model, [2]
 - (ii) a refinement of a model. [2]
2. Every week a social club holds a raffle in which each ticket has a probability of 0.05 that it will win a prize. Marjory decides to buy one ticket every week until she wins a prize, after which she will stop taking part. Find the probability that
- (i) she wins a prize in the third week, [3]
 - (ii) she is still taking part in the fifth week. [2]
3. The random variable X has the binomial distribution $B(n, p)$.
Given that $E(X) = k \text{ Var}(X)$, find an expression for p in terms of k . [3]
If $n = 8$ and $k = 4$, find
- (i) $P(X = 5)$, (ii) $P(X > 6)$. [5]

4. The length of time, in minutes, that visitors queued for a tourist attraction is given by the following table, where, for example, '20 - ' means from 20 up to but not including 30 minutes.

Queuing time (mins)	0 -	10 -	15 -	20 -	30 -	40 - 60
Number of visitors	15	24	x	13	10	y

A histogram is drawn to represent this data. The total area under the histogram is 36 cm^2 . The '10 - ' bar has width 1 cm and height 9.6 cm. The '15 - ' bar is ten times as high as the '40 - 60' bar.

- (i) Find the values of x and y . [6]
 - (ii) On graph paper, construct the histogram accurately. [4]
5. The discrete random variable X takes only the values 4, 5, 6, 7, 8 and 9. The probabilities of these values are given in the table:

x	4	5	6	7	8	9
$P(X = x)$	p	0.1	q	q	0.3	0.2

It is known that $E(X) = 6.7$. Find

- (i) the values of p and q , [6]
- (ii) $P(X < 7)$, [2]
- (iii) $\text{Var}(X)$. [3]

STATISTICS 1 (C) TEST PAPER 9 Page 2

6. Among the families with two children in a large city, the probability that the elder child is a boy is $\frac{5}{12}$ and the probability that the younger child is a boy is $\frac{9}{16}$. The probability that the younger child is a girl, given that the elder child is a girl, is $\frac{1}{4}$.

One of the families is chosen at random. Using a tree diagram, or otherwise,

(i) show that the probability that both children are boys is $\frac{1}{8}$. [4]

(ii) Find the probability that one child is a boy and the other is a girl. [3]

If three of the families are chosen at random,

(iii) find the probability that exactly two of the families have two boys. [3]

(iv) State an assumption that you have made in answering part (iii). [1]

7. The marks out of 75 obtained by a group of ten students in their first and second Statistics modules were as follows:

Student	A	B	C	D	E	F	G	H	I	J
Module 1 (x)	54	33	42	71	60	27	39	46	59	64
Module 2 (y)	50	22	44	58	42	19	35	46	55	60

(i) Find $\sum x$ and $\sum y$. [2]

Given that $\sum x^2 = 26\,353$ and $\sum xy = 22\,991$,

(ii) obtain the equation of the regression line of y on x . [3]

(iii) Estimate the Module 2 result of a student whose mark in Module 1 was (a) 65, (b) 5.

Explain why one of these estimates is less reliable than the other. [3]

The equation of the regression line of x on y is $x = 0.921y + 9.81$.

(iv) Deduce the product moment correlation coefficient between x and y , and briefly interpret its value. [3]

STATISTICS 1 (C) TEST PAPER 9 : ANSWERS AND MARK SCHEME

1. (i) A mathematical representation which uses probabilities to describe and predict the behaviour of a real-life situation B2
- (ii) An improved mathematical formulation of the problem which aims to represent the reality more closely. B2 4
2. (i) $P(\text{wins in third week}) = 0.95 \times 0.95 \times 0.05 = 0.045$ M1 A1 A1
- (ii) $0.95^4 = 0.815$ M1 A1 5
3. $np = knp(1-p)$ $k(1-p) = 1$ $p = 1 - \frac{1}{k}$ M1 A1 A1
- (i) $p = 0.75$ $P(X \leq 5) - P(X \leq 4) = 0.3215 - 0.1138 = 0.208$ B1 M1 A1
- (ii) $P(X > 6) = 1 - P(X \leq 6) = 1 - 0.6329 = 0.367$ M1 A1 8
4. (i) 10 - 15 has area 9.6 cm^2 , so 2.5 visitors : 1 cm^2 B1 M1
- and $36 \text{ cm}^2 = 90$ visitors $62 + x + y = 90$ $x + y = 28$ M1 A1
- Also $\frac{x}{2.5} = 10 \times \frac{y}{10}$ so $x = 2.5y$ Hence $x = 20, y = 8$ M1 A1
- (ii) Freq. densities 1.5, 4.8, 4, 1.3, 1, 0.4 Histogram drawn B1 B3 10
5. (i) $4p + 13q + 4.7 = 6.7$ $4p + 13q = 2$ M1 A1
- $p + 2q + 0.6 = 1$ $p + 2q = 0.4$ B1
- Solve: $p = 0.24, q = 0.08$ M1 A1 A1
- (ii) $P(X < 7) = p + q + 0.1 = 0.42$ M1 A1
- (iii) $E(X^2) = 48.54$ $\text{Var}(X) = 48.54 - 6.7^2 = 3.65$ M1 A1 A1 11
6. (i) Let $P(\text{younger child is a boy, given elder is a boy}) = p$
- $\frac{5}{12}p + \frac{7}{12} \times \frac{3}{4} = \frac{9}{16}$ $\frac{5}{12}p = \frac{1}{8}$ $p = \frac{3}{10}$ M1 A1
- $P(B, B) = \frac{5}{12} \times \frac{3}{10} = \frac{1}{8}$ M1 A1
- (ii) $P(B, G \text{ or } G, B) = \frac{5}{12} \times \frac{7}{10} + \frac{7}{12} \times \frac{3}{4} = \frac{35}{48}$ or 0.729 M1 A1 A1
- (iii) $\frac{1}{8} \times \frac{1}{8} \times \frac{7}{8} \times 3 = \frac{21}{512}$ or 0.0410 M1 A1 A1
- (iv) Assumed independence B1 11
7. (i) $\sum x = 495, \sum y = 431$ B1 B1
- (ii) $S_{xx} = 1850.5, S_{yy} = 1656.5$ $y - 43.1 = 0.895(x - 49.5)$ M1 A1
- $y = 0.895x - 1.21$ A1
- (iii) (a) 57, (b) 3 B1 B1
- (b) is less reliable as it is well outside the range of given values B1
- (iv) $r = \sqrt{(0.895 \times 0.921)} = 0.908$ Quite good positive correlation M1 A1 B1 11