

GCE

Edexcel GCE

Statistics S1 (6683)

Summer 2005

advancing learning, changing lives

Mark Scheme (Results)

June 2005
6683 Statistics S1
Mark Scheme

| Question Number | Scheme | Marks |
|-----------------|---|---|
| 1. | <p>Diagram A : y & x : $r = -0.79$; As x increases, y decreases <u>or</u> most points lie in the 2nd and 4th quadrant.</p> <p>Diagram B : v & u : $r = 0.08$; No real pattern. Several values of v for one value of u or points lie in all four quadrants, randomly scattered.</p> <p>Diagram C : t & s : $r = 0.68$; As s increases, t increases or most points lie in the 1st and 3rd quadrants</p> | <p>B1;B1dep</p> <p>B1;B1dep</p> <p>B1;B1dep (6)</p> |
| 2. (a) | Distance is a continuous. | continuous B1 (1) |
| (b) | F.D = freq/class width \Rightarrow 0.8, 3.8, 5.3, 3.7, 0.75, 0.1 | or the same multiple of M1 A1 (2) |
| (c) | $Q_2 = 50.5 + \frac{(67 - 23)}{53} \times 10 = 58.8$ $Q_1 = 52.48; \quad Q_3 = 67.12$ | <p>awrt 58.8/58.9 M1 A1</p> <p>awrt 52.5/52.6 67.1/67.3 A1 A1 (4)</p> |
| | Special case : no working B1 B1 B1 (\equiv A's on the open) | |
| (d) | $\bar{x} = \frac{8379.5}{134} = 62.5335\dots$ $s = \sqrt{\frac{557489.75}{134} - \left(\frac{8379.5}{134}\right)^2}$ $s = 15.8089\dots \quad (S_{n-1} = 15.86825\dots)$ | <p>awrt 62.5 B1</p> <p>M1 A1√</p> <p>awrt 15.8 (15.9) A1 (4)</p> |
| | Special case : answer only B1 B1 (\equiv A's on the open) | |
| (e) | $\frac{Q_3 - 2Q_2 + Q_1}{Q_3 - Q_1} = \frac{67.12 - 2 \times 58.8 + 52.48}{67.12 - 52.48}$ $= 0.1366 \Rightarrow ; +ve \text{ skew}$ | <p>subst their Q_1, Q_2 & Q_3 need to show working for A1√ and have reasonable values for quartiles awrt 0.14 M1 A1√ A1; B1 (4)</p> |
| (f) | <p>For +ve skew Mean > Median & $62.53 > 58.80$ <u>or</u> $Q_3 - Q_2 (8.32) > Q_2 - Q_1 (6.32)$ Therefore +ve skew</p> | <p>B1 (1)</p> |

| Question Number | Scheme | Marks |
|-----------------|--|--|
| 3. (a) | $S_{xy} = 8880 - \frac{130 \times 48}{8} = (8100)$ <p style="text-align: right;">may be implied</p> $S_{xx} = 20487.5$ $b = \frac{s_{xy}}{s_{xx}} = \frac{8100}{20487.5} = 0.395363\dots$ <p style="text-align: right;">allow use of their S_{xy} for M awrt 0.395</p> $a = \frac{48}{8} - (0.395363\dots) \frac{130}{8} = -0.424649\dots$ <p style="text-align: right;">allow use of their b for M awrt -0.425</p> $y = -0.425 + 0.395x$ <p style="text-align: right;">3s.f.</p> <p>Special case answer only B0 M0 B1 M0 B1 B1 (fully correct 3sf) (\equiv to B0 M0 A1 M0 A1 B1 on the open)</p> | <p>B1</p> <p>M1 A1</p> <p>M1 A1</p> <p>B1 \checkmark</p> <p>(6)</p> |
| (b) | $f - 100 = -0.424649\dots + 0.395\dots(m - 250)$ $f = 0.735 + 0.395m$ | <p style="text-align: right;">subst f - 100 & m - 250</p> <p>M1 A1 \checkmark</p> <p style="text-align: right;">3 s.f.</p> <p>A1</p> <p>(3)</p> |
| (c) | $m = 235 \Rightarrow f = 93.64489\dots$ | <p style="text-align: right;">awrt 93.6/93.7</p> <p>B1</p> <p>(1)</p> |

| | | |
|-------------|---|--|
| <p>4(a)</p> | <p> $1.5 (Q_3 - Q_1) = 1.5 (28 - 12) = 24$ $Q_3 + 24 = 52 \Rightarrow 63$ is an outlier $Q_1 - 24 < 0 \Rightarrow$ no outliers </p> | <p> may be implied att $Q_3 + \dots$ or $Q_1 - \dots$, 52 and -12 or < 0 or evidence of no lower outliers 63 is an outlier B1 M1, A1 A1 M1 A1 A1 (7) </p> |
| <p>(b)</p> | <p>Distribution is +ve skew; $Q_2 - Q_1 (5) < Q_3 - Q_2 (11)$;</p> | <p>B1; B1 (2)</p> |
| <p>(c)</p> | <p>Many delays are small so passengers should find these acceptable or sensible comment in the context of the question.</p> | <p>B1 (1)</p> |

| | | | |
|--------------|---|---|-----------------------------------|
| <p>5.(a)</p> | $k + 2k + 3k + 5k + 6k = 1$ $17k = 1$ $k = \frac{1}{17} = 0.0588$ | <p>use of $\sum P(X = x) = 1$</p> | <p>M1</p> <p>A1</p> <p>(2)</p> |
| <p>(b)</p> | $E(X) = 1 \times \frac{1}{17} + 2 \times \frac{2}{17} + \dots + 5 \times \frac{6}{17} = \frac{64}{17}$ $= 3 \frac{13}{17}$ | <p>use of $\sum xP(X = x)$ and at least 2 prob correct</p> <p>Do not ignore subsequent working</p> | <p>M1</p> <p>A1</p> |
| <p>(c)</p> | $E(X^2) = 1^2 \times \frac{1}{17} + 2^2 \times \frac{2}{17} + \dots + 5^2 \times \frac{6}{17} = \left(\frac{266}{17} = 15.6 \right)$ $\text{Var}(X) = \frac{266}{17} - \left(\frac{64}{17} \right)^2$ $= 1.4740\dots$ | <p>use of $\sum x^2 P(X = x)$</p> <p>and at least 2 prob correct</p> <p>use of $\sum x^2 P(X = x) -$</p> <p>awrt 1.47</p> | <p>M1 A1</p> <p>A1</p> <p>(4)</p> |
| <p>(d)</p> | $\text{Var}(4 - 3X) = 9 \text{Var}(X) = 9 \times 1.47 = 13.23 \Rightarrow 13.2$ $\text{or } 9 \times 1.4740\dots = 13.266 \Rightarrow 13.3$ | <p>cao $9 \text{Var} X$</p> | <p>M1 A1</p> <p>(2)</p> |

| <p>6(a)</p> <p>(b)</p> <p>(c)</p> | <p>$M \sim N(155, 3.5^2)$</p> <p>$P(M > 160) = P\left(z > \frac{160-155}{3.5}\right)$ $= P(z > 1.43)$ $= 0.0764$</p> <p>$P(150 \leq M \leq 157) = P(-1.43 \leq z \leq 0.57)$ $= 0.7157 - (1 - 0.9236)$ $= 0.6393$</p> <p>special case : answer only B0 B0 M1 A1</p> <p>$P(M \leq m) = 0.3 \Rightarrow \frac{m-155}{3.5} = -0.5244$ $m = 153.2$</p> | <p>standardising $\pm(160-155), \sigma, \sigma^2, \sqrt{\sigma}$</p> <p>M1 A1 A1 (3)</p> <p>awrt -1.43, 0.57 $p > 0.5$ 0.6393 - 0.6400 4dp B1 B1 M1 A1 (4)</p> <p>-0.5244 att stand = z value for A1 may use awrt to - 0.52. cao B1 M1 A1 A1 (4)</p> | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|--|--|-----------|-----------------------|--------|--|---------|----|-----------|----|--|------|-----------|-----------|-----------|-----------------------|------------|----|-----------|----|-----------------------|--------|----|-----------|-----|--|---|
| <p>7.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> | <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Glasses</th> <th style="text-align: center;">No Glasses</th> <th style="text-align: center;">Totals</th> <th></th> </tr> </thead> <tbody> <tr> <td>Science</td> <td style="text-align: center;">18</td> <td style="text-align: center;">12</td> <td style="text-align: center;">30</td> <td></td> </tr> <tr> <td>Arts</td> <td style="text-align: center;">27</td> <td style="text-align: center;">23</td> <td style="text-align: center;">50</td> <td>50 may be seen in (a)</td> </tr> <tr> <td>Humanities</td> <td style="text-align: center;">44</td> <td style="text-align: center;">24</td> <td style="text-align: center;">68</td> <td>23 may be seen in (b)</td> </tr> <tr> <td>Totals</td> <td style="text-align: center;">89</td> <td style="text-align: center;">59</td> <td style="text-align: center;">148</td> <td></td> </tr> </tbody> </table> <p>$P(\text{Arts}) = \frac{50}{148} = \frac{25}{74} = 0.338$</p> <p>$P(\text{No glasses} / \text{Arts}) = \frac{23/148}{50/148} = \frac{23}{50} = 0.46$</p> <p>$P(\text{Right Handed}) = \left(\frac{30}{148} \times 0.8\right) + \left(\frac{50}{148} \times 0.7\right) + \left(\frac{68}{148} \times 0.75\right)$ $= \frac{55}{74} = 0.743$</p> <p>$P(\text{Science} / \text{Right handed}) = \frac{\frac{30}{148} \times 0.8}{\frac{55}{74}} = \frac{12}{55} = 0.218$</p> | | Glasses | No Glasses | Totals | | Science | 18 | 12 | 30 | | Arts | 27 | 23 | 50 | 50 may be seen in (a) | Humanities | 44 | 24 | 68 | 23 may be seen in (b) | Totals | 89 | 59 | 148 | | <p>B1 B1</p> <p>a number/148 M1 A1 (4)</p> <p>prob their(a)prob or number their 50 M1 A1 (2)</p> <p>attempt add three prob A1 ✓ on their (a) awrt 0.743 A1 (3)</p> <p>✓ on their (c) M1 A1 ✓ A1 (3)</p> |
| | Glasses | No Glasses | Totals | | | | | | | | | | | | | | | | | | | | | | | | |
| Science | 18 | 12 | 30 | | | | | | | | | | | | | | | | | | | | | | | | |
| Arts | 27 | 23 | 50 | 50 may be seen in (a) | | | | | | | | | | | | | | | | | | | | | | | |
| Humanities | 44 | 24 | 68 | 23 may be seen in (b) | | | | | | | | | | | | | | | | | | | | | | | |
| Totals | 89 | 59 | 148 | | | | | | | | | | | | | | | | | | | | | | | | |

