



# Mathematics (MEI)

Advanced Subsidiary GCE

Unit 4766: Statistics 1

# Mark Scheme for June 2013

PMT

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2013

PMT

June 2013

### Annotations and abbreviations

Annotation in scoris	Meaning
√and <b>×</b>	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
<b>^</b>	Omission sign
MR	Misread
Highlighting	

Other abbreviations in mark	Meaning
scheme	
E1	Mark for explaining
U1	Mark for correct units
G1	Mark for a correct feature on a graph
M1 dep*	Method mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working

June 2013

## Subject-specific Marking Instructions for GCE Mathematics (MEI) Statistics strand

a. Annotations should be used whenever appropriate during your marking.

The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded.

For subsequent marking you must make it clear how you have arrived at the mark you have awarded.

b. An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct *solutions* leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly.

Correct but unfamiliar or unexpected methods are often signalled by a correct result following an *apparently* incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, award marks according to the spirit of the basic scheme; if you are in any doubt whatsoever (especially if several marks or candidates are involved) you should contact your Team Leader.

c. The following types of marks are available.

Μ

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, eg by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

## Α

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

### В

Mark for a correct result or statement independent of Method marks.

## Ε

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, eg wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d. When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep \*' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e. The abbreviation ft implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only — differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, exactly what is acceptable will be detailed in the mark scheme rationale. If this is not the case please consult your Team Leader.

Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.

f. Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.

Candidates are expected to give numerical answers to an appropriate degree of accuracy. 3 significant figures may often be the norm for this, but this always needs to be considered in the context of the problem in hand. For example, in quoting probabilities from Normal tables, we generally expect *some* evidence of interpolation and so quotation to 4 decimal places will often be appropriate. But even this does not always apply – quotations of the standard critical points for significance tests such as 1.96, 1.645, 2.576 (maybe even 2.58 – but not 2.57) will commonly suffice, especially if the calculated value of a test statistic is nowhere near any of these values. Sensible discretion *must* be exercised in such cases.

#### Mark Scheme

June 2013

Discretion must also be exercised in the case of small variations in the degree of accuracy to which an answer is given. For example, if 3 significant figures are expected (either because of an explicit instruction or because the general context of a problem demands it) but only 2 are given, loss of an accuracy ("A") mark is likely to be appropriate; but if 4 significant figures are given, this should not normally be penalised. Likewise, answers which are slightly deviant from what is expected in a very minor manner (for example a Normal probability given, after an attempt at interpolation, as 0.6418 whereas 0.6417 was expected) should not be penalised. However, answers which are *grossly* over- or under-specified should normally result in the loss of a mark. This includes cases such as, for example, insistence that the value of a test statistic is (say) 2.128888446667 merely because that is the value that happened to come off the candidate's calculator. Note that this applies to answers that are given as final stages of calculations; intermediate working should usually be carried out, and quoted, to a greater degree of accuracy to avoid the danger of premature approximation.

The situation regarding any particular cases where the accuracy of the answer may be a marking issue should be detailed in the mark scheme rationale. If in doubt, contact your Team Leader.

g. Rules for replaced work

If a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests.

If there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others.

NB Follow these maths-specific instructions rather than those in the assessor handbook.

h. Genuine misreading (of numbers or symbols, occasionally even of text) occurs. If this results in the object and/or difficulty of the question being considerably changed, it is likely that all the marks for that question, or section of the question, will be lost. However, misreads are often such that the object and/or difficulty remain substantially unaltered; these cases are considered below.

The simple rule is that *all* method ("M") marks [and of course all independent ("B") marks] remain accessible but at least some accuracy ("A") marks do not. It is difficult to legislate in an overall sense beyond this global statement because misreads, even when the object and/or difficulty remains unchanged, can vary greatly in their effects. For example, a misread of 1.02 as 10.2 (perhaps as a quoted value of a sample mean) may well be catastrophic; whereas a misread of 1.6748 as 1.6746 may have so slight an effect as to be almost unnoticeable in the candidate's work.

#### **Mark Scheme**

A misread should normally attract *some* penalty, though this would often be only 1 mark and should rarely if ever be more than 2. Commonly in sections of questions where there is a numerical answer either at the end of the section or to be obtained and commented on (eg the value of a test statistic), this answer will have an "A" mark that may actually be designated as "cao" [correct answer only]. This should be interpreted *strictly* – if the misread has led to failure to obtain this value, then this "A" mark must be withheld even if all method marks have been earned. It will also often be the case that such a mark is implicitly "cao" even if not explicitly designated as such.

On the other hand, we commonly allow "fresh starts" within a question or part of question. For example, a follow-through of the candidate's value of a test statistic is generally allowed (and often explicitly stated as such within the marking scheme), so that the candidate may exhibit knowledge of how to compare it with a critical value and draw conclusions. Such "fresh starts" are not affected by any earlier misreads.

A misread may be of a symbol rather than a number – for example, an algebraic symbol in a mathematical expression. Such misreads are more likely to bring about a considerable change in the object and/or difficulty of the question; but, if they do not, they should be treated as far as possible in the same way as numerical misreads, *mutatis mutandis*. This also applied to misreads of text, which are fairly rare but can cause major problems in fair marking.

The situation regarding any particular cases that arise while you are marking for which you feel you need detailed guidance should be discussed with your Team Leader.

Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

(	Questio	n	Answer	Marks		Guidance
1	(i)		Mean = $\frac{24940}{100}$ = 249.4g or 249g	B1	Ignore units	CAO NB 249.40 gets B0 for over- specification
			$Sxx = 6240780 - \frac{24940^2}{100} = 20744$	M1	For S <i>xx</i>	M1 for 6240780 - $100 \times \text{their mean}^2$ BUT NOTE M0 if their $S_{xx} < 0$
			$s = \sqrt{\frac{20744}{99}} = \sqrt{209.53} = 14.4751 = 14.5g$	A1	CAO ignore units	For s <sup>2</sup> of 210 (or better) allow M1A0 with or without working For RMSD of 14.4 (or better) allow M1A0 provided working seen For RMSD <sup>2</sup> of 207 (or better) allow M1A0 provided working seen Allow 14.48 but NOT 14.47
				[3]		
1	(ii)		New mean = $(0.9 \times 249.4) - 15 = 209.5g$	B1	FT their mean provided answer is positive	If candidate 'starts again' only award marks for CAO Allow 209
			New sd = $0.9 \times 14.48 = 13.03$ g	M1	FT their sd	Or for $0.9^2 \times 14.5^2$
				A1	FT Allow 13.0 to 13.1	Deduct at most 1 mark overall in whole question for over-specification of Mean and 1 mark overall for SD
				[3]		

(	Juestio	n	Answer	Marks		Guidance
2	(i)		$3 \times \frac{5}{10} \times \frac{4}{9} \times \frac{5}{8} = \frac{300}{720} = \frac{5}{12} = (0.4167)$	M1	For $5/10 \times 4/9$	Correct working but then multiplied or divided by some factor scores M1M1M0A0
				M1	For $\times$ 5/8	Zero for binomial Allow M2 for equivalent triple such as $\frac{5}{10} \times \frac{5}{9} \times \frac{4}{8}$
				M1	For $3 \times$ triple product	Or 3 separate equal triplets added
				A1	CAO (Fully simplified)	Answer must be a fraction
				[4]		
			Or $\binom{5}{\times} \binom{5}{\times}$	M1*	For $\binom{5}{2} \times \binom{5}{1}$	Seen
			$\frac{\binom{2}{\binom{10}{3}}}{\binom{10}{3}} = \frac{10 \times 5}{120} = \frac{5}{12}$	M1*	For $\begin{pmatrix} 10\\ 3 \end{pmatrix}$	Seen
				M1*	For whole fraction	Correct working but then multiplied or
				dep		divided by some factor scores M1M1M0A0
				A1	CAO (Fully simplified)	
2	( <b>ii</b> )		$(, 7 (5)^3 (5)^4)$	M1FT	For first probability	Allow <sup>4</sup> C <sub>3</sub>
			$4 \times \frac{12}{12} \times \left(\frac{12}{12}\right) + \left(\frac{12}{12}\right)$	M1FT	For $(5/12)^4$	
				M1FT	For sum of both correct probabilities	Provided sum <1
			= 0.169 + 0.030 = 0.199	A1	CAO	Alternative for 1- $(P(0)+P(1)+P(2))$
			$Or = \frac{875}{625} + \frac{625}{625} - \frac{1375}{625}$		Do not allow 0.2, unless	allow M1FT for two 'correct' probs, M1
			5184 20736 6912		fuller answer seen first	answer, A1 CAO
				[4]		

4766

(	Question		Answer	Marks		Guidance
3	(i)		X ~ B(50, 0.1)	M1	For $0.1^5 \times 0.9^{45}$	
			P(5 underweight) = $\binom{50}{5} \times 0.1^5 \times 0.9^{45} = 0.1849$	M1	For $\binom{50}{5} \times p^5 \times q^{45}$	With $p + q = 1$ Also for 2118760 × 8.73 × 10 <sup>-8</sup>
				A1 [ <b>3</b> ]	CAO	Allow 0.185 or better <u>NB 0.18 gets A0</u>
3	(ii)		$X \sim B(20, 0.1)$ P(X \ge 1) = 1 - P(X = 0) = 1 - 0.1216 = 0.8784	M1 A1 [2]	For 0.1216 CAO	Allow M1 for 0.9 <sup>20</sup> Allow 0.878 or better See tables at the website <u>http://www.mei.org.uk/files/pdf/formula</u> <u>_book_mf2.pdf</u>
3	(iii)		E( <i>X</i> ) = 48 × 0.8784 = 42.16 (= 42.2)	M1 A1 [2]	FT their probability from part (ii)	If any indication of rounding to 42 or 43 or to another integer on FT allow M1A0 SC1 for 48 $\times$ their <i>p</i> giving an integer answer. NB 0.6083 in (ii) leads to 29.20

(	Questio	0 <b>n</b>	Answer	Marks		Guidance
4	(i)		$P(X = 15) = \frac{3}{6} \times \frac{2}{5} \times \frac{1}{4}$	M1	For product of three correct fractions	
			$= = \frac{6}{120} = \frac{1}{20} = 0.05$ Or $\frac{1}{10} = \frac{1}{10} = 0.05$	A1	NB ANSWER GIVEN	Full marks for $3! \times \frac{1}{6} \times \frac{1}{5} \times \frac{1}{4} = \frac{6}{120} = 0.05$
			$\mathbf{Or}  \frac{_{6}C_{3}}{6!} = \frac{1}{20} = 0.05$		= 0.05  scores M0A0	Allow 3 × 2 in place of 3! SC1 for $6 \times \frac{1}{6} \times \frac{1}{5} \times \frac{1}{4} = \frac{6}{120} = 0.05$
				[2]		
4	(ii)		$ E(X) = (15 \times 0.05) + (1010 \times 0.45) + (2005 \times 0.45) + (3000 \times 0.05) $	M1	For $\Sigma rp$ (at least 3 terms correct)	
			= 1507.5 so 1508 (4sf)	A1	CAO	Allow 1507, 1510, 1507.5, 1507.50 or 3015/2
			$E(X^{2}) = (15^{2} \times 0.05) + (1010^{2} \times 0.45) + (2005^{2} \times 0.45) + (3000^{2} \times 0.05)$ = 2718067.5	M1	For $\Sigma r^2 p$ (at least 3 terms correct)	Use of $E(X-\mu)^2$ gets M1 for attempt at $(x-\mu)^2$ should see $(-1492.5)^2$ , $(-497.5)^2$ , $497.5^2$ , $1492.5^2$ , (if $E(X)$ wrong FT their $E(X)$ ) (all 4 correct for M1), then M1 for $\Sigma p(x-\mu)^2$ (at least 3 terms correct with their probabilities) Division by 4 or other spurious value at end gives max M1A1M1M1A0, or M1A0M1M1A0 if $E(X)$ also divided by 4. Unsupported correct answers get 5 marks
			Var (X) = $2718067.5 - (1507.5)^2$	M1	dep for – their $E(X)^2$	
			= 445511.25 so 445500 (4sf)	A1	FT their E(X) provided Var(X) > 0 (and of course E( $X^2$ ) is correct)	Allow 446000
				[5]		

(	Question		Answer	Marks	Guidance		
5	(i)		Because if people cannot make a correct identification, then the probability that they guess correctly will be 0.5 For 'equally likely to guess right or wrong' or 'two outcomes with equal probability' or '50:50 chance of success' or 'right one in two occasions on average' or 'two (equally likely)	E1 E1	For idea of a guess or 'chosen at random' For idea of two outcomes	NB The question includes the sentence 'She suspects that people do no better than they would by guessing.', so this on its own does not get the mark for the idea of a guess	
			outcomes' etc	[2]			
5	(ii)		'Because people may do better than they would by guessing' or similar	B1	For idea of selecting correctly /identifying /knowing	No marks if answer implies that it is because there are over half in the sample who make a correct identification	
				[1]			
5	(iii)		$P(X \ge 13) = 1 - P(X \le 12) = 1 - 0.8684 = 0.1316$ NB PLEASE ANNOTATE THE TOP AND BOTTOM OF THE EXTRA PAGE IF NOT USED	M1	For notation $P(X \ge 13)$ or $P(X > 12)$ or $1 - P(X \le 12)$	Notation $P(X = 13)$ scores M0. If they have the correct $P(X \ge 13)$ then give M1 and ignore any further incorrect notation.	
				B1*	For 0.1316	Or for 1 – 0.8684 indep of previous mark	
			0.1316 > 0.05	M1* dep	For comparison with 5%		
			So not significant	A1*		Allow 'accept H <sub>0</sub> ' or 'reject H <sub>1</sub> '	
			There is insufficient evidence to suggest that people can make a correct identification.	E1* dep	NB Point probabilities score zero.	Must include 'insufficient evidence' or something similar such as 'to suggest that' ie an element of doubt either in the A or E mark. Must be in context to gain E1 mark. Do not allow 'sufficient evidence to suggest proportion making correct identification is 0.5' or similar	

June	2013	
------	------	--

Question	Answer	Marks		Guidance
	ALTERNATIVE METHOD – follow method above unless some mention of CR seen		Must see some reference to CR to gain any marks	
	Critical region method UPPER TAIL $P(X \ge 14) = 1 - P(X \le 13) = 1 - 0.9423 = 0.0577 > 5\%$ $P(X \ge 15) = 1 - P(X \le 14) = 1 - 0.9793 = 0.0207 < 5\%$ So critical region is {15,16,17,18,19,20}	B1 M1* M1* dep	For either probability For a correct comparison with 5% cao dep on the two correct probabilities	Do not insist on correct notation as candidates have to work out two probabilities for full marks. Allow comparison in form of statement 'critical region at 5% level is' No marks if CR not justified Condone {15, 20}, $X \ge 15$ , oe but not P( $X \ge 15$ ,) etc
	13 not in CR so not significant	A1*	Must include '13 not in CR'	Allow 'accept H <sub>0</sub> ' or 'reject H <sub>1</sub> '
	There is insufficient evidence to indicate that people can make a correct identification.	E1* dep on A1 [5]	Ignore any work on lower critical region	NB If CR found correctly, <b>then</b> <b>P(X=13) subsequently found</b> , but cand says '13 not in CR' then allow up to all five marks. If do not say '13 not in CR' allow no marks

C	Juestion	Answer	Marks		Guidance
6	(i)	Median = $3.32 \text{ kg}$ Q1 (= 6.5th value) = 2.83 Q3 (= 19.5th value) = 3.71 Inter-quartile range = $3.71 - 2.83 = 0.88$	B1 B1 B1 [ <b>3</b> ]	For Q1 or Q3 For IQR dep on both quartiles correct	For Q1 allow 2.82 to 2.84 For Q3 allow 3.70 to 3.72 If no quartiles given allow B0B1 for IOR in range 0.86 to 0.90
	(ii)	1 1.5 2 2.5 3 3.5 4 4.5 Weight (kg)	G1 G1 G1	For reasonably linear scale shown. For boxes in approximately correct positions, with median just to right of centre For whiskers in approximately correct positions in proportion to the box FT their median and quartiles if sensible – guidance above is only for correct values	Dep on attempt at box and whisker plot with at least a box and one whisker. Condone lack of label. Do not award unless RH whisker significantly shorter than LH whisker Allow LH whisker going to 2.5 and outlier marked at 1.39
			[3]		
6	(iii)	Lower limit $2.83 - (1.5 \times 0.88) = 1.51$	B1	For 1.51 FT	Any use of median $\pm 1.5 \times IQR$ scores B0 B0 E0 No marks for $\pm 2$ or $3 \times IQR$ In this part FT their values from (i)or (ii) if sensibly obtained but not from location ie 6.5, 19.5
		Upper limit $3.71 + (1.5 \times 0.88) = 5.03$	B1	For 5.03 FT	Do not penalise over-specification as
		Exactly one baby weighs less than 1.51 kg and none weigh over 5.03 kg so there is exactly one outlier.	E1*	Dep on their 1.51 and 5.03	Do not allow unless FT leads to upper limit above 4.34 and lower limit between 1.39 and 2.50

Question		n	Answer	Marks	Guidance	
			'Nothing to suggest that this baby is not a genuine data value so she should not be excluded' or 'This baby is premature and therefore should be excluded'.	E1* Dep	Any sensible comment in context	For use of mean $\pm$ 2sd allow B1 For $3.27 + 2 \times 0.62 = 4.51$ B1 For $3.27 - 2 \times 0.62 = 2.03$ Then E1E1 as per scheme
6	(iv)		Median = $3.5 \text{ kg}$ Q1 = 50th value = $3.12$ Q3 = $150th \text{ value } = 3.84$ Inter-quartile range = $3.84 - 3.12 = 0.72$	B1 B1 B1 [3]	For Q1 or Q3 For IQR FT their quartiles	For Q1 allow 3.11 to 3.13 For Q3 allow 3.83 to 3.85 Dep on both quartiles correct If no quartiles given allow B0B1 for
6	( <b>v</b> )		Female babies have lower weight than male babies on the whole Female babies have higher weight variation than male babies	E1 FT E1 FT	Allow 'on average' or similar in place of 'on the whole' Allow 'more spread' or similar but not 'higher range' Condone less consistent	IQR in range 0.70 to 0.74 Do not allow lower median Do not allow higher IQR, but SC1 for both lower median and higher IQR, making clear which is which
6	(vi)		Male babies must weigh more than 4.34 kg		E 10 0 0	0 000 100 000 101 000 100
			Approx 10 male babies weigh more than this. Probability = $\frac{10}{200} \times \frac{9}{199} = \frac{90}{39800} = \frac{9}{3980} = 0.00226$ or $\frac{9}{200} \times \frac{8}{199} = \frac{72}{39800} = 0.00181$ or $\frac{8}{200} \times \frac{7}{199} = \frac{56}{39800} = \frac{7}{4975} = 0.00141$	M1* M1* dep A1	For 10 or 9 or 8 For first fraction multiplied by any other different fraction (Not a binomial probability) CAO Allow their answer to min of 2 sf	Or $200 - 190$ , $200 - 191$ or $200 - 192$ Allow any of these answersFor spurious factors, eg 2 × correctanswer allow M1M1A0SC1 for $n/200 \times (n-1)/199$

4766

Question		on	Answer	Marks	Guidance	
7	Questic	on	AnswerThirdSecond $0.2$ Hit $0.1$ Hit $0.2$ Hit $0.1$ Hit $0.2$ Hit $0.1$ Hit $0.2$ Hit $0.1$ Hit $0.05$ Hit $0.1$ Hit $0.2$ Hit $0.95$ Miss $0.2$ Hit $0.95$ Miss $0.2$ Hit $0.95$ Miss $0.2$ Hit $0.95$ Miss $0.05$ Hit $0.95$ Miss $0.05$ Hit $0.95$ Miss $0.95$ Miss $0.95$ Miss $0.95$	Marks G1 G1 G1 G1 [4]	For first set of branches For second set of branches (indep) For third set of branches (indep) For labels	Guidance         All probabilities correct         All probabilities correct         All probabilities correct         All correct labels for 'Hit' and 'Miss', 'H' and 'M' etc. Condone omission of First, Second, Third.         Do not allow misreads here as all FT
			ALTERNATIVE METHOD only if there is an attempt to add 7 probabilities At least three correct triple products Attempt to add 7 triple products FURTHER ALTERNATIVE METHOD $0.1 + 0.9 \times 0.05$ Above probability $+ 0.9 \times 0.95 \times 0.05$	M1 M1* dep A1 M1 M1 A1 M1 A1 M1 A1 [3]	For 1 – ans CAO CAO	<ul> <li>0.188 or better. Condone 0.1877</li> <li>Allow 751/4000</li> <li>(not necessarily correct triple products)</li> </ul>

Question		on	Answer	Marks	Guidance	
7	(ii)	B	P(Hits with exactly one)	M1	For two correct products	FT their tree for all three M marks,
			$= (0.1 \times 0.8 \times 0.95) + (0.9 \times 0.05 \times 0.8) + (0.9 \times 0.95 \times 0.05)$	M1	For all three correct products	provided three terms
			$= 0.076 + 0.036 + 0.04275 = \frac{19}{250} + \frac{9}{250} + \frac{171}{4000}$	M1	For sum of all three correct products	
			$=\frac{619}{4000}=0.15475$	A1	CAO	Allow 0.155 or better
				[4]		
7	(iii)		P(Hits with exactly one given hits with at least one)			If answer to $(B)$ > than answer to $(A)$
			$-\frac{P(\text{Hits with exactly one and hits with at least one})}{P(\text{Hits with exactly one and hits with at least one})}$			then max M1M0A0
			P(Hits with at least one)			
			0 15475	M1	For numerator FT	Both must be part of a fraction
			$=\frac{0.18775}{0.18775}$	M1	For denominator FT	1
			- 0.8242	Δ1	CAO	Allow 0.824 or better or 619/751
			- 0.0242	[3]		
7	(iv)		P(Hits three times overall) = $(0.1 \times 0.2 \times 0.2) + (0.9 \times 0.95 \times 0.95 \times 0.05 \times 0.2 \times 0.2)$	M1	For $0.1 \times 0.2 \times 0.2$ or $0.004$ or $1/250$	FT their tree for all three M marks
				M1	For $0.9 \times 0.95 \times 0.95 \times 0.05 \times 0.05 \times 0.2 \times 0.2$	provided three terms in first product and six in second product. Last three probs must be $0.05 \times 0.2 \times 0.2$ unless they extend their tree
			= 0.004 + 0.0016245	M1*	For sum of both	With no extras
				Dep		
				on h ath		
				both		
				M1's		
			= 0.0056245	A1	CAO	Allow 0.00562 or 0.00563 or 0.0056
				[4]		

## NOTE RE OVER-SPECIFICATION OF ANSWERS

If answers are grossly over-specified, deduct the final answer mark in every case. Probabilities should also be rounded to a sensible degree of accuracy. In general final non probability answers should not be given to more than 4 significant figures. Allow probabilities given to 5 sig fig.

## PLEASE HIGHLIGHT ANY OVER-SPECIFICATION

Please note that there are no G or E marks in scoris, so use B instead

# NB PLEASE ANNOTATE EVERY ADDITIONAL ANSWER SHEET EVEN IF FULL MARKS AWARDED OR THE PAGE IS BLANK

## Additional notes re Q5 part iii

Comparison with 95% method If 95% seen anywhere then M1 for P( $X \le 12$ ) B1 for 0.8684 M1\* for comparison with 95% dep on second B1 A1\* for not significant oe E1\*

Comparison with 95% CR method If 95% seen anywhere then B1 for 0.9423 or 0.9793 M1 for correct comparison with 95% M1dep for correct CR provided both probs correct then follow mark scheme for CR method

Smallest critical region method: Smallest critical region that 13 could fall into is {13, 14, 15, 16, 17, 18, 19, 20} gets B1 and has size 0.1316 gets B1, This is > 5% gets M1\*, A1\*, E1\* as per scheme NB These marks only awarded if 13 used, not other values.

<u>Use of *k* method with no probabilities quoted:</u> This gets zero marks.

Use of *k* method with one probability quoted: Mark as per scheme

Line diagram method and Bar chart method

No marks unless correct probabilities shown on diagram, then mark as per scheme..

PMT

OCR (Oxford Cambridge and RSA Examinations) 1 Hills Road Cambridge CB1 2EU

**OCR Customer Contact Centre** 

#### **Education and Learning**

Telephone: 01223 553998 Facsimile: 01223 552627 Email: general.qualifications@ocr.org.uk

#### www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee Registered in England Registered Office; 1 Hills Road, Cambridge, CB1 2EU Registered Company Number: 3484466 OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations) Head office Telephone: 01223 552552 Facsimile: 01223 552553 MAT OF THE CAMERIDGE ASSESSMENT

