


Question Number	Scheme	Marks
1.	<p>(a) Survey is less time consuming.</p> <p>(b) It is easier/quicker to analyse the results</p> <p>(c) List of members</p> <p>(d) The members</p>	<p>B1</p> <p>B1 (2)</p> <p>B1 (1)</p> <p>B1 (1)</p> <p>(4 marks)</p>
2.	<p>(a) Y is the random variable consisting of any function of the X_i that involves no other quantities.</p> <p>(b) $Y = \bar{X} = \frac{\sum X}{n}$</p> <p>(c) When all possible samples are taken and the values of Y found then the values form a probability distribution (known as the sampling distribution of Y)</p>	<p>B1 B1 (2)</p> <p>B1 (1)</p> <p>B1 B1 (2)</p> <p>(5 marks)</p>
3.	<p>(a) $E(R) = \frac{\alpha + \beta}{2} = 3, \Rightarrow \alpha + \beta = 6$</p> <p>(b) $\text{Var}(R) = \frac{(\beta - \alpha)^2}{12} = \frac{25}{3}, \Rightarrow (\beta - \alpha)^2 = 100$</p> <p>$\alpha = -2, \beta = 8$</p> <p>$P(R < 6.6) = \frac{1}{10} \times 8.6 = 0.86$</p>	<p>M1 A1</p> <p>M1 A1</p> <p>M1 A1 A1 (7)</p> <p>M1 A1 (2)</p> <p>(9 marks)</p>
4.	<p>(a) $H_0 : \rho = 0.20, H_1 : \rho < 0.20$</p> <p>$X =$ number buying single packets, $X \sim B(25, 0.20)$</p> <p>$P(X \leq 2) = 0.0982$</p> <p>$0.0982 > 5\%$, so not significant (comparison)</p> <p>No reason to suspect the percentage who bought crisps in single packets that day was lower than usual (context)</p> <p>$H_0 : \rho = 0.03, H_1 : \rho \neq 0.03$</p> <p>$Y =$ number buying bumper packs, $Y \sim B(300, 0.03) \Rightarrow Y \sim \text{Po}(9)$</p> <p>$P(Y \leq 3) = 0.0212$ and $P(Y \leq 15) = 0.9780 \Rightarrow P(Y \geq 16) = 0.0220$</p> <p>Critical region $Y \leq 3$ and $Y \geq 16$</p> <p>Significance level = $0.0212 + 0.0220 = 0.0432$</p>	<p>B1 B1</p> <p>M1 A1</p> <p>M1</p> <p>A1 ft (2)</p> <p>B1 B1</p> <p>M1</p> <p>M1 A1</p> <p>A1 (6)</p> <p>B1 ft (1)</p> <p>(13 marks)</p>

Question Number	Scheme	Marks
5.	<p>(a) $L \sim N(\mu, 0.3^2)$, $P(L < 150) = 0.05 \Rightarrow P\left(Z < \frac{150 - \mu}{0.3}\right) = 0.05$</p>  <p style="margin-left: 400px;">$\Rightarrow \frac{150 - \mu}{0.3} = -1.6449$</p> <p style="margin-left: 400px;">$\mu = 150.49347 = 150.5$</p> <p>(b) X represents number less than 150cm. $X \sim B(10, 0.05)$</p> <p style="margin-left: 400px;">$P(X \leq 2) = 0.9885$</p> <p>(c) Normal approximation $\mu = 500 \times 0.05 = 25$, $\sigma^2 = 23.75$ or 25</p> <p style="margin-left: 40px;">$P(X < 35) \approx P\left(Z < \frac{34.5 - 25}{\sqrt{23.75 \text{ or } 25}}\right)$ ± 0.5, standardise</p> <p style="margin-left: 100px;">$\approx P(Z < 1.95 \text{ or } 1.9)$</p> <p style="margin-left: 100px;">$\approx 0.9744 \text{ or } 0.9713$</p>	<p>M1 A1, B1</p> <p>A1 (4)</p> <p>B1</p> <p>M1 A1 (3)</p> <p>B1, B1</p> <p>M1, M1</p> <p>A1</p> <p>A1 (6)</p> <p style="text-align: right;">(13 marks)</p>
6.	<p>(a) X represents number of faults per 25 m $\Rightarrow X \sim \text{Po}(1.5)$</p> <p style="margin-left: 40px;">$P(X = 4) = 0.0471$</p> <p>(b) Y represents number of faults per 100 m $\Rightarrow Y \sim \text{Po}(6.0)$</p> <p style="margin-left: 40px;">$P(Y < 6) = P(Y \leq 5) = 0.4457$</p> <p>$R$ represents number of 100 m balls containing fewer than 6 faults</p> <p style="margin-left: 40px;">$R \sim B(3, 0.4457)$</p> <p style="margin-left: 40px;">$P(R = 1) = C_1^3 \times 0.4457 \times (1 - 0.4457)^2 = 0.41082$ accept 0.411</p> <p>(c) S represents number of faults in a 500 m ball $\Rightarrow S \sim \text{Po}(30)$</p> <p style="margin-left: 40px;">$P(23 \leq S \leq 33) \approx P\left(\frac{22.5 - 30}{\sqrt{30}} \leq Z \leq \frac{33.5 - 30}{\sqrt{30}}\right)$ ± 0.5, standardise</p> <p style="margin-left: 100px;">$\approx P(-1.37 \leq Z \leq 0.64)$</p> <p style="margin-left: 100px;">≈ 0.6536</p>	<p>B1</p> <p>B1 (2)</p> <p>B1</p> <p>B1</p> <p>M1 A1</p> <p>M1 A1 (6)</p> <p>B1</p> <p>M1, M1 A1</p> <p>A1</p> <p>A1 (6)</p> <p style="text-align: right;">(14 marks)</p>

Question Number	Scheme	Marks
7. (a)	<p>(b) (i) $F(x) = \int_0^x \frac{x}{15} dx = \frac{x^2}{30}$ for $0 \leq x \leq 2$</p> <p>$F(x) = \frac{12}{15} + \int_7^x \left(\frac{4}{9} - \frac{2x}{45}\right) dx = \frac{4x}{9} - \frac{x^2}{45} - \frac{11}{9}$ for $7 \leq x \leq 10$</p> <p>(ii) $F(x) = \frac{2}{15} + \int_2^x \frac{2}{15} dx = \frac{2x}{15} - \frac{2}{15}$ for $2 \leq x \leq 7$</p> <p>(iii) $F(x) = 0, x < 0, F(x) = 1, x > 10$</p> <p>(c) $P(X \leq 8.2) = F(8.2) = 0.928$</p> <p>(d) $E(X) = \int_0^2 \frac{x^2}{15} dx + \int_2^7 \frac{2x}{15} dx + \int_7^{10} \left(\frac{4x}{9} - \frac{2x^2}{45}\right) dx$</p> <p>$= \left[\frac{x^3}{45}\right]_0^2 + \left[\frac{x^2}{15}\right]_2^7 + \left[\frac{2x^2}{9} - \frac{2x^3}{125}\right]_7^{10} = 4.78$</p>	<p>B1 (labels) B1 (graph) B1 (axes)</p> <p>B1</p> <p>B1 M1 A1</p> <p>B1 M1 A1</p> <p>B1 (8)</p> <p>M1 A1 (2)</p> <p>M1 A1</p> <p>A1 A1 (4)</p> <p>(17 marks)</p>