

# Mark Scheme (Results) Summer 2010

GCE

GCE Statistics S2 (6684/01)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Summer 2010

Publications Code UA024768

All the material in this publication is copyright

© Edexcel Ltd 2010

June 2010  
Statistics S2 6684  
Mark Scheme

Question Number	Scheme	Marks
Q1	(a) A population is collection of all items	B1 (1)
	(b) (A random variable) that is a function of the sample which contains no unknown quantities/parameters.	B1 (1)
	(c) The voters in the town Percentage/proportion voting for Dr Smith	B1 B1 (2)
	(d) Probability Distribution of those voting for Dr Smith from all possible samples (of size 100)	B1 (1)
		[5]
	<p><b>Notes</b></p> <p>(a) <b>B1</b> – collection/group <b>all</b> items – need to have /imply all eg entire/complete/every</p> <p>(b) <b>B1</b> – needs <u>function/calculation(o.e.) of the sample/random variables/observations</u> <b>and no unknown quantities/parameters(o.e.)</b> NB do not allow unknown variables  e.g. “A calculation based <u>solely</u> on observations from a given sample.” B1  “A calculation based <u>only</u> on known data from a sample” B1  “A calculation based on known observations from a sample” B0</p> <p>(c) <b>B1</b> – Voters  Do not allow 100 voters.  <b>B1</b> – percentage/ proportion voting (for Dr Smith)  the <b>number</b> of people voting (for Dr Smith)  Allow 35% of people voting (for Dr Smith)  Allow 35 people voting (for Dr Smith)  Do <b>not</b> allow 35% or 35 alone</p> <p>(d) <b>B1</b> – answers must include all three of these features  (i) All possible samples,  (ii) their associated probabilities,  (iii) context of voting for Dr Smith.   e.g “It is all possible values of the percentage and their associated probabilities.” B0 no context</p>	Solely/only imply no unknown quantities

Question Number	Scheme	Marks
Q2 (a)	<p>Let <math>X</math> be the random variable the number of games Bhim loses.  <math>X \sim B(9, 0.2)</math></p> $P(X \leq 3) - P(X \leq 2) = 0.9144 - 0.7382 \quad \text{or} \quad (0.2)^3 (0.8)^6 \frac{9!}{3!6!}$ $= 0.1762 \quad \quad \quad = 0.1762$ <p>awrt 0.176</p> <p>(b) <math>P(X \leq 4) = 0.9804</math> awrt 0.98</p> <p>(c) Mean = 3    variance = 2.85, <math>\frac{57}{20}</math></p> <p>(d) Po(3) poisson</p> $P(X > 4) = 1 - P(X \leq 4)$ $= 1 - 0.8153$ $= 0.1847$	<p>B1</p> <p>M1 A1    (3)</p> <p>M1A1    (2)</p> <p>B1 B1    (2)</p> <p>M1</p> <p>M1</p> <p>A1    (3)</p> <p>[10]</p>
	<p><b>Notes</b></p> <p>(a) <b>B1</b> – writing or use of <math>B(9, 0.2)</math>  <b>M1</b> for writing/ using <math>P(X \leq 3) - P(X \leq 2)</math> or <math>(p)^3 (1-p)^6 \frac{9!}{3!6!}</math>  <b>A1</b> awrt 0.176</p> <p>(b) <b>M1</b> for writing or using <math>P(X \leq 4)</math>  <b>A1</b> awrt 0.98</p> <p>(c) <b>B1</b> 3  <b>B1</b> 2.85, or exact equivalent</p> <p>(d) <b>M1</b> for using Poisson  <b>M1</b> for writing or using <math>1 - P(X \leq 4)</math> NB <math>P(X \leq 4)</math> is 0.7254 Po(3.5) and 0.8912 Po(2.5)  <b>A1</b> awrt 0.185</p> <p><b>Special case :Use of Po(1.8) in (a) and (b)</b></p> <p>(a) can get B1 M1 A0 – B1 if written <math>B(9, 0.2)</math>, M1 for <math>\frac{e^{-1.8} 1.8^3}{3!}</math> <b>or</b> awrt to 0.161  If <math>B(9, 0.2)</math> is not seen then the only mark available for using Poisson is M1.  (b) can get M1 A0 - M1 for writing or using <math>P(X \leq 4)</math> or may be implied by awrt 0.964</p> <p><b>Use of Normal in (d)</b>  Can get M0 M1 A0.- for M1 they must write <math>1 - P(X \leq 4)</math> or get awrt 0.187</p>	

Question Number	Scheme			Marks
Q3	<p style="text-align: center;">Method 1</p> $P(X > 6) = \frac{1}{6}$ $P(X < 4) = \frac{1}{2}$ $\text{total} = \frac{1}{6} + \frac{1}{2} = \frac{2}{3}$	<p style="text-align: center;">Method 2</p> $P(4 < X < 6) = \frac{1}{3}$ $1 - \frac{1}{3} = \frac{2}{3}$	<p style="text-align: center;">Method 3</p> $P(X > 6) = \frac{1}{6}$ $Y \sim U[3,9] \quad P(Y > 6) = \frac{1}{2}$ $\text{total} = \frac{1}{6} + \frac{1}{2} = \frac{2}{3}$	<p>B1 M1</p> <p>A1</p> <p>M1dep B A1</p> <p style="text-align: right;">(5)</p> <p style="text-align: right;">[5]</p>
<p>Notes</p> <p><b>Methods 1 and 2</b></p> <p><b>B1</b> for 6 and 4 (allow if seen on a diagram on <math>x</math>-axis)</p> <p><b>M1</b> for <math>P(X &gt; 6)</math> or <math>P(6 &lt; X &lt; 7)</math>; or <math>P(X &lt; 4)</math> or <math>P(1 &lt; X &lt; 4)</math>; or <math>P(4 &lt; X &lt; 6)</math></p> <p>Allow <math>\leq</math> and <math>\geq</math> signs</p> <p><b>A1</b> <math>\frac{1}{6}</math>; or <math>\frac{1}{2}</math>; <math>\frac{1}{3}</math> must match the probability statement</p> <p><b>M1</b> for adding their “<math>P(X &gt; 6)</math>” and their “<math>P(X &lt; 4)</math>” or 1 - their “<math>P(4 &lt; X &lt; 6)</math>” dep on getting first B mark</p> <p><b>A1</b> cao <math>\frac{2}{3}</math></p> <p><b>Method 3 <math>Y \sim U[3, 9]</math></b></p> <p><b>B1</b> for 6 with <math>U[1,7]</math> and 6 with <math>U[3,9]</math></p> <p><b>M1</b> for <math>P(X &gt; 6)</math> or <math>P(6 &lt; X &lt; 7)</math> or <math>P(6 &lt; Y &lt; 9)</math></p> <p><b>A1</b> <math>\frac{1}{6}</math>; or <math>\frac{1}{2}</math>; must match the probability statement</p> <p><b>M1</b> for adding their “<math>P(X &gt; 6)</math>” and their “<math>P(Y &gt; 6)</math>” dep on getting first B mark</p> <p><b>A1</b> cao <math>\frac{2}{3}</math></p>				



Question Number	Scheme	Marks
Q4 (a)	$\frac{4}{9}(m^2 + 2m - 3) = 0.5$ $m^2 + 2m - 4.125 = 0$ $m = \frac{-2 \pm \sqrt{4 + 16.5}}{2}$ $m = 1.26, -3.264$ (median =) 1.26	M1 M1 A1 (3)
(b)	Differentiating $\frac{d\left(\frac{4}{9}(x^2 + 2x - 3)\right)}{dx} = \frac{4}{9}(2x + 2)$	M1 A1
(c)	$f(x) = \begin{cases} \frac{8}{9}(x+1) & 1 \leq x \leq 1.5 \\ 0 & \text{otherwise} \end{cases}$	B1ft (3)
(c)	$P(X \geq 1.2) = 1 - F(1.2)$ $= 1 - 0.3733$ $= \frac{47}{75}, 0.6267$ 0.627	M1 awrt A1 (2)
(d)	$(0.6267)^4 = 0.154$	awrt 0.154 or 0.155 M1 A1 (2)
<b>[10]</b>		
<u>Notes</u>		
(a)	<b>M1</b> putting $F(x) = 0.5$ <b>M1</b> using correct quadratic formula. If use calc need to get 1.26 (384... ) <b>A1</b> cao 1.26 must reject the other root.	
If they use Trial and improvement they have to get the correct answer to gain the second M mark.		
(b)	<b>M1</b> attempt to differentiate. At least one $x^n \rightarrow x^{n-1}$ <b>A1</b> correct differentiation <b>B1</b> must have both parts- follow through their $F'(x)$ Condone <	
(c)	<b>M1</b> finding/writing $1 - F(1.2)$ may use/write $\int_{1.2}^{1.5} \frac{8}{9}(x+1)dx$ or $1 - \int_1^{1.2} \frac{8}{9}(x+1)dx$ or $\int_{1.2}^{1.5}$ "their $f(x)$ " $dx$ . Condone missing $dx$	
(d)	<b>A1</b> awrt 0.627	
(d)	<b>M1</b> (c) <sup>4</sup> If expressions are not given you need to check the calculation is correct to 2sf. <b>A1</b> awrt 0.154 or 0.155	

Question Number	Scheme	Marks
Q5	<p>(a) Connecting occurs at random/independently, singly or at a constant rate</p> <p>(b) <math>P_0(8)</math></p> <p>(i) <math>P(X = 0) = 0.0003</math></p> <p>(ii) <math>P(X \geq 4) = 1 - P(X \leq 3)</math>  <math>= 1 - 0.0424</math>  <math>= 0.9576</math></p> <p>(c) <math>H_0: \lambda = 4</math> (48) <math>H_1: \lambda &gt; 4</math> (48)  <math>N(48, 48)</math>  Method 1  <math display="block">P(X \geq 59.5) = P\left(Z \geq \frac{59.5 - 48}{\sqrt{48}}\right)</math> <math display="block">= P(Z \geq 1.66)</math> <math display="block">= 1 - 0.9515</math> <math display="block">= 0.0485</math>   <math>0.0485 &lt; 0.05</math>  Reject <math>H_0</math>. Significant. 60 lies in the Critical region  The number of failed connections at the first attempt has increased.</p> <div style="display: inline-block; vertical-align: middle; border-left: 1px solid black; padding-left: 10px; margin-left: 20px;"> Method 2  <math display="block">\frac{x - 0.5 - 48}{\sqrt{48}} = 1.6449</math>    <math>x = 59.9</math> </div>	B1 (1) B1 M1A1 M1 A1 (5)  B1 M1 A1  M1 M1 A1  A1  M1 A1 ft (9) [15]
	<p><b>Notes</b></p> <p>(a) <b>B1</b> Any one of randomly/independently/singly/constant rate. Must have context of connection/logging on/fail</p> <p>(b) <b>B1</b> Writing or using <math>P_0(8)</math> in (i) or (ii)</p> <p>(i) <b>M1</b> for writing or finding <math>P(X = 0)</math>  <b>A1</b> awrt 0.0003</p> <p>(ii) <b>M1</b> for writing or finding <math>1 - P(X \leq 3)</math>  <b>A1</b> awrt 0.958</p> <p>(c) <b>B1</b> both hypotheses correct. Must use <math>\lambda</math> or <math>\mu</math>  <b>M1</b> identifying normal  <b>A1</b> using or seeing mean and variance of 48  These first two marks may be given if the following are seen in the standardisation formula : 48 and <math>\sqrt{48}</math> or awrt 6.93  <b>M1</b> for attempting a continuity correction (Method 1: <math>60 \pm 0.5</math> / Method 2: <math>x \pm 0.5</math>)  <b>M1</b> for standardising using their mean and their standard deviation and using either Method 1 [<math>59.5, 60</math> or <math>60.5</math>. accept <math>\pm z</math>.] Method 2 [<math>(x \pm 0.5)</math> and equal to a <math>\pm z</math> value]  <b>A1</b> correct z value awrt <math>\pm 1.66</math> or <math>\pm \frac{59.5 - 48}{\sqrt{48}}</math>, or <math>\frac{x - 0.5 - 48}{\sqrt{48}} = 1.6449</math>  <b>A1</b> awrt 3 sig fig in range 0.0484 – 0.0485, awrt 59.9  <b>M1</b> for “reject <math>H_0</math>” or “significant” maybe implied by “correct contextual comment”  If one tail hypotheses given follow through “their prob” and 0.05, <math>p &lt; 0.5</math>  If two tail hypotheses given follow through “their prob” with 0.025, <math>p &lt; 0.5</math>  If one tail hypotheses given follow through “their prob” and 0.95, <math>p &gt; 0.5</math>  If two tail hypotheses given follow through “their prob” with 0.975, <math>p &gt; 0.5</math>  If no <math>H_1</math> given they get M0  <b>A1</b> ft correct contextual statement followed through from their prob and <math>H_1</math>. need the words <u>number of failed connections/log ons has increased</u> o.e.  Allow “there are more failed connections”  <b>NB</b> A correct contextual statement <b>alone</b> followed through from their prob and <math>H_1</math> gets M1 A1</p>	

Question Number	Scheme	Marks
Q6	<p>(a) 2 outcomes/faulty or not faulty/success or fail A constant probability Independence Fixed number of trials (fixed <math>n</math>)</p> <p>(b) <math>X \sim B(50,0.25)</math> <math>P(X \leq 6) = 0.0194</math> <math>P(X \leq 7) = 0.0453</math> <math>P(X \geq 18) = 0.0551</math> <math>P(X \geq 19) = 0.0287</math>  CR <math>X \leq 6</math> and <math>X \geq 19</math></p> <p>(c) <math>0.0194 + 0.0287 = 0.0481</math></p> <p>(d) 8(It) is not in the Critical region or 8(It) is not significant or <math>0.0916 &gt; 0.025</math>; There is evidence that the probability of a faulty bolt is 0.25 or the company's claim is correct.</p> <p>(e) <math>H_0 : p = 0.25</math> <math>H_1 : p &lt; 0.25</math> <math>P(X \leq 5) = 0.0070</math> or CR <math>X \leq 5</math> <math>0.007 &lt; 0.01</math>, 5 is in the critical region, reject <math>H_0</math>, significant. There is evidence that the probability of faulty bolts has decreased</p>	<p>B1 B1  (2)</p> <p>M1     A1 A1 (3)</p> <p>M1A1 (2)</p> <p>M1; A1ft (2)</p> <p>B1B1 M1A1  M1 A1ft (6) [15]</p>
	<p>Notes</p> <p>(a) <b>B1 B1</b> one mark for each of any of the four statements. Give first B1 if only one correct statement given. No context needed.</p> <p>(b) <b>M1</b> for writing or using <math>B(50,0.25)</math> also may be implied by both CR being correct. Condone use of P in critical region for the method mark. <b>A1</b> <math>(X) \leq 6</math> o.e. <math>[0,6]</math> DO NOT accept <math>P(X \leq 6)</math> <b>A1</b> <math>(X) \geq 19</math> o.e. <math>[19,50]</math> DO NOT accept <math>P(X \geq 19)</math></p> <p>(c) <b>M1</b> Adding two probabilities for two tails. Both probabilities must be less than 0.5 <b>A1</b> awrt 0.0481</p> <p>(d) <b>M1</b> one of the given statements followed through from their CR. <b>A1</b> contextual comment followed through from their CR. NB A correct contextual comment <b>alone</b> followed through from their CR will get M1 A1</p> <p>(e) <b>B1</b> for <math>H_0</math> must use <math>p</math> or <math>\pi</math> (<math>\pi</math>) <b>B1</b> for <math>H_1</math> must use <math>p</math> or <math>\pi</math> (<math>\pi</math>) <b>M1</b> for finding or writing <math>P(X \leq 5)</math> or attempting to find a critical region or a correct critical region <b>A1</b> awrt 0.007/CR <math>X \leq 5</math> <b>M1</b> correct statement using their Probability and 0.01 if one tail test or a correct statement using their Probability and 0.005 if two tail test. The 0.01 or 0.005 needn't be explicitly seen but implied by correct statement compatible with their <math>H_1</math>. If no <math>H_1</math> given M0 <b>A1</b> correct contextual statement follow through from their prob and <math>H_1</math>. Need faulty bolts and decreased. NB A correct contextual statement <b>alone</b> followed through from their prob and <math>H_1</math> get M1 A1</p>	



Question Number	Scheme	Marks
Q7 (ai)	$f(y) \geq 0$ or $f(3) \geq 0$ $ky(a-y) \geq 0$ or $3k(a-3) \geq 0$ or $(a-y) \geq 0$ or $(a-3) \geq 0$ $a \geq 3$	M1  A1 cso
	(ii) $\int_0^3 k(ay - y^2) dy = 1$ $\left[ k \left( \frac{ay^2}{2} - \frac{y^3}{3} \right) \right]_0^3 = 1$ $k \left( \frac{9a}{2} - 9 \right) = 1$ $k \left[ \frac{9a-18}{2} \right] = 1$ $k = \frac{2}{9(a-2)}$ *	integration  M1  answer correct A1  answer = 1 M1  A1 cso 6)
(b)	$\int_0^3 k(ay^2 - y^3) dy = 1.75$ $\left[ k \left( \frac{ay^3}{3} - \frac{y^4}{4} \right) \right]_0^3 = 1.75$ $k \left( 9a - \frac{81}{4} \right) = 1.75$ $2 \left( 9a - \frac{81}{4} \right) = 15.75(a-2)$ $2.25a = -31.5 + \frac{81}{2}$ $a = 4$ * $k = \frac{1}{9}$	Int $\int xf(x)$ M1 Correct integration A1 $\int xf(x) = 1.75$ and limits 0,3 M1dep  subst $k$ M1dep  A1cso B1 (6)

Question Number	Scheme	Marks
(c)		B1 B1 (2)
(d)	mode = 2	B1 (1)
		<b>[15]</b>
(a) (i)	<p>Notes</p> <p><b>M1</b> for putting <math>f(y) \geq 0</math> or <math>f(3) \geq 0</math> or <math>ky(a - y) \geq 0</math> or <math>3k(a - 3) \geq 0</math> or <math>(a - y) \geq 0</math> or <math>(a - 3) \geq 0</math> or state in words the probability can not be negative o.e.</p> <p><b>A1</b> need one of <math>ky(a - y) \geq 0</math> or <math>3k(a - 3) \geq 0</math> or <math>(a - y) \geq 0</math> or <math>(a - 3) \geq 0</math> <b>and</b> <math>a \geq 3</math></p>	
(ii)	<p><b>M1</b> attempting to integrate (at least one <math>y^n \rightarrow y^{n+1}</math>) (ignore limits)</p> <p><b>A1 Correct integration.</b> Limits not needed. And equals 1 not needed.</p> <p><b>M1</b> dependent on the previous M being awarded. Putting equal to 1 and have the correct limits. Limits do not need to be substituted.</p> <p><b>A1</b> cso</p>	
(b)	<p><b>M1</b> for attempting to find <math>\int yf(y) dy</math> (at least one <math>y^n \rightarrow y^{n+1}</math>) (ignore limits)</p> <p><b>A1</b> correct Integration</p> <p><b>M1</b> <math>\int yf(y) = 1.75</math> and limits 0,3 dependent on previous M being awarded</p> <p><b>M1</b> subst in for <math>k</math>. dependent on previous M being awarded</p> <p><b>A1</b> cso 4</p> <p><b>B1</b> cao 1/9</p>	
(c)	<p><b>B1</b> correct shape. No straight lines. No need for patios.</p> <p><b>B1</b> completely correct graph. Needs to go through origin and the curve ends at 3.</p> <p><u>Special case:</u> If draw full parabola from 0 to 4 get B1 B0 Allow full marks if the portion between <math>x = 3</math> and <math>x = 4</math> is dotted and the rest of the curve solid.</p>	
		
(d)	<b>B1</b> cao 2	



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA024768 Summer 2010

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH