

GCE

Mathematics (MEI)

Advanced GCE


Unit **4767**: Statistics 2

Mark Scheme for June 2011

4767

Mark Scheme

June 2011

1 (i)		<p>G1 for axes</p> <p>G1 For values of x</p> <p>G1 for values of y</p>	3	<p>Condone axes drawn either way.</p> <p>Axes should show some indication of scale. If not then Max G1 if points ‘visibly correct’.</p> <p>If axes are scaled and only one point is incorrectly plotted, allow max G2/3.</p>
1 (ii)	<p>$\bar{x} = 60, \bar{y} = 4.26$</p> $b = \frac{S_{xy}}{S_{xx}} = \frac{1803 - 300 \times 21.3/5}{27000 - 300^2/5} = \frac{525}{9000} = 0.0583$ <p>OR $b = \frac{1803/5 - 60 \times 4.26}{27000/5 - 60^2} = \frac{105}{1800} = 0.0583$</p> <p>hence least squares regression line is:</p> $y - \bar{y} = b(x - \bar{x})$ $\Rightarrow y - 4.26 = 0.0583(x - 60)$ $\Rightarrow y = 0.0583x + 0.76$	<p>B1 for \bar{x} and \bar{y} used appropriately (SOI)</p> <p>M1 for attempt at gradient (b)</p> <p>A1 for 0.0583 cao</p> <p>M1 for equation of line</p> <p>A1 FT for complete equation</p>	5	<p>B1 for means can be implied by a correct value of b using either method. Allow $\bar{y} = 4.3$</p> <p>Attempt should be correct – e.g. evidence of either of the two suggested methods should be seen.</p> <p>Allow 0.058 Condone $0.058\dot{3}$ and $\frac{7}{120}$</p> <p>Dependent on first M1. Values must be substituted to earn M1. Condone use of their b for FT provided $b > 0$. Final equation must be simplified.</p> <p>$b = 0.058$ leads to $y = 0.058x + 0.78$</p>
1 (iii)	<p>Regression line plotted on graph</p> <p>The fit is good</p>	<p>G1</p> <p>G1</p> <p>E1 for good fit</p>	3	<p>Line must pass through their (\bar{x}, \bar{y}) and y-intercept.</p> <p>E0 for notably inaccurate graphs/lines</p>

4767

Mark Scheme

June 2011

1 (iv)	$x = 30 \Rightarrow$ predicted $y = 0.0583 \times 30 + 0.76 = 2.509$ Residual = $2.5 - 2.509 = -0.009$	B1 for prediction M1 for subtraction A1 FT	3	Using their equation Subtraction can be 'either way' but for the final mark the sign of the residual must be correct. FT sensible equations only – e.g. no FT for $y = 0.071x$ leading to +0.37. [$c = 0.78$ leads to a residual of -0.02]
1 (v)	(A) For $x = 45$, $y = 0.0583 \times 45 + 0.76 = 3.4$ (B) For $x = 150$, $y = 0.0583 \times 150 + 0.76 = 9.5$	M1 for at least one prediction attempted A1 for both answers (FT their equation provided their $b > 0$)	2	Prediction obtained from their equation.
1 (vi)	This is well below the predicted valuesuggesting that the model breaks down for larger values of x .	E1 for well below E1 extrapolation	2	Some indication that the value (8.7) is significantly below what is expected (9.5) is required for the first E1. Simply pointing out that it is 'below' is not sufficient. The second E1 is available for a suitable comment relating to the model being suitable only for values within the domain of the given points. Allow other sensible comments for either E1. E.g. The data might be better modelled by a curve', 'there may be other factors affecting yield',
			18	

4767

Mark Scheme


June 2011

2 (i)	Independently means that the arrival time of each car is unrelated to the arrival time of any other car. Randomly means that the arrival times of cars are not predictable. At a uniform average rate means that the average rate of car arrivals does not vary over time.	E1 E1 E1	3	NOTE Each answer must be ‘in context’ and ‘clear’ Allow sensible alternative wording. SC1 For ALL answers not in context but otherwise correct.
2 (ii)	$P(\text{At most 1 car}) = e^{-0.62} \frac{0.62^0}{0!} + e^{-0.62} \frac{0.62^1}{1!}$ $= 0.5379\dots + 0.3335\dots = 0.871$	M1 for either M1 for sum of both A1 CAO	3	$1.62e^{-0.62}$ Allow 0.8715 not 0.872 or 0.8714 Allow 0.87 without wrong working seen
2 (iii)	New $\lambda = 10 \times 0.62 = 6.2$ $P(\text{more than 5 in 10 mins}) = 1 - 0.4141 = 0.5859$	B1 for mean (SOI) M1 for probability A1 CAO	3	Use of $1 - P(X \leq 5)$ with any λ Allow 0.586
2 (iv)	Poisson with mean 37.2	B1 for Poisson B1 for mean 37.2	2	Dependent on first B1 Condone $P(37.2, 37.2)$
2 (v)	Use Normal approx with $\mu = \sigma^2 = \lambda = 37.2$ $P(X \geq 40) = P\left(Z > \frac{39.5 - 37.2}{\sqrt{37.2}}\right)$ $= P(Z > 0.377) = 1 - \Phi(0.377) = 1 - 0.6469$ $= 0.3531$	B1 for Normal (SOI) B1 for parameters B1 for 39.5 M1 for correct use of Normal approximation using correct tail A1 cao	5	Allow 0.353
			16	

4767

Mark Scheme

June 2011

3(iv)(B)	 <p style="text-align: center;">Cox's Braeburns</p>	<p>G1 for shape G1 for means, shown explicitly or by scale</p> <p>G1 for lower max height for Braeburns G1 for greater width (variance) for Braeburns</p>	4	<p>Ignore labelling of vertical axis.</p> <p>Two intersecting, adjacent Normal curves Means at 185 and 210.5</p>
		TOTAL	20	
4(a)(i)	<p>H_0: no association between amount spent and sex H_1: some association between amount spent and sex</p>	B1 for both	1	Hypotheses must be the right way round, in context and must not mention 'correlation'.
4(a)(ii)	<p>Expected frequency = $62 \times 102 \div 200 = 31.62$</p> <p>Contribution = $(34 - 31.62)^2 / 31.62 = 0.1791$</p>	<p>B1</p> <p>M1 A1 for valid attempt at $(O-E)^2/E$</p> <p>NB Answer given</p>	3	Do not allow 31.6

4767

Mark Scheme

June 2011

4(a)(iii)	<p>Refer to χ_4^2 Critical value at 5% level = 9.488 $3.205 < 9.488$ Result is not significant</p> <p>There is insufficient evidence to suggest any association between amount spent and sex.</p>	<p>B1 for 4 deg of freedom B1 CAO for cv M1 A1 for not significant</p> <p>E1</p>	5	<p>Allow $p = 0.524$ $0.524 > 0.05$ Conclusion must be stated to earn A1 here. Allow 'do not reject H_0' and condone 'accept H_0' or 'reject H_1'. FT if cv consistent with their d.o.f. Dependent on previous A1 and final comment must be in context and not mention correlation. SC1 for correct final conclusion where previous A1 omitted but M1 awarded.</p>
4 (b)	<p>$H_0: \mu = 400; H_1: \mu < 400$ Where μ denotes the population mean (weight of the loaves).</p> <p>$\bar{x} = 396.5$</p> <p>Test statistic = $\frac{396.5 - 400}{5.7/\sqrt{6}} = \frac{-3.5}{2.327} = -1.504$</p> <p>5% level 1 tailed critical value of $z = -1.645$</p> <p>$-1.504 > -1.645$ so not significant.</p> <p>There is insufficient evidence to reject H_0</p> <p>There is insufficient evidence to suggest that the true mean weight of the loaves is lower than the minimum specified value of 400 grams.</p>	<p>B1 for H_0 B1 for H_1 B1 for definition of μ</p> <p>B1 for sample mean</p> <p>M1 must include $\sqrt{6}$ A1FT their sample mean</p> <p>B1 for ± 1.645</p> <p>M1 for sensible comparison leading to a conclusion</p> <p>A1 for conclusion in context</p>	9	<p>Hypotheses in words must refer to population mean.</p> <p>Condone numerator reversed for M1 but award A1 only if test statistic of 1.504 is compared with a positive z-value.</p> <p>Dependent on previous M1</p> <p>FT their sample mean only if hypotheses are correct.</p>
		TOTAL	18	

4767

Mark Scheme

June 2011

Additional notes re Q1 parts (ii), (iv) and (v)

Part (ii) 'x on y' max B1

Part (iv) $x = 16.9y - 12.02$ leads to a prediction of $x = 30.23$ and a residual of -0.23 B1 M1 A1 available.

Part (v) 'x on y' not appropriate here so award 0 if 'x on y' used.

Additional notes re Q2 parts (i) & (v)

Part (i)

Independent – Allow 'not linked to' or 'no association' or 'unrelated' 'not affected by', 'not connected to', 'not influenced by'

DO NOT ACCEPT 'not together' or 'not dependent'

Random – Allow 'not predictable' or 'no pattern' or 'could happen at any time' or 'not specific time'

Uniform average rate – Allow 'average (rate) is constant over time' DO NOT ACCEPT 'average constant' or 'average rate and uniform' – be generous over defining 'average' and/or 'rate'.

Part (v) If Binomial distribution stated in part (iv), allow B1 B0 B1 M0 A0 max

Additional notes re Q3 part (iii) where $p > 0.1$ (iii) *B* – as scheme unless a Normal approximation is more suitable ($p > 0.1$). If so, award B1 B1 for Normal and correct parameters. The remaining marks are dependent on both these B1 marks being awarded. M1 for the correct continuity correction ($P(X < 1.5)$) and depM1 for the correct tail but award A0.(iii) *C* – ' n is large and p is not too small' or ' $np > 10$ '

4767

Mark Scheme

June 2011

Additional notes re Q4(b) σ estimated

sample mean, 7.079... used in place of 5.7, the given value of the population mean, leads to a test statistic of -1.212... This gets M1A0 & the remaining marks are still available.

Critical Value Method

$400 - 1.645 \times 5.7 \div \sqrt{6} \dots$ gets M1B1 ... = 396.17... gets A1

$400 + 1.645 \times 5.7 \div \sqrt{6}$ gets M1B1A0.

$396.5 > 396.2$ gets M1 for sensible comparison (and B1 for 396.5)

A1 still available for correct conclusion in words & context

90% Confidence Interval Method

CI centred on 396.5 (gets B1 for 396.5)

+ or - $1.645 \times 5.7 \div \sqrt{6}$ gets M1 B1

= (392.67, 400.33) A1

contains 400 gets M1

A1 still available for correct conclusion in words & context

Probability Method

Finding $P(\text{sample mean} < 396.5) = 0.0663$ gets M1 A1 (and B1 for 396.5)

$0.0663 > 0.05$ gets M1 for a sensible comparison if a conclusion is made and also gets the B1 for 0.0663 (to replace the B1 for $cv = 1.645$).

A1 still available for correct conclusion in words & context.

Condone $P(\text{sample mean} > 396.5) = 0.9337$ for M1 and B1 for 0.9337 but only allow A1 if later compared with 0.95 at which point the final M1 and A1 are still available

Two-tailed test

Max B1 B0 B1 B1 M1 A1 B1 (for $cv = -1.96$) M1 A0