



#### **Exercise 4A**

**1** a Sketching the function:



There are negative values for f(x) when  $-1 \le x < 0$ , so this is not a probability density function.


**b** Sketching the function:



There is no negative value of f(x)Area under  $f(x) = \int_{1}^{3} x^{2} dx = \left[\frac{x^{3}}{3}\right]_{1}^{3} = \frac{27}{3} - \frac{1}{3} = \frac{26}{3}$ 

Area is not equal to 1, therefore this is not a valid probability density function.

**c** When f(x) = 0,  $x = 2^{\frac{1}{3}} = 1.26$  (2 d.p.). So for  $-1 \le x < 1.26$ , f(x) < 0. As there are negative values for f(x), this is not a probability density function. Alternatively, reach this result by sketching the function:



2 The area under the curve must equal 1, so:

$$\int_{-4}^{-2} k(x^{2} - 1) dx = 1$$

$$k \left[ \frac{x^{3}}{3} - x \right]_{-4}^{-2} = 1$$


$$k \left( \left( -\frac{8}{3} + 2 \right) - \left( -\frac{64}{3} + 4 \right) \right) = 1$$

$$k \left( \frac{56}{3} - 2 \right) = 1$$


$$\frac{50}{3} k = 1$$

$$k = \frac{3}{50}$$

**3** a For the non-zero parts of the function, its graph is a straight line running from (2, 0) to (6, 0.5).



**b** For the non-zero parts of the function, its graph is a straight line running from  $\left(1, \frac{8}{15}\right)$  to  $\left(4, \frac{2}{15}\right)$ .



**4 a** The area under the curve must equal 1, so:

 $\int_{1}^{3} kx \, dx = 1$  $\left[\frac{kx^{2}}{2}\right]_{1}^{3} = 1$  $\frac{9k}{2} - \frac{k}{2} = 1$ 4k = 1 $k = \frac{1}{4}$ 

Pearson

#### **INTERNATIONAL A LEVEL**

#### Statistics 2

### Solution Bank




4 **b** 
$$\int_{0}^{3} kx^{2} dx = 1$$
$$\left[\frac{kx^{3}}{3}\right]_{0}^{3} = 1$$
$$\frac{27k}{3} = 1$$
$$9 k = 1$$
$$k = \frac{1}{9}$$
**c** 
$$\int_{0}^{2} k(1 + 1) k = \frac{1}{9}$$

$$\int_{-1}^{2} k(1+x^2) dx = 1$$
$$k \left[ x + \frac{x^3}{3} \right]_{-1}^{2} = 1$$
$$k \left( \left( 2 + \frac{8}{3} \right) - \left( -1 - \frac{1}{3} \right) \right) = 1$$
$$k \left( 3 + \frac{9}{3} \right) = 1$$
$$6k = 1$$
$$k = \frac{1}{6}$$

**5** a The area under the curve must equal 1, so:

$$\int_0^2 k(4-x)dx = 1$$
$$k \left[ 4x - \frac{x^2}{2} \right]_0^2 = 1$$
$$k(8-2) = 1$$
$$6k = 1$$
$$k = \frac{1}{6}$$

**b** For the non-zero parts of the function, its graph is a straight line running from  $\left(0, \frac{2}{3}\right)$  to  $\left(2, \frac{1}{3}\right)$ .



#### **INTERNATIONAL A LEVEL**

### **Statistics 2** Solution Bank



5 c 
$$P(X > 1) = \int_{1}^{2} \frac{1}{6} (4 - x) dx = \left[\frac{2}{3}x - \frac{1}{12}x^{2}\right]_{1}^{2}$$
  
=  $\left(\frac{4}{3} - \frac{1}{3}\right) - \left(\frac{2}{3} - \frac{1}{12}\right) = \frac{1}{3} + \frac{1}{12} = \frac{5}{12}$ 

**6 a** The area under the curve must equal 1, so:

$$\int_{0}^{2} kx^{2}(2-x) dx = 1$$

$$k \left[ \frac{2x^{3}}{3} - \frac{x^{4}}{4} \right]_{0}^{2} = 1$$

$$k \left( \frac{16}{3} - \frac{16}{4} \right) = 1$$

$$\frac{16k}{12} = 1$$

$$k = \frac{3}{4} = 0.75$$

**b** 
$$P(0 < X < 1) = \int_0^1 \frac{3}{4} x^2 (2 - x) dx = \left[\frac{1}{2}x^3 - \frac{3}{16}x^4\right]_0^1 = \frac{5}{16}$$

7 a The area under the curve must equal 1, so:


$$\int_{1}^{4} kx^{3} dx = 1$$
$$\left[\frac{kx^{4}}{4}\right]_{1}^{4} = 1$$
$$\frac{256k}{4} - \frac{k}{4} = 1$$
$$\frac{255k}{4} = 1$$
$$k = \frac{4}{255}$$

- **b**  $\int_{1}^{2} \frac{4}{255} x^{3} dx = \left[\frac{1}{255} x^{4}\right]_{1}^{2} = \frac{15}{255} = \frac{1}{17} = 0.0588 \ (4 \ \text{d.p.})$
- 8 a The area under the curve must equal 1, so:  $\int_{1}^{2} 1 dx = \int_{1}^{3} 1 dx$

$$\int_{0}^{2} k \, dx + \int_{2}^{3} k(2x-3) \, dx = 1$$
$$[kx]_{0}^{2} + [kx^{2} - 3kx]_{2}^{3} = 1$$
$$2k + [(9k - 9k) - (4k - 6k)] = 1$$
$$2k + 2k = 1$$
$$k = \frac{1}{4} = 0.25$$



8 b For the non-zero parts of the function, its graph is a horizontal line running from (0, 0.25) to (2, 0.25) and then a straight line from (2, 0.25) to (3, 0.75).




c  $P(X < 1) = \int_0^1 0.25 dx = [0.25x]_0^1 = 0.25$  $P(Y < 1) = \int_{-2}^1 \frac{3}{16} y^2 dy = \left[\frac{1}{16}y^3\right]_{-2}^1 = \frac{1}{16} - \left(-\frac{8}{16}\right) = \frac{9}{16}$ 

As X and Y are independent:

$$P(X < 1 \cap Y < 1) = P(X < 1) \times P(Y < 1) = \frac{1}{4} \times \frac{9}{16} = \frac{9}{64}$$


9 a 
$$P(X < 0.5) = \int_0^{0.5} \frac{1}{60} (x+1) dx = \frac{1}{60} \Big[ 0.5x^2 + x \Big]_0^{0.5}$$
  
=  $\frac{1}{60} \Big( \frac{1}{8} + \frac{1}{2} \Big) = \frac{1}{60} \times \frac{5}{8} = \frac{1}{96} = 0.0104 \ (4 \ d.p.)$ 

**b** For the non-zero parts of the function, its graph is a straight line running from  $\left(0, \frac{1}{60}\right)$  to





9 c By definition, every visitor would spend some time (however short) on the site, but the probability of spending a long time on the site would be very low but not become zero as x gets larger. So in reality the probability density might look like this:



11 a The area under the curve must equal 1, so:

$$\int_{1}^{5} \frac{k}{x} dx = 1$$
$$\left[k \ln x\right]_{1}^{5} = 1$$
$$k \ln 5 = 1$$
$$k = \frac{1}{\ln 5}$$

- **b**  $P(2 < X < 4) = \frac{1}{\ln 5} \int_{2}^{4} \frac{1}{x} dx = \frac{1}{\ln 5} [\ln x]_{2}^{4}$  $= \frac{1}{\ln 5} (\ln 4 - \ln 2) = \frac{\ln 2}{\ln 5}$
- 12 a The area under the curve must equal 1, so:

$$\int_{-1}^{4} \frac{k}{x+2} dx = 1$$

$$\begin{bmatrix} k \ln(x+2) \end{bmatrix}_{-1}^{4} = 1$$

$$k \ln 6 = 1$$

$$k = \frac{1}{\ln 6}$$

$$\mathbf{b} \quad P(1 < X < 3) = \frac{1}{\ln 6} \int_{1}^{3} \frac{1}{x+2} dx = \frac{1}{\ln 6} [\ln(x+2)]_{1}^{3}$$

$$= \frac{\ln 6^{31} x + 2}{\ln 6} (\ln 5 - \ln 3) = \frac{\ln 1.666...}{\ln 6} = 0.285 (3 \text{ d.p.})$$

#### **INTERNATIONAL A LEVEL**


## **Statistics 2** Solution Bank

**13 a** The area under the curve must equal 1, so:

$$\int_0^1 k \sin \pi x \, dx = 1$$
$$\left[ -\frac{k}{\pi} \cos \pi x \right]_0^1 = 1$$
$$\frac{k}{\pi} (1 - (-1)) = 1$$
$$k = \frac{\pi}{2}$$

**b** For the non-zero parts of the function, its graph is a sine curve of amplitude  $\frac{\pi}{2}$  running from (0, 0) to (1, 0).

Pearson





#### Challenge

**a** The area under the curve must equal 1, so:

$$\int_{1}^{\infty} \frac{k}{t^{3}} dt = 1 \Longrightarrow k \int_{1}^{\infty} \frac{1}{t^{3}} dt = 1 \Longrightarrow k \times \frac{1}{2} = 1 \Longrightarrow k = 2$$

**b** i 
$$P(0 < T < 3) = \int_{1}^{3} \frac{2}{t^{3}} dt = \left[-t^{-2}\right]_{1}^{3} = 1 - \frac{1}{9} = \frac{8}{9}$$

ii 
$$P(T > 20) = \int_{20}^{\infty} \frac{2}{t^3} dt = \left[-t^{-2}\right]_{20}^{\infty} = \frac{1}{20^2} = \frac{1}{400}$$

c 
$$P(p < T < 2p) = \int_{p}^{2p} \frac{2}{t^{3}} dt = \left[-t^{-2}\right]_{p}^{2p} = \frac{1}{p^{2}} - \frac{1}{4p^{2}}$$
  
So  $\frac{1}{p^{2}} - \frac{1}{4p^{2}} = 0.12$   
 $\Rightarrow \frac{3}{4p^{2}} = 0.12$   
 $\Rightarrow p^{2} = \frac{3}{4 \times 0.12} = \frac{1}{4 \times 0.04} = \frac{1}{0.16} = 6.25$   
 $\Rightarrow p = 2.5$