

Leave blank

3. A doctor is interested in the relationship between a person's Body Mass Index (BMI) and their level of fitness. She believes that a lower BMI leads to a greater level of fitness. She randomly selects 10 female 18 year-olds and calculates each individual's BMI. The females then run a race and the doctor records their finishing positions. The results are shown in the table.

Individual	A	B	C	D	E	F	G	H	I	J
BMI	17.4	21.4	18.9	24.4	19.4	20.1	22.6	18.4	25.8	28.1
Finishing position	3	5	1	9	6	4	10	2	7	8

- (a) Calculate Spearman's rank correlation coefficient for these data. **(5)**
- (b) Stating your hypotheses clearly and using a one tailed test with a 5% level of significance, interpret your rank correlation coefficient. **(5)**
- (c) Give a reason to support the use of the rank correlation coefficient rather than the product moment correlation coefficient with these data. **(1)**



Leave blank

- 4. A sample of size 8 is to be taken from a population that is normally distributed with mean 55 and standard deviation 3. Find the probability that the sample mean will be greater than 57.

(5)



Leave blank

5. The number of goals scored by a football team is recorded for 100 games. The results are summarised in Table 1 below.

Number of goals	Frequency
0	40
1	33
2	14
3	8
4	5

Table 1

- (a) Calculate the mean number of goals scored per game. (2)

The manager claimed that the number of goals scored per match follows a Poisson distribution. He used the answer in part (a) to calculate the expected frequencies given in Table 2.

Number of goals	Expected Frequency
0	34.994
1	r
2	s
3	6.752
≥ 4	2.221

Table 2

- (b) Find the value of r and the value of s giving your answers to 3 decimal places. (3)
- (c) Stating your hypotheses clearly, use a 5% level of significance to test the manager's claim. (7)



Leave
blank

Question 5 continued

A series of horizontal lines for writing the answer to Question 5.



Leave blank

6. The lengths of a random sample of 120 limpets taken from the upper shore of a beach had a mean of 4.97 cm and a standard deviation of 0.42 cm. The lengths of a second random sample of 150 limpets taken from the lower shore of the same beach had a mean of 5.05 cm and a standard deviation of 0.67 cm.

(a) Test, using a 5% level of significance, whether or not the mean length of limpets from the upper shore is less than the mean length of limpets from the lower shore. State your hypotheses clearly.

(8)

(b) State two assumptions you made in carrying out the test in part (a).

(2)



Leave blank

Question 7 continued

Lined area for writing the answer to Question 7.

(Total 11 marks)

Q7

--	--



