

Chapter Review 4

1 By the central limit theorem $\bar{X} \approx \sim N\left(5, \frac{1}{100}\right)$, i.e. $\bar{X} \approx \sim N(5, 0.01)$

$$P(\bar{X} > 5.2) = 1 - P(\bar{X} < 5.2) \approx 1 - 0.9772 = 0.0228 \text{ (4 d.p.)}$$

2 $E(X) = \frac{1}{6}(1+2+4+5+7+8) = \frac{27}{6} = 4.5$

$$\begin{aligned} \text{Var}(X) &= \frac{1}{6}(1+2^2+4^2+5^2+7^2+8^2) - 4.5^2 \\ &= \frac{159}{6} - \frac{729}{36} = \frac{225}{36} = \frac{25}{4} = 6.25 \end{aligned}$$

By the central limit theorem $\bar{X} \approx \sim N\left(4.5, \frac{6.25}{20}\right)$, i.e. $\bar{X} \approx \sim N(4.5, 0.3125)$

$$P(\bar{X} < 4) \approx 0.1855 \text{ (4 d.p.)}$$

3 $X \sim N(1, 1)$ and by the central limit theorem $\bar{X} \sim N\left(1, \frac{1}{\sqrt{n}}\right)$

Standardise the sample mean.

$$P(\bar{X} < 0) = P(Z < -\sqrt{n}) \text{ and so require } P(Z > -\sqrt{n}) < 0.05$$

Using the table for the percentage points of the normal distribution:

$$P(Z = -1.645) = 0.05$$

$$\Rightarrow -\sqrt{n} < -1.645$$

$$\Rightarrow n > 2.706$$

So minimum sample size is $n = 3$ for the probability of a negative sample mean being less than 5%

4 Let the random variable X denote the number of sixes thrown by a student in 10 rolls of the dice, so

$$X \sim B\left(10, \frac{1}{6}\right)$$

$$E(X) = np = 10 \times \frac{1}{6} = \frac{5}{3}$$

$$\text{Var}(X) = np(1-p) = \frac{5}{3} \times \frac{5}{6} = \frac{25}{18}$$

By the central limit theorem $\bar{X} \approx \sim N\left(\frac{5}{3}, \frac{25}{18 \times 20}\right)$, i.e. $\bar{X} \approx \sim N\left(\frac{5}{3}, \frac{5}{72}\right)$

$$P(\bar{X} > 2) = 1 - P(\bar{X} < 2) \approx 1 - 0.8970 = 0.1030 \text{ (4 d.p.)}$$

5 a Let X be the number of buses that arrive in a 10-minute period, then $X \sim \text{Po}(2)$

$$P(X = 3) = \frac{e^{-2} 2^3}{3!} = 0.1804 \text{ (4 d.p.)}$$

5 b Let T be the number of buses that arrive in a two-hour period, so $T = 12\bar{X}$

By the central limit theorem $\bar{X} \approx \sim N\left(2, \frac{2}{12}\right)$, i.e. $\bar{X} \approx \sim N\left(2, \frac{1}{6}\right)$

$$P(T \geq 25) = P\left(\bar{X} \geq \frac{25}{12}\right)$$

$$P\left(\bar{X} \geq \frac{25}{12}\right) = 1 - P\left(\bar{X} < \frac{25}{12}\right) \approx 1 - 0.5809 = 0.4191 \text{ (4 d.p.)}$$

6 a Let the random variable X be the mass of an egg, then $X \sim N(60, 25)$ and $\bar{X} \sim N\left(60, \frac{25}{48}\right)$

$$P(\bar{X} > 59) = 1 - P(\bar{X} < 59) = 1 - 0.0829 = 0.9171 \text{ (4 d.p.)}$$

b The answer in part **a** is not an estimate because the sample is taken from a population that is normally distributed.

c Let the random variable Y is the number of double yolk eggs in a crate of 48 eggs, so $Y \sim B(48, 0.1)$

$$E(Y) = np = 48 \times 0.1 = 4.8$$

$$\text{Var}(Y) = np(1-p) = 4.8 \times 0.9 = 4.32$$

By the central limit theorem $\bar{Y} \approx \sim N\left(4.8, \frac{4.32}{30}\right)$, i.e. $\bar{Y} \approx \sim N(4.8, 0.144)$

The probability that the sample of 30 crates will contain fewer than 150 double-yolk eggs is $P(\bar{Y} < 5)$ as $30 \times 5 = 150$

$$P(\bar{Y} < 5) \approx 0.7009 \text{ (4 d.p.)}$$

7 Consider a sample of 100 cups of coffee, so $\bar{S} \sim N(4.9, 0.0064)$. One pack of milk powder will be sufficient, if $100\bar{S} < 500$, i.e. $\bar{S} < 5$

$$P(\bar{S} < 5) = 0.8944 \text{ (4 d.p.)}$$

- 8 Let the random variable be X , so by the central limit theorem $\bar{X} \approx \sim N\left(40, \frac{9}{n}\right)$

Required to find minimum n such that $P(\bar{X} > 42) < 0.05$

Standardise the sample mean using $Z = \frac{\bar{X} - \mu}{\sigma}$, $\mu = 40$ and $\sigma = \frac{3}{\sqrt{n}}$

So for $\bar{X} = 42$, $Z = \frac{(42 - 40)\sqrt{n}}{3} = \frac{2\sqrt{n}}{3}$ and $P(\bar{X} > 42) = P\left(Z > \frac{2\sqrt{n}}{3}\right)$

Using the table for the percentage points of the normal distribution;

$$P(Z > 1.6449) = 0.05$$

$$\text{So } \frac{2\sqrt{n}}{3} > 1.6449$$

$$\Rightarrow \sqrt{n} > 2.46735$$

$$\Rightarrow n > 6.0878\dots$$

So $n = 7$ is the minimum sample size required for $P(\bar{X} > 42) < 0.05$

- 9 Let the random variable be X , so by the central limit theorem $\bar{X} \approx \sim N\left(35, \frac{9}{20}\right)$

$$P(\bar{X} > 37) = 1 - P(\bar{X} < 37) \approx 1 - 0.9986 = 0.0014 \text{ (4 d.p.)}$$

- 10 a The table describes the distribution of X

x	0	1
$P(X = x)$	0.4	0.6

$$E(X) = 0.6, \text{ Var}(X) = 0.6 - 0.6^2 = 0.24$$

- b By the central limit theorem $\bar{X} \approx \sim N\left(0.6, \frac{0.24}{500}\right)$, i.e. $\bar{X} \approx \sim N(0.6, 0.00048)$

$$P(\bar{X} > 0.63) + P(\bar{X} < 0.57) = 1 - P(\bar{X} < 0.63) + P(\bar{X} < 0.57)$$

$$\approx 1 - 0.91454 + 0.08545 = 0.1709 \text{ (4 d.p.)}$$

10 c Required to find minimum n such that $P(0.57 < \bar{X} < 0.63) > 0.95$

Standardise the sample mean using $Z = \frac{\bar{X} - \mu}{\sigma}$, $\mu = 0.6$ and $\sigma = \sqrt{\frac{0.24}{n}}$

So for $\bar{X} = 42$, $Z = \frac{(42 - 40)\sqrt{n}}{3} = \frac{2\sqrt{n}}{3}$ and $P(\bar{X} > 42) = P\left(Z > \frac{2\sqrt{n}}{3}\right)$

So require $P\left(-\frac{0.03\sqrt{n}}{\sqrt{0.24}} < Z < \frac{0.03\sqrt{n}}{\sqrt{0.24}}\right) > 0.95$

$\Rightarrow 1 - 2P\left(Z < -\frac{0.03\sqrt{n}}{\sqrt{0.24}}\right) > 0.95$ (by the symmetry of the normal distribution)

$\Rightarrow P\left(Z < -\frac{0.03\sqrt{n}}{\sqrt{0.24}}\right) < 0.025$

Using the table for the percentage points of the normal distribution

$$P(Z < -1.960) = 0.025$$

$$\Rightarrow -\frac{0.03\sqrt{n}}{\sqrt{0.24}} < -1.960$$

$$\Rightarrow \sqrt{n} > \frac{1.960 \times \sqrt{0.24}}{0.03} \Rightarrow \sqrt{n} > 32.0066\dots$$

$$\Rightarrow n > 1024.42\dots$$

So $n = 1025$

11 The sample of bands from the new supplier has

$$\begin{aligned}\bar{x} &= \frac{\sum X}{100} \\ &= \frac{4715}{100} \\ &= 47.15\end{aligned}$$

and

$$\begin{aligned}s^2 &= \frac{\sum X^2}{100} - \left(\frac{\sum X}{100}\right)^2 \\ &= \frac{222910}{100} - 47.15^2 \\ &= 5.978 \\ s &= 2.445\end{aligned}$$

$$H_0: \mu = 46.5$$

$$H_1: \mu > 46.5$$

Using the estimator above for s , under the null hypothesis the distribution of the mean breaking stress of a sample of 100 bands can be modelled as $N\left(46.5, \frac{(2.445)^2}{100}\right)$

The probability a sample of 100 bands having a mean breaking stress of at least 47.15 is

$$Z = \frac{47.15 - 46.5}{\frac{2.445}{10}} = 2.659$$

$$2.659 > 1.645$$

Therefore, there is evidence that the new manufacturer's bands are stronger.

12

x	x^2	f	fx	fx^2
12	144	5	60	720
13	169	42	546	7098
14	196	210	290	41160
15	225	31	465	6975
16	256	12	192	3072
Totals		300	4205	59025

$$\begin{aligned}\bar{x} &= \frac{\sum fx}{\sum f} \\ &= \frac{4203}{300} \\ &= 14.01 \text{ mg}\end{aligned}$$

$$\begin{aligned}s^2 &= \frac{\sum X^2}{100} - \left(\frac{\sum X}{100}\right)^2 \\ &= \frac{59025}{300} - \left(\frac{4203}{300}\right)^2 \\ &= 196.75 - 196.2801 \\ &= 0.4699\end{aligned}$$

The variance of the mean is therefore $\frac{0.4699}{300}$

And the standard error is $\sqrt{\frac{0.4699}{300}} = 0.0396$

$$13 \text{ N}\left(\mu_A - \mu_B, \frac{8.0^2}{25} + \frac{8.0^2}{20}\right)$$

$$H_0: \mu_A - \mu_B = 0$$

$$H_1: \mu_A - \mu_B > 0$$

$$\begin{aligned} Z_{\text{Test}} &= \frac{\bar{X}_A - \bar{X}_B - 0}{8.0 \sqrt{\frac{1}{25} + \frac{1}{20}}} \\ &= \frac{44.2 - 40.9}{8.0 \sqrt{\frac{1}{25} + \frac{1}{20}}} \\ &= 1.375 \end{aligned}$$

$$Z_{\text{Test}} = 1.375 < Z_{\text{Crit}} = 1.645$$

There is no evidence for a difference in means between the two schools.

$$14 \text{ H}_0: \mu_{2010} = \text{£}9.10$$

$$H_1: \mu_{2010} > \text{£}9.10$$

For the random sample of 100 individual sales of unleaded fuel in 2010, the mean is 9.71 and the

standard deviation is $\frac{3.25}{\sqrt{100}}$

$$\begin{aligned} Z_{\text{Test}} &= \frac{9.71 - 9.10}{\left(\frac{3.25}{\sqrt{100}}\right)} \\ &= 1.8769 \end{aligned}$$

$$Z_{\text{Test}} = 1.8769 > Z_{\text{Crit}} = 1.645$$

Therefore, there is evidence that the 2010 sales of unleaded are higher than the 2009 sales of unleaded.

$$15 \text{ a } H_0: \mu_{\text{Drive}} - \mu_{\text{Walk}} = 0$$

$$H_1: \mu_{\text{Drive}} - \mu_{\text{Walk}} \neq 0$$

$$\begin{aligned} Z_{\text{Test}} &= \frac{\bar{X}_{\text{Drive}} - \bar{X}_{\text{Walk}}}{\sqrt{\frac{\text{Var}_{\text{Drive}}}{n_{\text{Drive}}} + \frac{\text{Var}_{\text{Walk}}}{n_{\text{Walk}}}}} \\ &= \frac{52 - 47}{\sqrt{\frac{60.2}{30} + \frac{55.8}{36}}} \\ &= 2.651 \end{aligned}$$

$$Z_{\text{Test}} = 2.651 > Z_{\text{Crit}} = 1.645$$

Therefore, there is evidence that those who drive to work have a higher heart rate.

- b** Assume normal distribution or assume sample sizes large enough to use the central limit theorem; assume individual results are independent; assume $\sigma_2 = s_2$ for both populations.

$$\text{c } \bar{X}_{\text{Drive}} = 52$$

$$\sum X = 1560$$

$$s^2 = 60.2$$

$$\sum X^2 = s^2(n-1) + n\bar{X}_{\text{Drive}}^2 = 82865.8$$

$$\sum X_{\text{New}} = 1560 + 55 = 1615$$

$$\sum X_{\text{New}}^2 = 82865 + 55^2 = 85890.8$$

Therefore, we can calculate the unbiased estimator of the variance of the new sample using these values to find

$$s_{\text{New}}^2 = 58.5$$

Challenge

$X_1 + \dots + X_n \sim N(n\mu, n\sigma^2)$ and so

$$\bar{X} = \frac{1}{n}(X_1 + \dots + X_n) \sim N\left(\frac{n\mu}{n}, \frac{n\sigma^2}{n^2}\right) = N\left(\mu, \frac{\sigma^2}{n}\right)$$