



Mark Scheme (**Results**)

Summer 2016

Pearson Edexcel GCE Statistics
S4

(6686/01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2016

Publications Code: **6686_01_1606_MS**

All the material in this publication is copyright

© Pearson Education Ltd 2016

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. **Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.**
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

PEARSON EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - **d... or dep** – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper or ag- answer given
 - \square or d... The second mark is dependent on gaining the first mark
4. **All A marks are 'correct answer only' (cao.)**, unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

June 2016
6686 Statistics S4
Mark Scheme

Question Number	Scheme	Marks
1(a)	$d: 5\ 9\ 5\ 2\ 0\ 0\ 8\ 5\ 6\ 6$ $\bar{d} = \frac{\Sigma d}{n} = 4.6$ $s^2 = \frac{296 - 10 \times 4.6^2}{9} = 9.378$ $H_0: \mu_d = 2 \quad H_1: \mu_d > 2$ $t = \pm \frac{4.6 - 2}{\sqrt{\frac{9.378}{10}}} = \pm 2.6848$ $t_{9(5\%)} = \pm 1.833\dots$ There is evidence to reject H_0 . There is sufficient evidence to support the designers claim.	M1 M1 M1 B1 M1 A1 B1 A1ft (8)
(b)	The differences in weights are normally distributed.	B1 (1)
Notes		Total 9
(a)	M1 for attempting the ds M1 for attempting \bar{d} M1 for s_d or s_d^2 B1 for both hypotheses correct in terms of μ or μ_d .(allow a defined symbol) M1 for attempting the correct test statistic $\frac{\bar{d}}{s_d / \sqrt{10}}$ A1 awrt 2.68 B1 awrt 1.83 A1ft for a correct comment in context	
(b)	B1 for a comment that mentions “differences” and “normal” distribution	

Question Number	Scheme	Marks
<p>2. (a)</p> <p>(b)</p>	<p>$H_0 : \mu = 1.2 \quad H_1 : \mu > 1.2$</p> <p>$t_8(5\%) = 1.860$</p> <p>$\bar{m} = 1.28888..$</p> <p>$t = \frac{1.28... - 1.2}{\sqrt{\frac{0.031111}{9}}} = 1.511$ awrt 1.51</p> <p>Not significant. There is not sufficient evidence that the mean <u>weight of piglets</u> is greater than 1.2 kg</p> <p>$H_0 : \sigma^2 = 0.09 \quad H_1 : \sigma^2 \neq 0.09 \quad [H_0 : \sigma = 0.3 \quad H_1 : \sigma \neq 0.3]$</p> <p>$s^2 = \frac{15.2 - 9 \times \left(\frac{11.6}{9}\right)^2}{8} = 0.031111$</p> <p>$[\chi_8^2(0.25) = 17.535] \quad \chi_8^2(0.975) = 2.18$</p> <p>Critical region $\frac{(n-1)s^2}{\sigma^2} \sim \chi_8^2$ test statistic = 2.7654... awrt 2.77</p> <p>2.77 is not in the critical region. There is no evidence that the standard deviation of the weights of <u>piglets</u> is different to 0.3</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1 A1ft</p> <p>A1</p> <p>A1</p> <p>(7)</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>M1A1</p> <p>A1</p> <p>(6)</p>
Notes		Total 13
<p>(a)</p> <p>(b)</p>	<p>B1 both hypotheses</p> <p>M1 for attempting the correct statistic</p> <p>A1ft follow through their s^2</p> <p>A1 awrt 1.51</p> <p>B1 both hypotheses, must be two tail</p> <p>B1 awrt 0.0311</p> <p>B1 NB allow 2.733 for one tail hypotheses. (no hypotheses gains B0)</p> <p>M1 for a correct test statistic</p> <p>NB one tail test can get B0 B1 B1 (2.733)B0 M1 A1 A1</p>	

Question Number	Scheme	Marks
<p>3. (a)</p> <p>(b)</p> <p>(c)</p>	<p>$X =$ No of soft centres. $X \sim B(20, 0.5)$ Critical region $X \leq 5$ or $X \geq 15$</p> <p>$P(\text{Type I error}) = P(X \leq 5 p = 0.5) + P(X \geq 15 p = 0.5)$ $= 0.0207 + 0.0207 = 0.0414$</p> <p>$P(\text{Type II error}) = P(X < 15 p = 0.25) - P(X < 6 p = 0.25)$ $= 1 - 0.6172 = 0.3828$</p>	<p>B1B1 (2)</p> <p>M1 A1 (2)</p> <p>M1 A1 (2)</p>
Notes		Total 6
<p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>B1 $X \leq 5$ B1 $X \geq 15$</p> <p>M1 Adding their two CR together or a correct answer A1 awrt 0.0414</p> <p>M1 FT their CR A1 awrt 0.383</p>	

Question Number	Scheme	Marks
<p>4. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>Size of test $A = P(Y \leq 2)$ $= 0.0547$</p> <p>Size of test $B = P(\text{Rejecting } H_0 p = 0.5)$ $= P(X = 0) + (1 - P(X = 0)) \times P(X = 0)$ $= 0.5^5 + (1 - 0.5^5)(0.5^5)$ $= 0.03125 + (0.96875)(0.03125)$ $= 0.0615/0.0614$</p> <p>Power function of test $B = P(0 \text{ long screws in first } 5) + P(0 \text{ long screws in second } 5 > 0 \text{ long screws in first } 5)$ $= P(X = 0 p) + [1 - P(X = 0 p)] P(X = 0 p)$ $= (1 - p)^5 + [1 - (1 - p)^5](1 - p)^5$ $= 2(1 - p)^5 - (1 - p)^{10}$</p> <p>$r = 0.68$</p> <p>Test A as it is more powerful for values of $p < 0.4$</p>	<p>B1 (1)</p> <p>M1 A1 A1 (3)</p> <p>M1 A1 (2)</p> <p>B1 (1)</p> <p>M1 A1 (2)</p>
	Notes	Total 9
<p>(b)</p> <p>(c)</p> <p>(e)</p>	<p>M1 for a correct expression/selection of probabilities A1 for a correct expression in terms of probabilities. Allow $0.0312 + (0.9688)(0.0312)$</p> <p>M1 for a correct expression A1 for a correct expression in terms of p</p> <p>M1 for reason based on the power function A1 test A</p>	

Question Number	Scheme	Marks
<p>5. (a)</p> <p>(b)</p> <p>(c)</p>	<p>$H_0 : \sigma^2_X = \sigma^2_Y \quad H_1 : \sigma^2_X \neq \sigma^2_Y$</p> <p>$F_{8,5} = \frac{6.76^2}{5.42^2} = 1.556$</p> <p>$F_{8,5}$ is 4.82</p> <p>There is evidence that the variances are the same.</p> <p>$H_0 : \mu_X = \mu_Y + 5 \quad H_1 : \mu_X > \mu_Y + 5$</p> <p>$s_p^2 = \frac{8 \times 6.76^2 + 5 \times 5.42^2}{13}, = 39.42... \quad \text{or} \quad s_p = 6.278...$</p> <p>$(t_{13} =)(\pm) \frac{14.8 - 7.2 - 5}{s_p \sqrt{\frac{1}{9} + \frac{1}{6}}} = (\pm)0.78578...$ awrt 0.786</p> <p>Critical value $t_{13} (2.5\%) = 1.771$</p> <p>There is no evidence to Reject H_0</p> <p>There is evidence that the fire brigade in X does not take more than 5 minutes longer than those in Y.</p> <p>Test in part (b) requires the variances to be equal. The test in part (a) showed that the variances could be assumed to be equal.</p>	<p>B1</p> <p>M1A1</p> <p>B1</p> <p>A1</p> <p>(5)</p> <p>B1</p> <p>M1 A1</p> <p>M1</p> <p>M1dA1</p> <p>B1</p> <p>A1cso</p> <p>(8)</p> <p>B1</p> <p>(1)</p>
	notes	Total 14
<p>(a)</p> <p>(b)</p>	<p>B1 both hypotheses</p> <p>M1 Allow use of 6.76 and 5.42 instead of 6.76^2 and 5.42^2</p> <p>A1 awrt 1.56</p> <p>B1 both hypotheses</p> <p>M1 allow use of 6.76 and 5.42 instead of 6.76^2 and 5.42^2</p> <p>A1 awrt 39.4 or 6.28</p> <p>B1 allow p value 0.650 instead of critical value</p> <p>M1 use of correct formula with their S_p – condone missing 5</p> <p>M1 use of correct formula with their S_p</p>	

Question Number	Scheme	Marks
6.(a)	$E(Y) = 2E(\bar{X})$ $= 2 \times \frac{a}{2}$ $= a$	M1 A1cso (2)
(b)	$E(M) = \int_0^a \frac{nm^n}{a^n} dm$ $= \left[\frac{nm^{n+1}}{a^n(n+1)} \right]_0^a$ $= \frac{na}{n+1}$	M1 A1 (2)
(c)	$\text{Var}(M) = \int_0^a \frac{nm^{n+1}}{a^n} dm - \left(\frac{na}{n+1} \right)^2$ $= \left[\frac{nm^{n+2}}{a^n(n+2)} \right]_0^a - \frac{n^2 a^2}{(n+1)^2}$ $= na^2 \left(\frac{(n+1)^2 - n(n+2)}{(n+1)^2(n+2)} \right)$ $= \frac{na^2}{(n+2)(n+1)^2}$	M1A1 M1d A1cso (4)
(d)	$E(S) = \frac{n+1}{n} E(M) = \frac{n+1}{n} \times \frac{na}{n+1} = a$ $\text{Var}(S) = \left(\frac{n+1}{n} \right)^2 \frac{na^2}{(n+2)(n+1)^2} = \frac{a^2}{n(n+2)}$ $\text{Var}(Y) = 4 \text{Var}(\bar{X})$ $= 4 \times \frac{a^2}{12n}$ $= \frac{a^2}{3n}$ <p>As $n > 1$ $n(n+2) > 3n$; therefore $\text{Var}(S) < \text{Var}(Y)$ $\therefore S$ is the better estimator</p>	B1 B1 M1 A1 M1;M1 A1cso (7)
Total 15		

	notes	
(a)	M1 for $2E(\bar{X})$ A1 For $2 \times \frac{a}{2}$ leading to a	
(b)	M1 attempting to integrate correct expression	
(c)	M1 for attempting to integrate a correct expression for $E(X^2)$ A1 correct $E(X^2)$ M1d dependent on previous M mark, using correct formula for $\text{Var}(M)$	
(d)	B1 for $\frac{n+1}{n}E(M) = a$ or $\frac{n+1}{n} \times \frac{na}{n+1} = a$ M1 using $4 \text{Var}(\bar{X})$ NB Failure to show S is unbiased gains a maximum of 5/7 lose first B1 and final A1	

Question Number	Scheme	Marks
7	$\bar{x} - 2.262 \frac{s}{\sqrt{10}} = 28.5$ $\bar{x} + 2.262 \frac{s}{\sqrt{10}} = 48.7$ $2\bar{x} = 48.7 + 28.5 \text{ or } 2.262 \frac{s}{\sqrt{10}} = \frac{1}{2}(48.7 - 28.5)$ $s = 14.1198... \text{ (} s^2 = 199.36 \text{)}$ $\left\{ \frac{9(14.1198^2)}{23.589}, \frac{9(14.1198^2)}{1.735} \right\}$ $= (76.0659..., 1034.19...)$	B1 M1 A1 M1 A1 M1 B1 B1 A1 (9)
	notes	Total 9
	B1 awrt 2.262 M1 $\bar{x} - t \text{ value } \frac{s}{\sqrt{10}} = 28.5$ A1 both equations correct M1 solving simultaneous leading to a value for \bar{x} or s A1 awrt 14.1 or awrt 199 M1 $\frac{9(s^2)}{\chi^2 \text{ value}}$ B1 23.589 B1 1.735 A1 awrt 76.1 and awrt 1030	

