

1. In a factory, an inspector checks a random sample of 30 mugs from a large batch and notes the number, X , which are defective. He then deals with the batch as follows.
- If $X < 2$, the batch is accepted.
 - If $X > 2$, the batch is rejected.
 - If $X = 2$, the inspector selects another random sample of only 15 mugs from the batch. If this second sample contains 1 or more defective mugs, the batch is rejected. Otherwise the batch is accepted.

It is given that 5% of mugs are defective.

- i.
- a. Find the probability that the batch is rejected after just the first sample is checked. [3]
- b. Show that the probability that the batch is rejected is 0.327, correct to 3 significant figures. [5]
- ii. Batches are checked one after another. Find the probability that the first batch to be rejected is either the 4th or the 5th batch that is checked. [3]

2. Froox sweets are packed into tubes of 10 sweets, chosen at random. 25% of Froox sweets are yellow.

- i. Find the probability that in a randomly selected tube of Froox sweets there are
- a. exactly 3 yellow sweets, [3]
- b. at least 3 yellow sweets. [2]
- ii. Find the probability that in a box containing 6 tubes of Froox sweets, there is at least 1 tube that contains at least 3 yellow sweets. [3]

3. (a) Write down and simplify the first four terms in the expansion of $(x + y)^7$.
Give your answer in ascending powers of x . [2]
- (b) $\frac{x}{y}$ [2]
Given that the terms in x^2y^5 and x^3y^4 in this expansion are equal, find the value of $\frac{x}{y}$.
- (c) A hospital consultant has seven appointments every day. The number of these appointments which start late on a randomly chosen day is denoted by L . The variable L is modelled by the distribution $B\left(7, \frac{3}{8}\right)$. Show that, in this model, the hospital consultant is equally likely to have two appointments start late or three appointments start late. [3]

4. On average, 40% of candidates pass a certain test on the first attempt.
- Three candidates take the test. The number who pass on the first attempt is denoted by X .
- (a) State an appropriate model for X , including the values of any parameters. [1]
- (b) State an appropriate model for X , including the values of any parameters. [2]
- (c) Suggest a reason why one of these assumptions might not be true in practice. [1]

You should now assume that both these assumptions are true.

- (d) Find the probability that exactly 2 of the 3 candidates pass the test. [1]

All candidates who fail the test take a re-test and, on average, 60% of these candidates pass. Assume that the same two assumptions are satisfied as for the original test.

- (e) Find the probability that all three candidates pass, either on the test or on the re-test. [3]

5. A bag contains 100 black discs and 200 white discs. Paula takes five discs at random, without replacement. She notes the number X of these discs that are black.

(a) Find $P(X = 3)$. [2]

Paula decides to use the binomial distribution as a model for the distribution of X .

(b) Explain why this model will give probabilities that are approximately, but not exactly, correct. [3]

(c) Paula uses the binomial model to find an approximate value for $P(X = 3)$. Calculate the percentage by which her answer will differ from the answer in part (b). [2]

Paula now assumes that the binomial distribution is a good model for X . She uses a computer simulation to generate 1000 values of X . The number of times that $X = 3$ occurs is denoted by Y .

(d) Calculate estimates of the limits between which two thirds of the values of Y will lie. [3]

6. The probability that Janice sees a kingfisher on any particular day is 0.3. She notes the number, X , of days in a week on which she sees a kingfisher.

(a) State one necessary condition for X to have a binomial distribution. [1]

Assume now that X has a binomial distribution.

(b) Find the probability that, in a week, Janice sees a kingfisher on exactly 2 days. [1]

Each week Janice notes the number of days on which she sees a kingfisher.

(c) Find the probability that Janice sees a kingfisher on exactly 2 days in a week during at least 4 of 6 randomly chosen weeks. [3]

END OF QUESTION paper

Mark scheme

| Question | Answer/Indicative content | Marks | Part marks and guidance |
|----------|---|---|---|
| 1 | <p>i (a) $X \sim B(30, 0.05)$ seen or implied</p> <p>$P(X > 2) = 1 - 0.8122$ alone or $1 - (0.95^{30} + 30 \times 0.95^{29} \times 0.05 + {}^{30}C_2 \times 0.95^{28} \times 0.05^2)$</p> <p>i = 0.1878 or 0.188 (3 sfs)</p> <p>i (b) Addition method: $X \sim B(30, 0.05)$ & $Y \sim B(15, 0.05)$ stated or implied</p> <p>i $P(X = 2) = (0.8122 - 0.5535)$ or ${}^{30}C_2 \times 0.95^{28} \times 0.05^2$ or 0.2587/6 OR $P(Y > 1) = (1 - 0.95^{15})$ or 0.5367</p> | <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> | <p>eg by 0.8122 or $1 - 0.5535$ or $0.95^r \times 0.05^s$ ($r, s > 1$) Allow $B(30, 0.95)$ or $B(30, 0.5)$ for B1 30×0.05 alone insufficient for B1</p> <p>nC_r: insufficient for B1</p> <p>Examiner's Comments</p> <p>A few misread the question, using $n = 15$ throughout, instead of both $n = 30$ and $n = 15$. The problem here was not generally the use of the binomial distribution but the need to see through the context and recognise that a binomial calculation was appropriate. In this part there was a distinct advantage to using tables rather than the formula (which is a much longer method in this case). Some candidates chose to use the formula and made arithmetical errors. Using either method, some candidates used $B(30, 0.95)$ instead of $B(30, 0.05)$. Others used the correct distribution but, in order to find $P(X < 2)$, they found $1 - P(X = 1)$ instead of $1 - P(X \leq 1)$. Some omitted the binomial coefficients.</p> <p>NB eg 0.0362 implies $B(15, 0.05)$ see below</p> <p>$P(X = 2) = (0.8122 - 0.5535)$ or ${}^{30}C_2 \times 0.95^{28} \times 0.05^2$ or 0.2587/6</p> <p>If $n = 15$; $B(15, 0.05)$ B1</p> <p>$1 - (0.95^{15} + 15 \times 0.95^{14} \times 0.05 + {}^{15}C_2 \times 0.95^{13} \times 0.05^2)$ M1</p> <p>= 0.0362 A0</p> <p>Subtraction methods: $X \sim B(30, 0.05)$ & $Y \sim B(15, 0.05)$ stated or impl B1</p> |

| | | | | | |
|--|--|--|-------------------------------|---|--|
| | | <p>i "0.2587/6" x "0.5367" or 0.1388</p> <p>i $P(X > 2) + P(X = 2) \times P(Y > 1)$ = "0.1878" + "0.1388" alone</p> <p>i = 0.327 (3 sf) AG</p> <p>i For A1 must see correct wking or 0.3265/6...</p> <p>i</p> <p>i Alternative scheme for the case where $n = 15$ is used for both distr's</p> <p>i</p> | <p>M1</p> <p>M1</p> <p>A1</p> | <p>fully correct method for $P(X = 2) \times P(Y \geq 1)$</p> <p>[their (a) + any p] alone, but dep 1st M1</p> <p>If ans 0.327, check whether it comes from a correct method (possibly not in MS) or clearly comes from an incorrect method eg $(0.4465 + 0.2587) \times 0.4633 = 0.327$ (ie $(P(X > 2) + P(X = 2)) \times P(Y = 0)$ B1M1M0M0A0</p> <p><u>Examiner's Comments</u></p> | <p>OR $P(Y = 0) = 0.95^{15}$ Binomial Distribution or 0.4633 M1</p> <p>fully correct method for $P(X = 2) \times P(Y = 0)$ "0.2587" x "0.4633" or 0.1199/8 M1</p> <p>$1 - (P(X = 0, 1) + P(X = 2) \times P(Y = 0)) = 1 -$ ("0.5535" + "0.1199") OR $P(X > 2) - P(X = 2) \times P(Y = 0) = (1 -$ "0.5535") - "0.1199" dep 1st M1 M1</p> <p>= 0.327 (3 sf) AG A1</p> <p>Do not use marks from a mixture of 3rd column and other columns. Decide which column would give most marks and mark according to that method.</p> <p>If $n = 15$ for both distr's, see next page NB If 0.1392 seen, it comes from given answer - (i)(a) (ie 0.3270 - 0.1878).</p> <p>If $n = 15$ for both distr's B(15, 0.05) B0</p> <p>$P(X = 2) = {}^{15}C_2 \times 0.05^2 \times 0.95^{13}$ or 0.1348 OR $P(Y > 1) = 1 - 0.95^{15}$ or 0.5367 M1 "0.1348"x"0.5367" or 0.0723 correct method M1 their (i)(a) + "0.0732" Dep 1st M1 M1 = 0.1085 A0 NB Also mark subtraction methods if seen.</p> |
|--|--|--|-------------------------------|---|--|

| | | | | | |
|--|----|--|-----------|--|---|
| | | | | <p>A few misread the question, using $n = 15$ throughout, instead of both $n = 30$ and $n = 15$.</p> <p>Few candidates were able to work their way through this part correctly. Some candidates found $P(Y \geq 1) = 1 - P(Y = 0 \text{ or } 1)$ or $P(Y \geq 1) = 1 - P(Y = 1)$. Some candidates were close to being correct, but found $P(X > 2) + P(X = 2) \times P(Y = 0)$ instead of $P(X > 2) + P(X = 2) \times (1 - P(Y = 0))$. Many complicated matters by using the same letter (X) for the number of defective mugs in the second sample as well as in the first.</p> | Binomial Distribution |
| | ii | Any use of 0.327 or their (i)(b) for 1 st M1 | M1 | | |
| | | $(1 - 0.327)^3 \times 0.327 + (1 - 0.327)^4 \times 0.327$ | | | $1 - 0.673^5 - (1 - 0.673^3)$ oe |
| | ii | Allow "correct" use of their (i)(a) or (i)(b) for 2 nd M1 | M1 | $(0.5535 + 0.2586 \times 0.4633)^3 \times 0.327 +$ $(0.5535 + 0.2586 \times 0.4633)^4 \times 0.327$ | Allow any use of their (i)(b) for 1 st M1 then if "correct" use, also 2 nd M1 |
| | ii | = 0.167 (3 sf) | A1 | | Allow use of their (i)(a) in "correct" method for M0M1A0 No marks for use of 0.95 & 0.05 |
| | ii | | | <p>Examiner's Comments</p> <p>A few misread the question, using $n = 15$ throughout, instead of both $n = 30$ and $n = 15$.</p> <p>This is an example of a common, "two-layered" type of question, requiring the use of a previously obtained figure as the value of p in a geometric (or binomial) calculation. Candidates would benefit from being taught to look out for such questions. Many candidates used the correct geometric structure but with $p = 0.05$ or with a probability equal to their answer to part (i)(a) instead of (i)(b). Others attempted to use a binomial distribution instead of a geometric.</p> | |
| | | Total | 11 | | |

| | | | | | |
|---|----|--|--|---|---|
| 2 | i | <p>(a) Binomial seen or implied</p> <p>0.7759 – 0.5256 or ${}^{10}C_3 \times (1 - 0.25)^7 \times 0.25^3$</p> <p>= 0.250 (3 sf)</p> <p>(b) 1 – 0.5256 or</p> <p>$1 - ((1 - 0.25)^{10} + 10(1 - 0.25)^9 \times 0.25 + {}^{10}C_2(1 - 0.25)^8 \times 0.25^2)$</p> <p>= 0.4744 or 0.474 (3 sf)</p> | B1 M1 A1 M1 A1 | <p>by tables or ${}^{10}C_3$ or ${}^{10}C_7$</p> <p>Allow 0.25</p> <p>Examiner's Comments</p> <p>This question was usually answered correctly, usually by the formula and sometimes using tables. A few candidates omitted the binomial coefficient while others only read one value (0.7759) from the table, i.e $P(X \leq 3)$). Others found $1 - 0.7759$.</p> <p>or $P(X = 3,4,5,6,7,8,9,10)$ all correct terms</p> <p>Allow ${}^{10}C_8$ instead of ${}^{10}C_2$</p> <p>Examiner's Comments</p> <p>A few candidates found $1 - 0.7759$ (i.e. $P(X > 3)$), or just gave 0.7759 as the answer. Others found $1 - P(X = 2)$. Some (correctly) used the formula to find $1 - P(X = 0, 1 \text{ or } 2)$. This is not a particularly long method, but it is significantly longer than finding $1 - P(X \leq 2)$ using the tables. Others attempted various methods not involving the binomial distribution at all.</p> | <p>Binomial Distribution or by $0.25^a \times 0.75^b$ ($a + b = 10$)</p> <p>NOT $1 - 0.7759$ ($P(X > 3)$ from table)</p> |
| | ii | <p>0.4744 or 0.474) or 0.5256 or 0.526 seen</p> <p>$1 - (1 - "0.4744")^6$ oe</p> <p>= 0.979 (3 sf)</p> | M1 M1 A1f | <p>Their (i)(b) seen, or result of $1 - (i)(b)$ seen</p> <p>or $P(X = 1,2,3,4,5,6)$ all correct terms seen</p> <p>fit from (i)(b)</p> <p>Examiner's Comments</p> <p>A good number of candidates answered this question</p> | <p>eg $B(6, 0.474)$ or $P(X \geq 3) = 0.474$</p> |


| | | | | | | | |
|---|---|--|---|--|---|--|--|
| | | | | <p>correctly. Many, however, found this very typical question difficult, not appreciating its two-layered structure. These candidates gave answers such as</p> $\frac{1}{6} \times {}^{10}C_3 \times 0.25^3 \times 0.75^7$ <p>Other candidates used their answer to part (i)(b) (thus gaining one mark) but used it in incorrect ways, such as dividing it by 6 or by raising it to the power of 6. Some subtracted it from 1, which is correct, but failed to take the next two steps correctly. A few candidates used their answer to part (i)(a) instead of (i)(b).</p> | | | |
| | | Total | 8 | | | | |
| 3 | a | $y^7 + 7xy^6 + 21x^2y^5 + 35x^3y^4$ | B2 (AO1.1) 1.1) [2] | <table border="1"> <tr> <td>B1 for three terms correct</td> <td></td> </tr> </table> | B1 for three terms correct | | |
| B1 for three terms correct | | | | | | | |
| | b | $21x^2y^5 = 35x^3y^4$ $\frac{x}{y} = \frac{3}{5}$ <small>or 0.6</small> | M1 (AO3.1a) A1 (AO1.1) [2] | <table border="1"> <tr> <td>Equate their terms in x^2y^5 and x^3y^4</td> <td></td> </tr> </table> | Equate their terms in x^2y^5 and x^3y^4 | | |
| Equate their terms in x^2y^5 and x^3y^4 | | | | | | | |
| | c | $P(L=2) = {}_7C_2 \left(\frac{3}{8}\right)^2 \left(\frac{5}{8}\right)^5$ and $P(L=3) = {}_7C_3 \left(\frac{3}{8}\right)^3 \left(\frac{5}{8}\right)^4$ Let $p = \frac{3}{8}$ and $1-p = \frac{5}{8}$, then $\frac{p}{1-p} = \frac{\frac{3}{8}}{\frac{5}{8}} = \frac{3}{5}$ | B1 (AO3.3) M1 (AO3.4) E1 (AO2.1) | <table border="1"> <tr> <td>Connect to part (b)</td> <td></td> </tr> </table> | Connect to part (b) | | |
| Connect to part (b) | | | | | | | |

| | | | | | |
|---|---|--|---------------------------------|--|-----------------------|
| | | so from part (b) $21p^2(1-p)^5 = 35p^3(1-p)^4$ and $P(L = 2) = P(L = 3)$ | [3] | AG | Binomial Distribution |
| | | Total | 7 | | |
| 4 | a | Bin(3, 0.4) | B1(AO 3.3) [1] | | |
| | b | P(Pass) same for all candidates Each candidate's result is indep of the other results | B1(AO 2.4) B1(AO 2.4) [2] | oe oe Not just "Independent" | |
| | c | eg Some candidates may be more able than others | B1(AO 3.5b) [1] | oe; or eg 'work harder' | |

| | | | | | | |
|---|--|--------------|---|--|---------------------|-----------------------|
| | | d | 0.288 | B1(AO 3.4) [1] | BC | Binomial Distribution |
| | | e | 0 & 3, 1 & 2, 2 & 1, 3 seen or implied $0.216 \times 0.216 + 0.432 \times 0.36 + 0.288 \times 0.6 + 0.064$ $= 0.439$ (3 sf) | M1(AO 3.1a) A1(AO 3.4) A1(AO 1.1) [3] | all correct | |
| | | Total | | 8 | | |
| 5 | | a | $P(X = 3) = {}^5C_3 \times \frac{100}{300} \times \frac{99}{299} \times \frac{98}{298} \times \frac{200}{297} \times \frac{199}{296}$ $= 0.164318883 = 0.164$ (3 sf) | M1(AO 1.1a) A1(AO 1.1) [2] | or equiv methods | |
| | | b | P(disc is black) changes each trial (because no replacement) oe But change in prob is small oe Hence bin gives approx, but not exact, probs oe | E1(AO 2.4) E1(AO 2.4) E1(AO 3.5b) [3] | | |
| | | c | $P(X = 3 \text{ using bin}) = {}^5C_3 \times \left(\frac{2}{3}\right)^2 \left(\frac{1}{3}\right)^3$ <input type="text"/> <input type="text"/> (= 0.164609053) | | | |

| | | | | | |
|---|--------------|---|--|--|--|
| | | $\frac{'0.164609053' - '0.164318883'}{'0.164318883'} \times 100$ <p>= 0.177%</p> | M1(AO 3.4) A1(AO 1.1) [2] | ft their values for M1 cao | Binomial Distribution |
| d | X ~ Normal | $\mu = 1000 \times 0.164609053 (= 164.609053)$ $\sigma^2 = 1000 \times 0.164609053 \times (1 - 0.164609053)$ <input type="text" value="137.5129127"/> (= 137.5129127) | M1(AO 3.3) M1(AO 1.1a) A1(AO 1.1) [3] | both np and npq correct method ft their μ and $\sqrt{\sigma^2}$. Allow rounding to 3 sf Allow (150 – 155) to (174 – 180) | ft their 0.164.6..... Allow use of 0.1643.... $165 \pm \sqrt{138}$ or better |
| | Total | | 10 | | |

| | | | | | | | |
|------------------------|---|---|---|--|------------------------|---|-----------------------|
| 6 | a | <p>Prob of seeing a kingfisher is the same each day</p> <p>OR Seeing a kingfisher on one day is independent of other days</p> | <p>B1 (AO3.5b)</p> <p>[1]</p> | <table border="1" data-bbox="1128 73 1646 424"> <tr> <td data-bbox="1128 73 1386 424">oe In context</td> <td data-bbox="1386 73 1646 424">Not: Prob of seeing kingfisher is indep</td> </tr> </table> <p><u>Examiner's Comments</u></p> <p>Despite almost identical questions having been asked in many examples of paper 4732 over many years, most candidates did not score the mark. Many quoted from their textbooks, to no avail, for example "Repeated trials" or "Only two possible outcomes, seeing or not seeing a kingfisher" or "There must be a fixed number of days". Many gave a correct answer, but not in context, such as "Independent trials" or "Constant probability of success." None of the above scored the mark. Some came a little nearer, but confused the two correct conditions, e.g. "The probability of seeing a kingfisher on a certain day is independent of other days."</p> | oe In context | Not: Prob of seeing kingfisher is indep | Binomial Distribution |
| oe In context | Not: Prob of seeing kingfisher is indep | | | | | | |
| | b | 0.318 (3 sf) | <p>B1 (AO3.4)</p> <p>[1]</p> | <table border="1" data-bbox="1128 1035 1646 1153"> <tr> <td data-bbox="1128 1035 1386 1153">BC Allow 0.32 or 0.317</td> <td data-bbox="1386 1035 1646 1153"></td> </tr> </table> <p><u>Examiner's Comments</u></p> <p>Most candidates understood how to calculate a binomial probability. Some used the formula, while others used the calculator function, giving only the answer. This is acceptable because "detailed reasoning" is not asked for.</p> | BC Allow 0.32 or 0.317 | | |
| BC Allow 0.32 or 0.317 | | | | | | | |

| | | | | | |
|--|--|--|---|--|--|
| | | | | Many candidates truncated, rather than rounded, their three significant figure answer. A few candidates just found 0.3^2 . | Binomial Distribution |
| | | | | <div style="border: 1px solid black; padding: 5px;"> <p>or B(6, their (b)) stated</p> <p>or $P(X = 4, 5, 6)$ attempted using $p = 0.318$ or their (b)</p> <p>Allow 0.0845 to 0.0875 Allow 2 sf BC</p> </div> | |
| | | | c | <p>0.318 or their (b) used in a calculation</p> <p>$1 - P(X \leq 3)$ using $p = 0.318$ or their (b)</p> <p>= 0.0854</p> | <p>M1 (AO3.1b)</p> <p>M1 (AO1.1)</p> <p>A1 (AO1.1)</p> <p>[3]</p> <p><u>Examiner's Comments</u></p> <p>Many candidates understood that they needed to use their answer from part (b), but many did not do so correctly. Some used the formula without the relevant coefficients.</p> <div style="text-align: center;">  </div> <p>Candidates should be aware that they can use the binomial distribution function on their calculator, thus avoiding the possibility arithmetic errors in binomial problems.</p> <p>Some candidates found $P(\text{exactly 4 days})$ instead of $P(\text{at least 4 days})$. Others just found 0.318^4. A few candidates misunderstood the question and found $P(\text{at least 8 days out of 42})$.</p> |

| | | | | | |
|--|--|--|-------|---|-----------------------|
| | | | Total | 5 | Binomial Distribution |
|--|--|--|-------|---|-----------------------|