



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

0606/12

Paper 1

February/March 2020

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Blank pages are indicated.

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series

$$u_n = a + (n-1)d$$

$$S_n = \frac{1}{2}n(a + l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series

$$u_n = ar^{n-1}$$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY*Identities*

$$\begin{aligned} \sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A \end{aligned}$$

Formulae for $\triangle ABC$


$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} bc \sin A$$

3

- 1 (a) On the axes below sketch the graph of $y = -3(x-2)(x-4)(x+1)$, showing the coordinates of the points where the curve intersects the coordinate axes. [3]



- (b) Hence find the values of x for which $-3(x-2)(x-4)(x+1) > 0$. [2]

4

2 Find the values of k for which the line $y = kx + 3$ is a tangent to the curve $y = 2x^2 + 4x + k - 1$. [5]

5


- 3 The first 3 terms in the expansion of $(3-ax)^5$, in ascending powers of x , can be written in the form $b-81x+cx^2$. Find the value of each of a , b and c . [5]

6

- 4 The tangent to the curve $y = \ln(3x^2 - 4) - \frac{x^3}{6}$, at the point where $x = 2$, meets the y -axis at the point P . Find the exact coordinates of P . [6]

5 DO NOT USE A CALCULATOR IN THIS QUESTION.

In this question all lengths are in centimetres.



The diagram shows the isosceles triangle ABC , where $AB = AC$ and $BC = 2 + 4\sqrt{3}$. The height, AD , of the triangle is $5 - \sqrt{3}$.

- (a) Find the area of the triangle ABC , giving your answer in the form $a + b\sqrt{3}$, where a and b are integers. [2]

- (b) Find $\tan \angle ABC$, giving your answer in the form $c + d\sqrt{3}$, where c and d are integers. [3]

- (c) Find $\sec^2 \angle ABC$, giving your answer in the form $e + f\sqrt{3}$, where e and f are integers. [2]

6 Solutions by accurate drawing will not be accepted.

The points A and B have coordinates $(-2, 4)$ and $(6, 10)$ respectively.

- (a) Find the equation of the perpendicular bisector of the line AB , giving your answer in the form $ax + by + c = 0$, where a , b and c are integers. [4]

The point C has coordinates $(5, p)$ and lies on the perpendicular bisector of AB .

- (b) Find the value of p . [1]

It is given that the line AB bisects the line CD .

- (c) Find the coordinates of D . [2]

7 $p(x) = ax^3 + 3x^2 + bx - 12$ has a factor of $2x + 1$. When $p(x)$ is divided by $x - 3$ the remainder is 105.

(a) Find the value of a and of b . [5]

(b) Using your values of a and b , write $p(x)$ as a product of $2x + 1$ and a quadratic factor. [2]

(c) Hence solve $p(x) = 0$. [2]

8 In this question all distances are in km.

A ship P sails from a point A , which has position vector $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$, with a speed of 52 kmh^{-1} in the direction of $\begin{pmatrix} -5 \\ 12 \end{pmatrix}$.

(a) Find the velocity vector of the ship. [1]

(b) Write down the position vector of P at a time t hours after leaving A . [1]

At the same time that ship P sails from A , a ship Q sails from a point B , which has position vector $\begin{pmatrix} 12 \\ 8 \end{pmatrix}$, with velocity vector $\begin{pmatrix} -25 \\ 45 \end{pmatrix} \text{ kmh}^{-1}$.

(c) Write down the position vector of Q at a time t hours after leaving B . [1]

(d) Using your answers to **parts (b) and (c)**, find the displacement vector \overrightarrow{PQ} at time t hours. [1]

11

(e) Hence show that $PQ = \sqrt{34t^2 - 168t + 208}$. [2]

(f) Find the value of t when P and Q are first 2 km apart. [2]

12

- 9 (a) (i) Find how many different 4-digit numbers can be formed using the digits 2, 3, 5, 7, 8 and 9, if each digit may be used only once in any number. [1]
- (ii) How many of the numbers found in **part (i)** are divisible by 5? [1]
- (iii) How many of the numbers found in **part (i)** are odd and greater than 7000? [4]

- (b) The number of combinations of n items taken 3 at a time is $92n$. Find the value of the constant n .
[4]

10 (a) Solve $\tan(\alpha + 45^\circ) = -\frac{1}{\sqrt{2}}$ for $0^\circ \leq \alpha \leq 360^\circ$. [3]

(b) (i) Show that $\frac{1}{\sin \theta - 1} - \frac{1}{\sin \theta + 1} = a \sec^2 \theta$, where a is a constant to be found. [3]

(ii) Hence solve $\frac{1}{\sin 3\phi - 1} - \frac{1}{\sin 3\phi + 1} = -8$ for $-\frac{\pi}{3} \leq \phi \leq \frac{\pi}{3}$ radians. [5]

Question 11 is on the next page.

11 Given that $\int_1^a \left(\frac{2}{2x+3} + \frac{3}{3x-1} - \frac{1}{x} \right) dx = \ln 2.4$ and that $a > 1$, find the value of a . [7]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.