

|--|

## **INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the boxes above. Please write clearly and in capital letters.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.

- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Your answers should be supported with appropriate working. Marks may be given for a correct method even if the answer is incorrect.
- Write your answer to each question in the space provided. Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Do **not** write in the bar codes.

# INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- Your quality of written communication is assessed in questions marked with an asterisk (\*).
- The total number of marks for this paper is **60**.
- This document consists of **16** pages. Any blank pages are indicated.



PMT

# Formulae Sheet: Higher Tier













In any triangle ABC Sine rule  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ Cosine rule  $a^2 = b^2 + c^2 - 2bc \cos A$ Area of triangle  $= \frac{1}{2}ab\sin C$ 

**Volume of prism** = (area of cross-section) × length

Volume of sphere =  $\frac{4}{3}\pi r^3$ Surface area of sphere =  $4\pi r^2$ 

Volume of cone =  $\frac{1}{3}\pi r^2 h$ Curved surface area of cone =  $\pi rl$ 

### The Quadratic Equation

The solutions of  $ax^2 + bx + c = 0$ , where  $a \neq 0$ , are given by

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

# PLEASE DO NOT WRITE ON THIS PAGE

# Answer all the questions.

(a) This table gives the number of sunspots observed on 1<sup>st</sup> January each year from 2000 (recorded as 00) to 2014 (recorded as 14).

Year	00	01	02	03	04	05	06	07	08
Number of sunspots	89	136	31	32	32	25	22	7	0
Year	09	10	11	12	13	14			
Number of sunspots	12	33	44	65	87	74			

(i) Plot the last 6 values and complete the time series graph.



- (ii) Describe the trend in the number of sunspots between 2000 and 2014.
  - \_\_\_\_\_ [1]
- (b) This time series graph shows the number of sunspots each year during the last century.



Explain how the graph shows that peaks in sunspot activity occur roughly every 11 years.



2 (a) Solve this inequality.

$$5x-7 \leq 33$$



[2]

3 (a) Helen's DIY store sells brass screws of different sizes. She picks out 4 screws. Their lengths, in inches, are as follows.

5	1	1	3
8	$^{1}\frac{1}{4}$	2	4

Put these lengths in order of size, starting with the smallest.



(b) Many years ago people did not know an exact value for  $\pi$  so approximations were used.

People (Date)	Their approximation to $\pi$
Egyptian (1800 BC)	3.1250
Indian (600 BC)	3.088
Chinese (100 AD)	3.14
Jewish (150 AD)	3.142857
Modern (2016 AD)	3.141 593

(i) Write down any of these values that are larger than the Modern value.

(b)(i) \_\_\_\_\_ [1]

(ii) Work out the difference between the value used in 100 AD and the Modern value. Give your answer correct to 2 significant figures.

(c) The average speed for a journey can be found using this formula.

Average speed = 
$$\frac{\text{distance}}{\text{time}}$$

For each of the following journeys decide if the average speed will be a terminating decimal (T) or a recurring decimal (R).

Write T or R in the final column.

Distance (miles)	Time (hours)	
13	8	
425	24	
5300	450	
14277	1250	

[3]

4 (a) Triangles P and Q are drawn on a coordinate grid.



(i) Use a column vector to describe the translation from P to Q.

		(a)(i)		[2]
	(ii) Reflect triangle <b>P</b> in the line $x = 1$ .			[2]
(b)	Alex rotates a shape. Lizzie enlarges the same shape. They both end up with exactly the same image. Whatever shape they use, their two images are always exactly th	e same.		
	Describe fully their transformations.			
	Alex's rotation		 	
	Lizzie's enlargement		 	[3]

5\* An isosceles triangle has one angle of 126°.The base of this triangle is one of the sides of a regular hexagon.

Calculate the size of the angle marked *m*.



6 The time period, *T* seconds, of a pendulum of length *L* metres is given by this formula.

$$T = 6.28319 \sqrt{\frac{L}{9.81}}$$

A pendulum has length 5.106 metres.

Work out an **estimate** of the time period of this pendulum. Show clearly the estimates you use.

\_\_\_\_\_ seconds [4]

A 6 m long ladder rests against a vertical wall.
The foot of the ladder is 2 m from the wall on horizontal ground.
A horizontal platform of length 0.8 m is attached to the ladder and reaches the wall.

Calculate *x*, the distance along the ladder from the platform to the top of the ladder.



8 The line *L* passes through the points (0, 3) and (5, -7).



(a) Calculate the gradient of line *L*.

(a) \_\_\_\_\_ [2]

**(b)** Write down the equation of line *L*.

(b) \_\_\_\_\_ [2]

(c) Line *M* is perpendicular to line *L* and also passes through the point (0, 3).Write down the equation of line *M*.

Turn over

**9** (a) Solve, algebraically, these simultaneous equations.

$$8x + 6y = 17$$
$$y = -4x - \frac{1}{2}$$



(b) The graph of 8x + 6y = 17 is drawn on the grid.

Draw another graph on the grid and use it to comment on your answer to part (a).



Comment \_\_\_

\_ [3]

**10** In quadrilateral OABC,  $\overrightarrow{OA} = \mathbf{a}$ ,  $\overrightarrow{OB} = \mathbf{b}$  and  $\overrightarrow{OC} = \mathbf{c}$ . M, N, P and Q are the midpoints of the four sides OA, AB, BC and CO.



(a) Find, in terms of **a**, **b** and **c**, an expression for  $\overrightarrow{MQ}$ .

		(a)	['	1]
(b)	Find	l, in terms of <b>a</b> , <b>b</b> and <b>c</b> , an expression for		
	(i)	AB,		
		(b)(i)	[′	1]
	(ii)	BC.		
		(ii)	ſ	11
		('')	L	.,
(c)	Use	your answers to part (b) to show that $\overrightarrow{NP} = \overrightarrow{MQ}$	. ['	1]

- **11** A function is given by  $f(x) = x^{\frac{1}{2}}$ .
  - (a) Work out f(36).
  - (b) Work out  $[f(64)]^{-\frac{1}{3}}$ . Give your answer in its simplest form.

(b) \_\_\_\_\_ [3]

(a) \_\_\_\_\_ [1]

(c) Simplify  $\frac{10}{f(2)}$ . Give your answer in the form  $k\sqrt{2}$  where *k* is an integer.

(c) \_\_\_\_\_ [2]

END OF QUESTION PAPER

## PLEASE DO NOT WRITE ON THIS PAGE



#### **Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series. If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

opportunity.