

1	$y = \frac{7-5x}{2}$ or $y = \frac{7}{2} - \frac{5}{2}x$ or $y = 3.5 - 2.5x$ or $2y = 7 - 5x$ oe		2	M1 for making y or $2y$ the subject Allow $y = -\frac{5}{2}x + c$ oe
		-2.5		A1 for $-\frac{5}{2}$ or -2.5
Total 3 marks				

2	$\left(\frac{-1+5}{2}, \frac{6-4}{2}\right)$ or $\left(\frac{4}{2}, \frac{2}{2}\right)$ or (2, 1)		6	M1 for finding midpoint
	$\frac{-4-6}{5-1}$ or $\frac{6-4}{-1-5}$ or $-\frac{10}{6}$ or $-\frac{5}{3}$			M1 indep for finding the gradient of PQ
	$\frac{-1}{-\frac{10}{6}}$ or $\frac{6}{10}$ or $\frac{-1}{-\frac{5}{3}}$ or $\frac{3}{5}$ or 0.6			M1 for finding the perpendicular gradient to PQ (ft their stated gradient)
	$1 = \frac{3}{5}(2) + c$ or $c = -\frac{1}{5}$ or $c = -\frac{2}{10}$ or $c = -0.2$			M1 dep on 1st and 3rd M1 for substituting '(2, 1)' into $y = \frac{3}{5}x + c$ or find the value of c oe eg $y - 1 = \frac{3}{5}(x - 2)$
	$y = \frac{3}{5}x - \frac{1}{5}$ or $y = 0.6x - 0.2$ or $5y = 3x - 1$			A1 for a correct equation in any form
		$3x - 5y - 1 = 0$		A1 for $3x - 5y - 1 = 0$ or $5y - 3x + 1 = 0$ or $6x - 10y - 2 = 0$ oe accept in the form $ax + by = -c$ eg $3x - 5y = 1$ or $5y - 3x = -1$ oe
Total 6 marks				

Alternative Mark Scheme for Q2

2	$(x+1)^2 + (y-6)^2$ or $(x-5)^2 + (y+4)^2$		6	M1
	$(x+1)^2 + (y-6)^2 = (x-5)^2 + (y+4)^2$			M1 using $PA^2 = QA^2$ (for some point A on the line)
	$x^2 + 2x + 1 + y^2 - 12y + 36$ or $x^2 - 10x + 25 + y^2 + 8y + 16$			M1
	$x^2 + 2x + 1 + y^2 - 12y + 36 = x^2 - 10x + 25 + y^2 + 8y + 16$			M1
	eg $2x + 1 - 12y + 36 = -10x + 25 + 8y + 16$ or $12x + 37 = 20y + 41$			A1 for a correct linear equation in x and y
		$3x - 5y - 1 = 0$		A1 for $3x - 5y - 1 = 0$ oe
Total 6 marks				

3	$y = -\frac{7}{2}x(+10)$ or (gradient $=$) $-\frac{7}{2}$		4	B1 for correct gradient which may be seen in an equation. Condone $-\frac{7}{2}x$
	$'-\frac{7}{2}, m = -1$ or $(m =) \frac{2}{7}$			M1 ft their gradient for use of $m_1 \times m_2 = -1$
	$-11 = \frac{2}{7} \times 6 + c$ or $y - 11 = \frac{2}{7}(x - 6)$ oe			M1 ft dep on M1
		$\left(0, -\frac{89}{7}\right)$		A1 accept $\left(0, -12\frac{5}{7}\right)$ must be exact values
Total 4 marks				

4				M1 for $y = 3x + c$ oe or $y = mx - 2$ oe or $3x - 2$ or eg $L = 3x - 2$ or $y = 3(x \pm a)$
		$y = 3x - 2$	2	A1 oe eg $y - 4 = 3(x - 2)$ $y - 1 = 3(x - 1)$ $y - a = 3(x - b)$ where (a, b) is any coordinate on the line
Total 2 marks				


5	(a)	$y = -4x + k$ (oe)	1	B1 for $y = -4x$ or $y = -4x + k$ where k is any numerical value $k \neq 7$ Could be written in another form e.g. $3y + 12x = 20$
---	-----	--------------------	---	--

6	eg $\frac{4-(-1)}{6-4} (= \frac{5}{2} = 2.5)$		4	M1	for a method to find the gradient of L
	eg $\frac{-1}{2.5}$ ($= -\frac{2}{5} = -0.4$) or $\frac{-1}{\text{their gradient}}$ oe			M1	fit for a method to find the gradient of M if <i>their</i> gradient of L clearly stated (even if no method shown for gradient of L)
	$y = -0.4x + 8$ oe eg $y - 8 = -\frac{2}{5}(x - 0)$ or $(8 \div 2) \times 5 (= 20)$ oe or $8 \div (-\text{their gradient of M})$			M1	dep on previous M1 for substitution of (0, 8) into equation for a line or use of $(8 \div 2) \times 5 (= 20)$ (maybe on diagram) NB: 20 gains M3 if clearly intended as x coordinate (stated or on a diagram)
	<i>Working not required, so correct answer scores full marks (unless from obvious incorrect working)</i>	(20, 0)		A1	
Total 4 marks					

7	$\left(\frac{-1+2}{2}, \frac{5+10}{2}\right)$ or (0.5, 7.5) oe		5	M1	
	$\frac{10-5}{2-(-1)} (= \frac{5}{3})$ oe			M1	
	$m \times \frac{5}{3} = -1$ oe or $m = -\frac{3}{5}$ oe			M1	fit their gradient for use of $m_1 \times m_2 = -1$
	$'7.5' = -\frac{3}{5} \times '0.5' + c$ or $c = 7.8$ oe or $y - '7.5' = -\frac{3}{5}(x - '0.5')$			M1	fit dep on first M1 and third M1
		$5y + 3x = 39$		A1	oe where p, q and r must be integers
Total 5 marks					

8	$3 \div 2 (= 1.5 \text{ or } \frac{3}{2})$ or eg $\frac{5-1}{4-(-0)}$ or $c = -1$		3	M1	for correct method to find gradient or the correct value of c for gradient, may see a correct calculation or $\frac{3}{2}$ oe or $1.5x (+c)$ oe for value of c , allow $c = -1, y = -1, (L =) mx - 1$ oe
	$y = "1.5"x (+c)$ or $y = mx - 1$ or eg $y - 5 = m(x - 4)$			M1	for use of $y = mx + c$ with either m or c correct (NB: $m \neq 0$) or for $(L =) 1.5x - 1$ oe
		$y = \frac{3}{2}x - 1$		A1	oe eg $y = 1.5x - 1$
Total 3 marks					

9	(gradient $AM =$) $\frac{4-2}{-3-0}$ oe ($= -\frac{2}{3}$)		7	M1	A correct method to find gradient of AM
	$y = \frac{3}{2}x + 2$ or eg $\frac{y-2}{x} = \frac{3}{2}$ oe			M1	For the correct equation of the line passing through BD or for a correct expression involving the x and y coordinates of point B or point D
	$(x-(-3))^2 + (y-4)^2 = 6.5^2$ or $(x-0)^2 + (y-2)^2 = 6.5^2 - [(-3-0)^2 + (4-2)^2]$ oe eg $x^2 + (y-2)^2 = 29.25$			M1	A correct equation in x and y to find the coordinates of B and D
	eg $x^2 + 6x + 9 + y^2 - 8y + 16 - 42.25 = 0$ oe or $x^2 + y^2 - 4y + 4 - 29.25 = 0$ oe			M1	Brackets expanded
	eg $x^2 + 6x + 9 + \left(\frac{3}{2}x + 2\right)^2 - 8\left(\frac{3}{2}x + 2\right) + 16 - 42.25 = 0$ $\left(\frac{2y-4}{3}\right)^2 + y^2 - 4y + 4 - 29.25 = 0$ oe			M1	For a correct substitution into a correct equation to get an equation in either x only or y only
	eg $\frac{13}{4}x^2 = \frac{117}{4}$ or $13y^2 - 52y - 211.25 = 0$ oe			M1	A fully correct simplified equation in x or in y - all brackets expanded and like terms grouped.
		(3, 6.5) (-3, -2.5)		A1	correct coordinates SCB3 for one pair of correct coordinates or both x values correct or both y values correct
Total 7 marks					

9 Alt 1	$(AM = \sqrt{3^2 + 2^2} = \sqrt{13} = 3.605\dots)$ or $(AM^2 = 3^2 + 2^2 = 13)$		7	M1	Use of Pythagoras for point A to point M
	$(BM = \sqrt{6.5^2 - \sqrt{13}^2} = \sqrt{29.25} = \frac{3\sqrt{13}}{2} = 5.4083\dots)$			M1	A correct method to find the length of BM or DM
	$(SF = \frac{\sqrt{29.25}}{\sqrt{13}} = \frac{3}{2}$ oe or $MN = x, BN = 1.5x$ (see diag) or $(LAM = \sin^{-1} \frac{3}{\sqrt{13}} (= 56.3\dots))$ oe or $(LMA = \cos^{-1} \frac{3}{\sqrt{13}} (= 33.6\dots))$ or			M1	A correct method to find the SF of the enlargement of the sides AM to BM or angle LAM OR LMA 
	eg $\overline{MB}_x = \frac{3}{2} \times 2$ or $\overline{MB}_y = \frac{3}{2} \times 3$ or $\overline{MD}_x = -\frac{3}{2} \times 2$ or $\overline{MD}_y = -\frac{3}{2} \times 3$ oe or $x^2 + (1.5x)^2 = \sqrt{29.25}^2$ or $MN = \sqrt{29.25} \cos 56.3\dots (= 3)$ oe or $BN = \sqrt{29.25} \sin 56.3\dots (= 4.5)$ oe <i>turn over</i>			M1	A correct method to find the translation of at least one component of MB or MD (need not be written in vector form) OR correct Pythagoras statement using the SF to find x coordinates OR 1 correct trig statement to find translations from M
	$\overline{MB}_x = \frac{3}{2} \times 2$ and $\overline{MB}_y = \frac{3}{2} \times 3$ or $\overline{MD}_x = -\frac{3}{2} \times 2$ and $\overline{MD}_y = -\frac{3}{2} \times 3$ oe or $x^2 + 2.25x^2 = 29.25$ or $MN = \frac{3\sqrt{13}}{2} \cos 56.309\dots (= 3)$ and $BN = \frac{3\sqrt{13}}{2} \sin 56.309\dots (= 4.5)$ oe			M1	A correct method to find the translation of both components of MB or MD (need not be written in vector form) OR correct Pythagoras statement with no brackets using the SF to find x coordinates OR 2 correct trig statements to find translations from M

	eg $(0, 2)$ is translated $\begin{pmatrix} 3 \\ 4.5 \end{pmatrix}$ or $(0+3, 2+4.5) (= (3, 6.5))$ or $(0, 2)$ is translated $\begin{pmatrix} -3 \\ -4.5 \end{pmatrix}$ or $(0-3, 2-4.5) (= (-3, -2.5))$ oe or $3.25x^2 = 29.25$			M1	correct method to find the coordinates of B or D or one pair of correct coordinates or a correct method to find both x coordinates or both y coordinates OR a fully correct simplified equation in x all brackets expanded and like terms grouped.
		$(3, 6.5)$ $(-3, -2.5)$		A1	correct coordinates SCB3 for one correct coordinate or both x values correct or both y values correct
Total 7 marks					

10 (a)		$y = -3x + 5$ oe	2	B2	fully correct equation eg $y = -3x + 5$ or $y - 5 = -3(x - 0)$ If not B2 then B1 for $y = -3x + a$ with $a \neq 5$ or $y = bx + 5$ ($b \neq 0, -3$) or $(L =) -3x + 5$
--------	--	---------------------	---	----	---

11	$\left(\frac{dy}{dx} = \right) 16x - 14$		5	M1	Differentiation to obtain 2 terms with at least 1 correct
	$16x - 14 = 10$			M1	their $dy/dx = 10$ dep on M1
	$(1.5, -9)$ or $x = 1.5, y = -9$			A1	coordinates of point on curve at which gradient is 10 – allow given as coordinates or as x worked out and y worked out if meaning is clear
	eg $y - -9 = -\frac{1}{10}(x - \frac{3}{2})$ oe or eg $-9 = -\frac{1}{10} \times 1.5 + c$ oe			M1	A correct method to find the equation for line Q using $(1.5, -9)$
	Correct answer scores full marks (unless from obvious incorrect working)	$2x + 20y + 177 = 0$		A1	oe where a, b, c are integers eg $10x + 100y + 885 = 0$
Total 5 marks					

12	$y = \frac{3}{5}x \left(+ \frac{6}{5} \right)$ or $y = 0.6x (+1.2)$ or (gradient =) $\frac{3}{5}$ or 0.6 oe		6	M1 for correct gradient which may be seen in an equation. Condone $\frac{3}{5}x$ or $0.6x$
	$k = -2$			B1 for $k = -2$
	$\left(\frac{-2+8}{2}, \frac{1+7}{2} \right)$ oe or (3, 4)			M1 for finding the midpoint (use of their k where $k < 0$)
	$\frac{3}{5}m = -1$ or $(m =) -\frac{5}{3}$			M1 ft their gradient for use of $m_1 \times m_2 = -1$ Allow $-\frac{5}{3} = -1.67$ or better
	$"4" = -\frac{5}{3} \times "3" + c$ or $c = 9$ or $y - "4" = -\frac{5}{3} ("x - "3")$			M1 dep on M3
	<i>Working required</i>	$5x + 3y = 27$		A1 allow equation in any form where p, q and r are integers
Total 6 marks				

13	(d) $y = mx + 4$ where $m \neq 0$ oe (eg $y = 2x + 4$) or $y = -2x + c$ or $y + 2x = c$ oe or $-2x + 4$ or $f(x) = -2x + 4$ oe		2	M1
	<i>Correct answer scores full marks (unless from obvious incorrect working)</i>	$y = -2x + 4$		A1 oe eg $y + 2x = 4$

14	eg (gradient =) $\frac{12 - -48}{-5 - 19} (= -2.5)$ oe		3	M1 for a method to find the gradient
	eg $12 = "-2.5" \times -5 + c$ oe $y - 12 = "-2.5"(x - -5)$ oe			M1 ft their gradient
	<i>Correct answer scores full marks (unless from obvious incorrect working)</i>	$y = -2.5x - 0.5$		A1 oe eg $y - 12 = -2.5(x + 5)$ or $2y + 5x + 1 = 0$
Total 3 marks				

15	$\frac{-5-10}{4-2} (= -\frac{5}{2})$		6	M1 A correct method to find the gradient of AC
	$y - 10 = -\frac{5}{2}(x + 2)$ oe eg $y = -\frac{5}{2}x + 5$ or $y - -5 = -\frac{5}{2}(x - 4)$ oe or $5x + 2y = 10$ oe			M1 ft (if M1 scored) correct equation of AC
	$y - 4 = \frac{2}{5}(x - -\frac{27}{5})$ oe or $4 = \frac{2}{5}(-\frac{27}{5}) + c$ ($y = \frac{2}{5}x + 6.16$) $\frac{4-y}{-\frac{27}{5}-x} = \frac{2}{5}$ oe or $5y - 2x = \frac{154}{5}$ oe			M1 ft (if first M1 scored) equation of BD or correct equation using gradient of BD
	solves $-\frac{5}{2}x + 5 = \frac{2}{5}x + 6.16$ oe eg $10x + 4y = 20$ oe, with operation of addition or $-10x + 25y = 154$ oe, with operation of subtraction or $25x + 10y = 50$ oe, with operation of subtraction or $-4x + 10y = 61.6$ oe, with operation of subtraction or $x = \frac{5}{2}y - \frac{154}{10}$ oe or $y = \frac{2}{5}x + \frac{154}{25}$ oe substituted in other equation			M1 Solve equation OR Solve simultaneously the correct equations of lines of AC and BD or correct equation from gradient or other correct equation. If elimination: same coefficient of x or y with suitable sign used to eliminate. If substitution: x or y substituted into other equation.
	Coordinates of intersection of AC and BD : $x = -\frac{2}{5}, y = 6$			M1 oe value of x and y at intersection of AC and BD
	<i>Correct answer scores full marks (unless from obvious incorrect working)</i>	(4.6, 8)		A1 oe coordinates of D
See next page for working with $AD = AB, CD = CB$ or gradients				
Total 6 marks				

15	$\text{eg } (10-4)^2 + \left(-2 + \frac{27}{5}\right)^2 (= 47.56) \text{ } (AB = 6.896\dots) \text{ or}$ $\text{eg } (-5-4)^2 + \left(4 + \frac{27}{5}\right)^2 (= 169.36) \text{ } (CB = 13.013\dots) \text{ or}$ $\text{eg } \frac{-5-10}{4-2} \text{ or } \frac{4-y}{-\frac{27}{5}-x} \text{ oe}$		6	M1 A correct method to find AB^2 or CB^2 or AB or CB or a correct gradient expression for AC or DB
	$\text{eg } (y-10)^2 + (x+2)^2 = (10-4)^2 + \left(-2 + \frac{27}{5}\right)^2 \text{ or}$ $\text{eg } (y+5)^2 + (x-4)^2 = (-5-4)^2 + \left(4 + \frac{27}{5}\right)^2 \text{ or}$ $\frac{-5-10}{4-2} \times \frac{4-y}{-\frac{27}{5}-x} = -1 \text{ oe eg } -60 + 15y = 6x + 32.4$			M1 Using $D(x, y)$ form a correct equation $AD^2 = AB^2$ or $CD^2 = CB^2$ or gradients $AC \times DB = -1$ (Using $D(x, y)$)
	$\text{eg } 2x - 5y = -30.8 \text{ or } x = 2.5y - 15.4 \text{ or } y = 0.4x + 6.16 \text{ oe}$			M1 uses rearrangement or solving simultaneous equations to find a correct 3 term linear equation
	$\text{eg } (y-10)^2 + (2.5y-15.4+2)^2 = (10-4)^2 + \left(-2 + \frac{27}{5}\right)^2$ $\text{eg } (0.4x+6.16+5)^2 + (x-4)^2 = (-5-4)^2 + \left(4 + \frac{27}{5}\right)^2$			M1 uses substitution to obtain a correct quadratic equation in one unknown
	$7.25y^2 - 87y + 232 = 0 \text{ oe or } 1.16x^2 + 0.928x - 28.8144 = 0 \text{ oe}$			M1 for a 3 term quadratic that can be used to find the value of x or the value of y at D
		(4.6, 8)		A1 oe coordinates of D
Total 6 marks				