


Q1.

Diagram **NOT** accurately drawn

The sketch shows a curve with equation

$$y = ka^x$$

where k and a are constants, and $a > 0$

The curve passes through the points (1, 7) and (3, 175).

Calculate the value of k and the value of a .

$$k = \dots\dots\dots$$

$$a = \dots\dots\dots$$

(Total 3 marks)

M1.

Working	Answer	Mark	Additional Guidance
$7 = ka^1; 175 = ka^3$ $k = \frac{7}{a}, 175 = \frac{7a^3}{a}, 175 = 7a^2$ $a^2 = 25, \text{ so } a = 5, k = 1.4$ Or $7^3 = k^3a^3, 175 = ka^3$ $k^3 = \frac{7}{175}, k = 1.4, a = 5$	$k = 1.4$ $a = 5$	3	M1 either $a^2 = 25$ or $7 = ka$ (or $7 = ka^1$) and $175 = ka^3$ A1 $k = 1.4$ oe A1 $a = 5$ SC Either $a = 5$ or $k = 1.4$ oe gets B2
			Total for Question: 3 marks

- E1.** Exponential growth is generally found to be a hard topic at GCSE and this question was no different. Many candidates started sensibly and substituted the values of x and y to get the pair of equations $7 = ka^1$ and $175 = ka^3$. However, things then went badly wrong, mainly through poor use of index laws. For example, was evaluated as 1, leading to $k = 7$ and ignoring the second equation, or, the 2 equations were combined to eliminate k giving $a^2 = 175/7$, followed by a cube root. There appeared to be little evidence of candidates checking the value of a and the value of k in both equations.