

1. Simplify the following expressions fully.

(a)  $(x^6)^{\frac{1}{3}}$

$$(x^a)^b = x^{ab}$$

(1)

(b)  $\sqrt{2}(x^3) \div \sqrt{\frac{32}{x^2}}$

(2)

a)  $(x^6)^{\frac{1}{3}} = x^{6/3} = x^2$

b)  $\sqrt{2} x^3 \div \sqrt{\frac{32}{x^2}} = \sqrt{2} x^3 \times \frac{\sqrt{x^2}}{\sqrt{32}}$

$$= \sqrt{2} x^3 \times \frac{x}{\sqrt{16 \times 2}}$$

$$= \sqrt{2} x^3 \times \frac{x}{4\sqrt{2}}$$

$$= \frac{x^4}{4}$$

2.




Figure 1

Figure 1 shows a sketch of part of the graph of  $y = \frac{12}{\sqrt{x^2 - 2}}$ ,  $x \geq 2$

The table below gives values of  $y$  rounded to 3 decimal places.

$x$	2	5	8	11
$y$	8.485	2.502	1.524	1.100

- (a) Use the trapezium rule with all the values of  $y$  from the table to find an approximate value, to 2 decimal places, for

$$\int_2^{11} \frac{12}{\sqrt{x^2 - 2}} dx$$

$$h = 5 - 2 = 3$$

(4)

- (b) Use your answer to part (a) to estimate a value for

$$\int_2^{11} \left( 1 + \frac{6}{\sqrt{x^2 - 2}} \right) dx$$

(3)

$$\begin{aligned} \text{a) } \int_2^{11} \frac{12}{\sqrt{x^2 - 2}} dx &= \frac{1}{2} h [y_0 + y_3 + 2(y_1 + y_2)] \\ &= \frac{1}{2} (3) [8.485 + 1.1 + 2(2.502 + 1.524)] \\ &= 26.46 \text{ (2 dp)} \end{aligned}$$

$$\begin{aligned} \text{b) } \int_2^{11} \left( 1 + \frac{6}{\sqrt{x^2 - 2}} \right) dx &= \int_2^{11} 1 dx + \frac{1}{2} \int_2^{11} \frac{12}{\sqrt{x^2 - 2}} dx \\ &= [x]_2^{11} + \frac{1}{2} (26.46) = 9 + 13.23 = 22.23 \text{ (2 dp)} \end{aligned}$$

3.




Figure 2


Figure 2 shows a sketch of part of the curve with equation  $y = f(x)$ .  
The curve crosses the coordinate axes at the points  $(2.5, 0)$  and  $(0, 9)$ , has a stationary point at  $(1, 11)$ , and has an asymptote  $y = 3$

On **separate** diagrams, sketch the curve with equation

(a)  $y = 3f(x)$  (3)

(b)  $y = f(-x)$  (3)

On each diagram show clearly the coordinates of the points of intersection of the curve with the two coordinate axes, the coordinates of the stationary point, and the equation of the asymptote.



4. (a) Find the first 4 terms in ascending powers of  $x$  of the binomial expansion of

$$\left(2 + \frac{x}{4}\right)^{10}$$

giving each term in its simplest form.

(4)

- (b) Use your expansion to find an estimated value for  $2.025^{10}$ , stating the value of  $x$  which you have used and showing your working.

$$\begin{aligned} \text{a) } \left(2 + \frac{x}{4}\right)^{10} &\approx 2^{10} + {}^{10}C_1 2^9 \left(\frac{x}{4}\right)^1 + {}^{10}C_2 2^8 \left(\frac{x}{4}\right)^2 \\ &\quad + {}^{10}C_3 2^7 \left(\frac{x}{4}\right)^3 \end{aligned} \quad (3)$$

$$= 1024 + 1280x + 720x^2$$

$$+ 240x^3$$

$$\text{b) } \text{When } x = 0.1,$$

$$\left(2 + \frac{x}{4}\right)^{10} = 2.025^{10}$$

$$= 1024 + 1280(0.1) + 720(0.1)^2$$

$$+ 240(0.1)^3$$

$$= 1159.44$$

5. (a) Prove that the sum of the first  $n$  terms of an arithmetic series is given by the formula

$$S_n = \frac{n}{2}[2a + (n-1)d]$$

where  $a$  is the first term of the series and  $d$  is the common difference between the terms.

(4)

- (b) Find the sum of the integers which are divisible by 7 and lie between 1 and 500

(3)

$$a) \quad S_n = a + (a+d) + (a+2d) + \dots + [a+(n-2)d] + [a+(n-1)d]$$

$$S_n = [a+(n-1)d] + [a+(n-2)d] + \dots + (a+2d) + (a+d) + a$$

$$2S_n = [a+a+(n-1)d] + [a+d+a+(n-2)d] + \dots + [a+a+(n-1)d]$$

$$= [2a+(n-1)d]n$$

$$S_n = \frac{n}{2}[2a+(n-1)d]$$

$$b) \quad S = 7 + 14 + \dots + 497$$

$$a = 7, \quad l = 497, \quad d = 7, \quad n = \frac{497}{7} = 71$$

$$\Rightarrow S = \frac{71}{2}(7+497)$$

$$S = \frac{n}{2}(a+l)$$

$$= 17892$$

6. Given that

$$2 \log_4(2x + 3) = 1 + \log_4 x + \log_4(2x - 1), \quad x > \frac{1}{2}$$

(a) show that

$$4x^2 - 16x - 9 = 0 \tag{5}$$

(b) Hence solve the equation

$$2 \log_4(2x + 3) = 1 + \log_4 x + \log_4(2x - 1), \quad x > \frac{1}{2} \tag{2}$$

a)  $2 \log_4(2x+3) = 1 + \log_4 x + \log_4(2x-1)$

$$\log_4(2x+3)^2 - \log_4 x - \log_4(2x-1) = 1$$

$$\log_4 \left[ \frac{(2x+3)^2}{x(2x-1)} \right] = 1$$

$$\frac{(2x+3)^2}{x(2x-1)} = 4^1$$

$$(2x+3)^2 = 4x(2x-1)$$

$$4x^2 + 12x + 9 = 8x^2 - 4x$$

$$4x^2 - 16x - 9 = 0$$

b)  $4x^2 - 16x - 9 = 0$

$$(2x+1)(2x-9) = 0$$

$$x = -\frac{1}{2} \text{ or } \frac{9}{2}$$

But given  $x > \frac{1}{2}$ ,  $x = \frac{9}{2}$

$$\begin{aligned} a \log b &= \log b^a \\ \log a + \log b &= \log(ab) \\ \log a - \log b &= \log\left(\frac{a}{b}\right) \end{aligned}$$

7. The circle  $C$  has equation

$$x^2 + y^2 + 10x - 6y + 18 = 0$$

Find

(a) the coordinates of the centre of  $C$ , (2)

(b) the radius of  $C$ . (2)

The circle  $C$  meets the line with equation  $x = -3$  at two points.

(c) Find the exact values for the  $y$  coordinates of these two points, giving your answers as fully simplified surds. (4)

$$x^2 + y^2 + 10x - 6y + 18 = 0$$

$$(x+5)^2 - 25 + (y-3)^2 - 9 + 18 = 0$$

$$(x+5)^2 + (y-3)^2 = 16$$

a) Centre:  $(-5, 3)$

b) Radius =  $\sqrt{16} = 4$

$$(x-a)^2 + (y-b)^2 = r^2$$

c) When  $x = -3$ ,

$$(-3+5)^2 + (y-3)^2 = 16$$

$$(y-3)^2 = 12$$

$$y-3 = \pm\sqrt{12}$$

$$y = 3 \pm \sqrt{12}$$

$$= 3 \pm 2\sqrt{3}$$

8. A sequence is defined by

$$\begin{aligned} u_1 &= k \\ u_{n+1} &= 3u_n - 12, \quad n \geq 1 \end{aligned}$$

where  $k$  is a constant.

(a) Write down fully simplified expressions for  $u_2$ ,  $u_3$  and  $u_4$  in terms of  $k$ .

(4)

Given that  $u_4 = 15$

(b) find the value of  $k$ ,

(2)

(c) find  $\sum_{i=1}^4 u_i$ , giving an exact numerical answer.

(3)

a)  $u_1 = k$   $u_2 = 3u_1 - 12 = 3k - 12$   $u_3 = 3u_2 - 12 = 3(3k - 12) - 12$

$u_2 = 3u_1 - 12$   $= 9k - 48$

$= 3k - 12$   $u_4 = 3u_3 - 12 = 3(9k - 48) - 12$

$= 27k - 156$

b)  $15 = 27k - 156$

$171 = 27k$

$k = \frac{19}{3}$

c)  $\sum_{i=1}^4 u_i = u_1 + u_2 + u_3 + u_4$

$= k + 3k - 12 + 9k - 48 + 27k - 156$

$= 40k - 216$

$= 40\left(\frac{19}{3}\right) - 216$

$= 112/3$



9.




Figure 3

In Figure 3, the points  $A$  and  $B$  are the centres of the circles  $C_1$  and  $C_2$  respectively. The circle  $C_1$  has radius 10 cm and the circle  $C_2$  has radius 5 cm. The circles intersect at the points  $X$  and  $Y$ , as shown in the figure.

Given that the distance between the centres of the circles is 12 cm,

- (a) calculate the size of the acute angle  $XAB$ , giving your answer in radians to 3 significant figures, (2)
- (b) find the area of the major sector of circle  $C_1$ , shown shaded in Figure 3, (3)
- (c) find the area of the kite  $AYBX$ . (3)

$$a) \quad \cos \widehat{XAB} = \frac{10^2 + 12^2 - 5^2}{2 \times 10 \times 12} \quad \text{cosine rule}$$

$$\widehat{XAB} = 0.421 \text{ (3 sf)}$$

$$b) \quad \widehat{XAY} = 2 \widehat{XAB} = 2 \times 0.421 = 0.843^\circ$$

$$\begin{aligned} \text{Area of major sector} &= \frac{1}{2} (10)^2 (2\pi - 0.843) \\ &= 272 \text{ cm}^2 \end{aligned} \quad A = \frac{1}{2} r^2 \theta$$

$$c) \quad \text{Area of } \triangle XAB = \frac{1}{2} \times 10 \times 12 \sin(0.421) = 24.5$$

$$\text{Area of kite} = 2 \times 24.5$$

$$= 49.0 \text{ cm}^2$$

$$A = \frac{1}{2} ab \sin \theta$$

10.

$$f(x) = 6x^3 + ax^2 + bx - 5$$

where  $a$  and  $b$  are constants.

When  $f(x)$  is divided by  $(x + 1)$  there is no remainder.

When  $f(x)$  is divided by  $(2x - 1)$  the remainder is  $-15$

(a) Find the value of  $a$  and the value of  $b$ .

(5)

(b) Factorise  $f(x)$  completely.

(4)

$$a) \quad f(-1) = 6(-1)^3 + a(-1)^2 + b(-1) - 5 = 0$$

$$-6 + a - b - 5 = 0$$

$$a - b = 11 \quad (1)$$

$$f\left(\frac{1}{2}\right) = 6\left(\frac{1}{2}\right)^3 + a\left(\frac{1}{2}\right)^2 + b\left(\frac{1}{2}\right) - 5 = -15$$

$$\frac{3}{4} + \frac{a}{4} + \frac{b}{2} - 5 = -15$$

$$a + 2b = -43 \quad (2)$$

$$(1) - (2): -3b = 54$$

$$b = -18$$

$$\ln (1): a = 11 + (-18) = -7$$

$$b) \quad f(x) = 6x^3 - 7x^2 - 18x - 5$$

$$= (x+1)(6x^2 - 13x - 5) \quad \text{by inspection}$$

$$= (x+1)(3x+1)(2x-5)$$

Remainder  
theorem

11.




Figure 4

Figure 4 shows a sketch of the curve  $C$  with equation  $y = \sin(x - 60^\circ)$ ,  $-360^\circ \leq x \leq 360^\circ$

(a) Write down the exact coordinates of the points at which  $C$  meets the two coordinate axes.

(3)

(b) Solve, for  $-360^\circ \leq x \leq 360^\circ$ ,

$$4 \sin(x - 60^\circ) = \sqrt{6} - \sqrt{2}$$

showing each stage of your working.

(5)

a) Meets the x-axis at:  $(-300, 0)$ ;  $(-120, 0)$ ;  
 $(60, 0)$ ;  $(240, 0)$

Transform  $y = \sin x$   
When  $x = 0$ ,  $y = \sin(-60) = -\sin 60 = \frac{-\sqrt{3}}{2}$

$\therefore$  Meets the y-axis at  $(0, -\sqrt{3}/2)$

b)  $4 \sin(x - 60) = \sqrt{6} - \sqrt{2}$

$$\sin(x - 60) = \frac{\sqrt{6} - \sqrt{2}}{4}$$

$$x - 60 = -345, -195, 15, 165$$

$$x = -285^\circ, -135^\circ, 75^\circ, 225^\circ$$



12. A business is expected to have a yearly profit of £275 000 for the year 2016. The profit is expected to increase by 10% per year, so that the expected yearly profits form a geometric sequence with common ratio 1.1

(a) Show that the difference between the expected profit for the year 2020 and the expected profit for the year 2021 is £40 300 to the nearest hundred pounds. (3)

(b) Find the first year for which the expected yearly profit is more than one million pounds. (4)

(c) Find the total expected profits for the years 2016 to 2026 inclusive, giving your answer to the nearest hundred pounds. (3)

$$a) \quad u_n = ar^{(n-1)}$$

$$u_5 = 275000 \times 1.1^4 = 402628$$

$$u_6 = 275000 \times 1.1^5 = 442890$$

$$u_6 - u_5 = 442890 - 402628 = 40300$$

$$b) \quad 275000 \times 1.1^{(n-1)} = 1000000$$

$$1.1^{(n-1)} = 3.64$$

$$n-1 = \frac{\ln 3.64}{\ln 1.1} = 13.6$$

$$\therefore \text{Year is } 2016 + 14 = 2030$$

$$c) \quad S_n = \frac{a(r^n - 1)}{r - 1}$$

$$S_{11} = \frac{275000(1.1^{11} - 1)}{1.1 - 1}$$

$$= 5096100$$

13. The curve  $C$  has equation

$$y = 3x^2 - 4x + 2$$

The line  $l_1$  is the normal to the curve  $C$  at the point  $P(1, 1)$

(a) Show that  $l_1$  has equation

$$x + 2y - 3 = 0 \quad (5)$$

The line  $l_1$  meets curve  $C$  again at the point  $Q$ .

(b) By solving simultaneous equations, determine the coordinates of the point  $Q$ . (4)

Another line  $l_2$  has equation  $kx + 2y - 3 = 0$ , where  $k$  is a constant.

(c) Show that the line  $l_2$  meets the curve  $C$  once only when

$$k^2 - 16k + 40 = 0 \quad (4)$$

(d) Find the two exact values of  $k$  for which  $l_2$  is a tangent to  $C$ . (2)

a)  $y = 3x^2 - 4x + 2$

$$\frac{dy}{dx} = 6x - 4$$

$$\text{At } (1, 1), \frac{dy}{dx} = 6(1) - 4 = 2$$

$$\therefore \text{Grad of } l_1 \text{ is } -\frac{1}{2}$$

$$y - y_1 = m(x - x_1)$$

$$y - 1 = -\frac{1}{2}(x - 1)$$

$$2y - 2 = -x + 1$$

$$x + 2y - 3 = 0$$

Question 13 continued

$$b) \quad y = 3x^2 - 4x + 2 \quad (1) \quad x + 2y - 3 = 0 \quad (2)$$

$$(1) \text{ in } (2): x + 6x^2 - 8x + 4 - 3 = 0$$

$$6x^2 - 7x + 1 = 0$$

$$(6x - 1)(x - 1) = 0$$

$$x = 1 \text{ or } \frac{1}{6}$$

$$\begin{aligned} \text{When } x = \frac{1}{6}, y &= 3\left(\frac{1}{6}\right)^2 - 4\left(\frac{1}{6}\right) + 2 \\ &= \frac{1}{12} - \frac{2}{3} + 2 \\ &= \frac{17}{12} \end{aligned}$$

$$\therefore Q\left(\frac{1}{6}, \frac{17}{12}\right)$$

$$c) \quad kx + 2y - 3 = 0 \quad (3)$$

$$(1) \text{ in } (3): kx + 6x^2 - 8x + 4 - 3 = 0$$

$$6x^2 + (k - 8)x + 1 = 0$$

For one solution,  $\Delta = 0$

$$(k - 8)^2 - 4(6)(1) = 0$$

$$k^2 - 16k + 40 = 0$$

$$d) \quad k = \frac{16 \pm \sqrt{16^2 - 4(40)}}{2} = 8 \pm \frac{\sqrt{96}}{2} = 8 \pm$$

$$= 8 \pm \frac{1}{2} \sqrt{16 \times 6} = 8 \pm 2\sqrt{6}$$

14. In this question, solutions based entirely on graphical or numerical methods are not acceptable.

(i) Solve, for  $0 \leq x < 360^\circ$ ,

$$3 \sin x + 7 \cos x = 0$$

Give each solution, in degrees, to one decimal place.

(4)

(ii) Solve, for  $0 \leq \theta < 2\pi$ ,

$$10 \cos^2 \theta + \cos \theta = 11 \sin^2 \theta - 9$$

Give each solution, in radians, to 3 significant figures.


(6)

i)

$$3 \sin x = -7 \cos x$$

$$\tan x = -\frac{7}{3}$$

$$x = 113.2^\circ, 293.2^\circ$$



ii)



$$10 \cos^2 \theta + \cos \theta = 11(1 - \cos^2 \theta) - 9$$

$$= 11 - 11 \cos^2 \theta - 9$$

$$21 \cos^2 \theta + \cos \theta - 2 = 0$$

$$(7 \cos \theta - 2)(3 \cos \theta + 1)$$

$$\cos \theta = \frac{2}{7} \text{ or } -\frac{1}{3}$$



$$\theta = 1.28, 1.91, 4.37, 5.00 \text{ (3sf)}$$

15.




Figure 5

Figure 5 shows a sketch of part of the curve  $C$  with equation

$$y = x^3 + 10x^{\frac{3}{2}} + kx, \quad x \geq 0$$

where  $k$  is a constant.

- (a) Find  $\frac{dy}{dx}$  (2)

The point  $P$  on the curve  $C$  is a minimum turning point.  
Given that the  $x$  coordinate of  $P$  is 4

- (b) show that  $k = -78$  (2)

The line through  $P$  parallel to the  $x$ -axis cuts the  $y$ -axis at the point  $N$ .

The finite region  $R$ , shown shaded in Figure 5, is bounded by  $C$ , the  $y$ -axis and  $PN$ .

- (c) Use integration to find the area of  $R$ . (7)

a)  $y = x^3 + 10x^{\frac{3}{2}} + kx$

$$\frac{dy}{dx} = 3x^2 + 15x^{\frac{1}{2}} + k$$

b) When  $x = 4$ ,  $\frac{dy}{dx} = 3(4)^2 + 15(4)^{\frac{1}{2}} + k = 0$

$$48 + 30 + k = 0$$

$$k = -78$$



Question 15 continued

$$\begin{aligned} \text{c) When } x=4, y &= (4)^3 + 10(4)^{3/2} - 78(4) \\ &= 64 + 80 - 312 \\ &= -168 \end{aligned}$$

$$\therefore |0N| = 168$$

$$R + Q = 168 \times 4 = 672$$

$$-Q = \int_0^4 (x^3 + 10x^{3/2} - 78x) dx$$

$$= \left[ \frac{x^4}{4} + 4x^{5/2} - 39x^2 \right]_0^4$$

$$= \left[ \frac{(4)^4}{4} + 4(4)^{5/2} - 39(4)^2 \right] - 0$$

$$= 64 + 128 - 624$$

$$Q = 432$$

$$\therefore R = 672 - 432$$

$$= 240$$