

1. (a) Express  $\frac{2}{4r^2 - 1}$  in partial fractions.

(2)

(b) Hence use the method of differences to show that

$$\sum_{r=1}^n \frac{1}{4r^2 - 1} = \frac{n}{2n+1} \quad (3)$$

a)  $\frac{2}{(2r-1)(2r+1)} = \frac{A}{2r-1} + \frac{B}{2r+1} \Rightarrow 2 = A(2r+1) + B(2r-1)$   
 $r = \frac{1}{2} \Rightarrow A = 1 \quad r = -\frac{1}{2} \quad B = -1$

$$\therefore = \frac{1}{2r-1} - \frac{1}{2r+1}$$

b)  $\sum_{r=1}^n \frac{2}{4r^2-1} = \left(\frac{1}{1} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{5}\right) + \left(\frac{1}{5} - \frac{1}{7}\right) + \dots + \left(\frac{1}{2n-3} - \frac{1}{2n-1}\right) + \left(\frac{1}{2n-1} - \frac{1}{2n+1}\right)$   
 $= 1 - \frac{1}{2n+1} = \frac{2n+1-1}{2n+1} = \frac{2n}{2n+1}$

$$\therefore 2 \sum_{r=1}^n \frac{1}{4r^2-1} = \frac{2n}{2n+1} \quad \therefore \sum_{r=1}^n \frac{1}{4r^2-1} = \frac{n}{2n+1} \quad \text{**}$$

2. Using algebra, find the set of values of  $x$  for which


$$3x - 5 < \frac{2}{x}$$

(5)

$$3x - 5 - \frac{2}{x} < 0 \Rightarrow \frac{3x^2 - 5x - 2}{x} < 0$$

$$\Rightarrow \frac{(3x + 1)(x - 2)}{x} < 0$$

$$\therefore x < -\frac{1}{3} \text{ or } 0 < x < 2$$



3. (a) Find the general solution of the differential equation

$$\frac{dy}{dx} + 2y \tan x = e^{4x} \cos^2 x, \quad -\frac{\pi}{2} < x < \frac{\pi}{2}$$

giving your answer in the form  $y = f(x)$ .

(6)

(b) Find the particular solution for which  $y = 1$  at  $x = 0$

(2)

$$I.F = e^{\int \tan x dx} = (e^{\ln(\sec x)})^2 = \sec^2 x$$

$$\Rightarrow \sec^2 x \frac{dy}{dx} + 2y \tan x \sec^2 x = e^{4x} \cos^2 x \sec^2 x$$

$$\therefore \frac{d}{dx}(y \sec^2 x) = e^{4x} \Rightarrow y \sec^2 x = \int e^{4x} dx = \frac{1}{4} e^{4x} + C$$

$$\therefore y = \frac{1}{4} e^{4x} \cos^2 x + C \cos^2 x$$

$$x=0, y=1 \quad 1 = \frac{1}{4}(1)(1)^2 + C(1)^2$$

$$1 = \frac{1}{4} + C \quad \therefore C = \frac{3}{4}$$

$$\therefore y = \frac{1}{4} \cos^2 x (e^{4x} + 3)$$

4.




Figure 1

Figure 1 shows the curve  $C$  with polar equation

$$r = 2 \cos 2\theta, \quad 0 \leq \theta \leq \frac{\pi}{4}$$

The line  $l$  is parallel to the initial line and is a tangent to  $C$ .

Find an equation of  $l$ , giving your answer in the form  $r = f(\theta)$ . (9)

$$\frac{dy}{d\theta} = 0 \quad y = r \sin \theta = 2 \cos 2\theta \sin \theta = 2(1 - 2 \sin^2 \theta) \sin \theta$$

$$\therefore y = 2 \sin \theta - 2 \sin^3 \theta$$

$$\frac{dy}{d\theta} = 2 \cos \theta - 6 \sin^2 \theta \cos \theta = 0 \Rightarrow 2 \cos \theta = 6 \sin^2 \theta \cos \theta$$

$$\therefore \sin^2 \theta = \frac{1}{3} \quad \therefore \sin \theta = \pm \frac{1}{\sqrt{3}}$$

$$\theta = 0.6154\dots$$

$$\begin{aligned} r &= 2(1 - 2 \sin^2 \theta) = 2\left(1 - 2\left(\frac{1}{3}\right)\right) \\ &= 2\left(\frac{1}{3}\right) = \frac{2}{3} \end{aligned}$$

$$y = r \sin \theta = \frac{2}{3} \left(\frac{1}{\sqrt{3}}\right) = \frac{2\sqrt{3}}{9} \quad \therefore L \Rightarrow y = \frac{2\sqrt{3}}{9}$$

$$r = \frac{y}{\sin \theta} \Rightarrow r = y \operatorname{cosec} \theta$$

$$\therefore r = \frac{2\sqrt{3}}{9} \operatorname{cosec} \theta$$

5.

$$y \frac{d^2y}{dx^2} + 2\left(\frac{dy}{dx}\right)^2 + 2y = 0$$

- (a) Find an expression for  $\frac{d^3y}{dx^3}$  in terms of  $\frac{d^2y}{dx^2}$ ,  $\frac{dy}{dx}$  and  $y$ .

(4)

Given that  $y = 2$  and  $\frac{dy}{dx} = 0.5$  at  $x = 0$ ,

- (b) find a series solution for  $y$  in ascending powers of  $x$ , up to and including the term in  $x^3$ .

(5)

$$\frac{d}{dx} \left( y \frac{d^2y}{dx^2} \right) + 2 \frac{d}{dx} \left( \frac{dy}{dx} \right)^2 + 2 \frac{d}{dx}(y) = 0$$

$$y \frac{d^3y}{dx^3} + \frac{dy}{dx} \frac{d^2y}{dx^2} + 4\left(\frac{dy}{dx}\right) \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 0$$

$$y \frac{d^3y}{dx^3} + 5\left(\frac{dy}{dx}\right) \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} = 0$$

$$x_0 = 0 \quad y_0 = 2 \quad y'_0 = \frac{1}{2}$$

$$2y''_0 + 2\left(\frac{1}{2}\right)^2 + 2(2) = 0 \Rightarrow 2y''_0 = -\frac{9}{2} \therefore y''_0 = -\frac{9}{4}$$

$$2y'''_0 + 5\left(\frac{1}{2}\right)\left(-\frac{9}{4}\right) + 2\left(\frac{1}{2}\right) = 0 \Rightarrow 2y'''_0 = -1 + \frac{45}{8} = \frac{37}{8}$$

$$y''''_0 = \frac{37}{16}$$

$$\therefore y = 2 + \frac{1}{2}x + -\frac{9}{8}x^2 + \frac{37}{96}x^3$$

6. The transformation  $T$  maps points from the  $z$ -plane, where  $z = x + iy$ , to the  $w$ -plane, where  $w = u + iv$ .

The transformation  $T$  is given by

$$w = \frac{z}{iz + 1}, \quad z \neq i$$

The transformation  $T$  maps the line  $l$  in the  $z$ -plane onto the line with equation  $v = -1$  in the  $w$ -plane.

- (a) Find a cartesian equation of  $l$  in terms of  $x$  and  $y$ .

(5)

The transformation  $T$  maps the line with equation  $y = \frac{1}{2}$  in the  $z$ -plane onto the curve  $C$  in the  $w$ -plane.

- (b) (i) Show that  $C$  is a circle with centre the origin.

- (ii) Write down a cartesian equation of  $C$  in terms of  $u$  and  $v$ .

(6)

$$\begin{aligned} u + iv &= \frac{x+iy}{i(x+iy)+1} = \frac{x+iy}{(1-y)+ix} = \left[ \frac{x+iy}{(1-y)+ix} \right] \left[ \frac{(1-y)-ix}{(1-y)-ix} \right] \\ &= \frac{(x(1-y)+xy) + i(y(1-y)-x^2)}{(1-y)^2+x^2} \end{aligned}$$

$$v = -1 \Rightarrow \frac{y-y^2-x^2}{(1-y)^2+x^2} = -1 \Rightarrow y - y^2 - x^2 = -1 + 2y - y^2 - x^2 \Rightarrow y = 1$$

$$\text{5) } wiz + w = z \Rightarrow w = z - wiz \Rightarrow w = z(1 - wi)$$

$$\begin{aligned} z &= \frac{w}{1-wi} \Rightarrow x+iy = \frac{u+iv}{1-(u+iv)i} = \left[ \frac{u+iv}{(1+v)-ui} \right] \left[ \frac{(1+v)+ui}{(1+v)+ui} \right] \\ x+iy &\equiv \frac{(u(1+v)-uv) + i(v(1+v)+u^2)}{(1+v)^2+u^2} \end{aligned}$$

$$y = \frac{1}{2} \quad \therefore \frac{v+v^2+u^2}{(1+v)^2+u^2} = \frac{1}{2} \Rightarrow \frac{2v+2v^2+2u^2}{(1+v)^2+u^2} = 1 \Rightarrow u^2+v^2 = 1$$

ii)  $u^2 + v^2 = 1$

Circle centre  $O$ ,  $r = 1$

7. (a) Use de Moivre's theorem to show that

$$\sin 5\theta \equiv 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta$$

(5)

- (b) Hence find the five distinct solutions of the equation

$$16x^5 - 20x^3 + 5x + \frac{1}{2} = 0$$

giving your answers to 3 decimal places where necessary.

(5)

- (c) Use the identity given in (a) to find

$$\int_0^{\pi} (4 \sin^5 \theta - 5 \sin^3 \theta) d\theta$$

expressing your answer in the form  $a\sqrt{2} + b$ , where  $a$  and  $b$  are rational numbers.

(4)

$$(\cos \theta + i \sin \theta)^5 = \cos 5\theta + i \sin 5\theta$$

$$\begin{array}{r} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ 10 \\ 11 \\ 12 \\ 13 \\ 14 \\ 15 \\ 16 \\ 17 \\ 18 \\ 19 \\ 20 \\ 21 \\ 22 \\ 23 \\ 24 \\ 25 \\ 26 \\ 27 \\ 28 \\ 29 \\ 30 \\ 31 \end{array}$$

$$(\cos \theta + i \sin \theta)^5 = \cos^5 \theta + 5i \cos^4 \theta \sin \theta - 10 \cos^3 \theta \sin^2 \theta - 10i \cos^2 \theta \sin^3 \theta + 5 \cos \theta \sin^4 \theta + i \sin^5 \theta$$

Comparing Imaginary parts

$$\begin{aligned} \Rightarrow \sin 5\theta &= 5 \cos^4 \theta \sin \theta - 10 \cos^2 \theta \sin^3 \theta + \sin 5\theta \\ &= 5(1 - 2 \sin^2 \theta + \sin^4 \theta) \sin \theta - 10(1 - \sin^2 \theta) \sin^3 \theta + \sin 5\theta \\ &= (5 - 10 \sin^2 \theta + 5 \sin^4 \theta) \sin \theta + (10 \sin^2 \theta - 10) \sin^3 \theta + \sin 5\theta \\ &= 5 \sin \theta - 10 \sin^3 \theta + 5 \sin^5 \theta + 10 \sin^3 \theta - 10 \sin^3 \theta + \sin 5\theta \end{aligned}$$

$$\therefore \sin 5\theta = 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta$$

$$b) \text{ if } x = \sin \theta \Rightarrow 16 \sin^5 \theta - 20 \sin^3 \theta + 5 \sin \theta = -\frac{1}{2}$$

$$\Rightarrow \sin 5\theta = -\frac{1}{2} \Rightarrow 5\theta = \frac{7\pi}{6}, \frac{11\pi}{6}, \frac{19\pi}{6}, \frac{23\pi}{6}, \frac{31\pi}{6}$$

$$\therefore \theta = \frac{7}{30}\pi, \frac{11}{30}\pi, \frac{19}{30}\pi, \frac{23}{30}\pi, \frac{31}{30}\pi, \dots$$

$$x = \sin \theta \therefore x = 0.823,$$

$$x = 0.669, 0.914, -0.105, -0.51, -0.978$$

2

$$b) \frac{1}{4} \sin 5\theta = 4 \sin^5 \theta - 5 \sin^3 \theta + \frac{5}{4} \sin \theta$$

$$\therefore 4 \sin^5 \theta - 5 \sin^3 \theta = \frac{1}{4} (\sin 5\theta - 5 \sin \theta)$$

$$\frac{1}{4} \int_0^{\frac{\pi}{4}} \sin 5\theta - 5 \sin \theta d\theta = \frac{1}{4} \left[ -\frac{1}{5} \cos 5\theta + 5 \cos \theta \right]_0^{\frac{\pi}{4}}$$

$$= \frac{1}{4} \left[ \left( \frac{\sqrt{2}}{10} + \frac{5\sqrt{2}}{2} \right) - \left( -\frac{1}{5} + 5 \right) \right]$$

$$= \frac{13}{20} \sqrt{20} - \frac{6}{5}$$

2

8. (a) Show that the substitution  $x = e^z$  transforms the differential equation

$$x^2 \frac{d^2y}{dx^2} + 2x \frac{dy}{dx} - 2y = 3 \ln x, \quad x > 0 \quad (\text{I})$$

into the equation

$$\frac{d^2y}{dz^2} + \frac{dy}{dz} - 2y = 3z \quad (\text{II})$$

(7)

- (b) Find the general solution of the differential equation (II). (6)

- (c) Hence obtain the general solution of the differential equation (I) giving your answer in the form  $y = f(x)$ . (1)

$$\frac{dx}{dz} = e^z \Rightarrow \frac{dz}{dx} = e^{-z} \quad x = e^z \quad x^2 = e^{2z}$$

$$\ln x = \ln e^z = z$$

$$\frac{dy}{dx} = \frac{dy}{dz} \frac{dz}{dx} = e^{-z} \frac{dy}{dz}$$

$$\frac{d^2y}{dx^2} = e^{-z} \left[ \frac{d}{dz} \left( \frac{dy}{dz} \right) \right] + \frac{du}{dz} \left[ \frac{d}{dx} e^{-z} \right]$$

$$= e^{-z} \frac{d^2y}{dz^2} \frac{dz}{dx} + \frac{dy}{dz} - e^{-z} \frac{dz}{dx}$$

$$= e^{-2z} \frac{d^2y}{dz^2} - e^{-2z} \frac{dy}{dz} = e^{-2z} \left( \frac{d^2y}{dz^2} - \frac{dy}{dz} \right)$$

$$\therefore x^2 \frac{d^2y}{dx^2} + 2x \frac{dy}{dx} - 2y = 3 \ln x \Rightarrow$$

$$e^{2z} \left( e^{-2z} \left( \frac{d^2y}{dz^2} - \frac{dy}{dz} \right) \right) + 2e^z \left( e^{2z} \frac{dy}{dz} \right) - 2y = 3z$$

$$= \frac{d^2y}{dz^2} + \frac{dy}{dz} - 2y = 3z \quad *$$

$$\textcircled{4} \quad y = Ae^{Mz}$$

$$y' = Ame^{Mz}$$

$$y'' = Am^2e^{Mz}$$

$$y'' + y' - 2y = 0$$

$$Ae^{Mz}(m^2 + m - 2) = 0$$

$$\neq 0 \quad (m+2)(m-1) = 0$$

$$\therefore m = -2, m = 1$$

$$y_{GS} = Ae^z + Be^{-2z}$$

(PI)

$$y = a + bz + cz^2$$

$$y' = b + 2cz$$

$$y'' = 2c$$

$$\begin{aligned} -2y &= -2a - 2bz - 2cz^2 \\ +y' &= b + 2cz \\ \hline \frac{+y''}{3z} &= \frac{2c}{(-2a+b+2c)z + (-2b+2c)} - 2cz^2 \end{aligned}$$

$$\therefore c = 0 \quad -2b + 2c = 3 \quad \therefore b = -\frac{3}{2}$$

$$-2a - \frac{3}{2} + 0 = 0 \quad \therefore 2a = -\frac{3}{2} \quad \therefore a = -\frac{3}{4}$$

$$\therefore y_{PI} = -\frac{3}{4} - \frac{3}{2}z$$

$$\therefore y_{GS} = Ae^z + Be^{-2z} - \frac{3}{2}z - \frac{3}{4} \quad \hookrightarrow x = e^z \therefore \ln x = z$$

$$\therefore y = Ae^{\ln x} + Be^{-2\ln x} - \frac{3}{2}\ln x - \frac{3}{4}$$

$$y = Ax + \frac{B}{x^2} - \frac{3}{2}\ln x - \frac{3}{4}$$