

CAMBRIDGE INTERNATIONAL EXAMINATIONS
GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2012 series

9702 PHYSICS

9702/22

Paper 2 (AS Structured Questions), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9702	22

- 1 (a) units for D identified as kg m s^{-2} M1
 all other units shown: units for A : m^2 units for v^2 : $\text{m}^2 \text{s}^{-2}$ units for ρ : kg m^{-3}
- $$C = \frac{\text{kg m s}^{-2}}{\text{kg m}^{-3} \text{m}^2 \text{m}^2 \text{s}^{-2}}$$
- with cancelling / simplification to give
- C
- no units A1 [2]
- (b) (i) straight line from (0,0) to (1,9.8) \pm half a square B1 [1]
- (ii) $\frac{1}{2} mv^2 = mgh$ or using $v^2 = 2as$ C1
 $v = (2 \times 9.81 \times 1000)^{1/2} = 140 \text{ m s}^{-1}$ A1 [2]
- (c) (i) weight = drag (D) (+ upthrust) B1 [1]
 Allow mg or W for weight and D or expression for D for drag
- (ii) 1. $mg = 1.4 \times 10^{-5} \times 9.81$ C1
 $1.4 \times 10^{-5} \times 9.81 = 0.5 \times 0.6 \times 1.2 \times 7.1 \times 10^{-6} \times v^2$ M1
 $v = 7.33 \text{ m s}^{-1}$ A0 [2]
2. line from (0,0) correct curvature to a horizontal line at velocity of 7 m s^{-1} M1
 line reaches 7 m s^{-1} between 1.5s and 3.5s A1 [2]
- 2 (a) (resultant) force = rate of change of momentum / allow proportional to B1 [1]
 or change in momentum / time (taken)
- (b) (i) $\Delta p = (-) 65 \times 10^{-3} (5.2 + 3.7)$ C1
 $= (-) 0.58 \text{ N s}$ A1 [2]
- (ii) $F = 0.58 / 7.5 \times 10^{-3}$
 $= 77(.3) \text{ N}$ A1 [1]
- (c) (i) 1. force on the wall from the ball is equal to the force on ball from the wall M1
 but in the opposite direction A1 [2]
 (statement of Newton's third law can score one mark)
2. momentum change of ball is equal and opposite to momentum change B1 [1]
 of the wall / change of momentum of ball and wall is zero
- (ii) kinetic energy (of ball and wall) is reduced / not conserved so inelastic B1 [1]
 (Allow relative speed of approach does not equal relative speed of separation.)

Page 3	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9702	22

- 3 (a) metal: regular / repeated / ordered arrangement / pattern / lattice or long range order (of atoms / molecules / ions) B1
- polymer: tangled chains (of atoms / molecules) or long chains (of atoms / molecules / ions) B1
- amorphous: disordered / irregular arrangement or short range order (of atoms / molecules / ions) B1 [3]
- (b) metal: straight line or straight line then curving with less positive gradient B1
- polymer: curve with decreasing gradient with steep increasing gradient at end B1 [2]
- 4 (a) waves (travels along tube) reflect at closed end / end of tube B1
- incident and reflected waves or these two waves are in opposite directions M1
- interfere or stationary wave formed if tube length equivalent to $\lambda / 4, 3\lambda / 4, \text{etc.}$ A1 [3]
- (b) (i) 1. no motion (as node) / zero amplitude B1 [1]
2. vibration backwards and forwards / maximum amplitude along length B1 [1]
- (ii) $\lambda = 330 / 880 (= 0.375 \text{ m})$ C1
- $L = 3\lambda / 4$ C1
- $L = 3 / 4 \times (0.375) = 0.28 (0.281) \text{ m}$ A1 [3]
- 5 (a) (i) $I_1 = I_2 + I_3$ B1 [1]
- (ii) $I = V / R$ or $I_2 = 12 / 10 (= 1.2 \text{ A})$ C1
- $R = [1/6 + 1 / 10]^{-1}$ [total $R = 3.75 \Omega$] or $I_3 = 12 / 6 (= 2.0 \text{ A})$ C1
- $I_1 = 12 / 3.75 = 3.2 \text{ A}$ or $I_1 = 1.2 + 2.0 = 3.2 \text{ A}$ A1 [3]
- (iii) power = VI or I^2R or V^2 / R C1
- $$x = \frac{\text{power in wire}}{\text{power in series resistors}} = \frac{I_2^2 R_w}{I_3^2 R_s} \text{ or } \frac{VI_2}{VI_3} \text{ or } \frac{V^2 / R_w}{V^2 / R_s}$$
- $x = 12 \times 1.2 / 12 \times 2.0 = 0.6(0)$ allow 3 / 5 or 3:5 A1 [3]
- (b) p.d. BC: $12 - 12 \times 0.4 = 7.2 \text{ (V)}$ / p.d. AC = 4.8 (V) C1
- p.d. BD: $12 - 12 \times 4 / 6 = 4.0 \text{ (V)}$ / p.d. AD = 8.0 (V) C1
- p.d. = 3.2V A1 [3]
- 6 (a) extension is proportional to force / load B1 [1]
- (b) $F = mg$ C1
- $x = (mg / k) = 0.41 \times 9.81 / 25 = (4.02 / 25)$ M1
- $x = 0.16 \text{ m}$ A0 [2]

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – October/November 2012	9702	22

- (c) (i) weight and (reaction) force from spring (which is equal to tension in spring) B1 [1]
- (ii) $F = \text{weight}$ or $0.06 \times 25 = ma$ C1
 $F = 0.2209 \times 25 = 5.52 \text{ (N)}$ or $0.22 \times 25 = 5.5$
 $a = (5.52 - 0.41 \times 9.81) / 0.41$ or $1.5 / 0.41$ and $(5.5 - 4.02)$ C1
 $a = 3.7 \text{ (3.66) m s}^{-2}$ gives 3.6 m s^{-2} A1 [3]
- (d) elastic potential energy / strain energy to kinetic energy and gravitational potential energy B1
stretching / extension reduces and velocity increases / height increases B1 [2]
- 7 (a) ${}^3_2\text{He} + {}^3_2\text{He} \rightarrow {}^4_2\text{He} + 2 {}^1_1\text{p} + Q$
A numbers correct (4 and 1) B1
Z numbers correct (2 and 1) B1 [2]
- (b) both nuclei have 2 protons B1
the two isotopes have 1 neutron and two neutrons B1 [2]
[allow 1 for 'same number of protons but different number of neutrons']
- (c) proton number and neutron number B1
energy – mass B1
momentum B1 [2]
- (d) (i) γ radiation B1 [1]
- (ii) product(s) must have kinetic energy B1 [1]
- (e) $13.8 \text{ MeV} = 13.8 \times 1.6 \times 10^{-19} \times 10^6 (= 2.208 \times 10^{-12})$ C1
 $60 = n \times 13.8 \times 1.6 \times 10^{-13}$
 $n = 2.7(2) \times 10^{13} \text{ s}^{-1}$ A1 [2]