



Mark Scheme (Results)

Summer 2014

Pearson Edexcel GCE
in Physics (6PH04)
Paper 01 Physics on the Move

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2014

Publications Code UA039725

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Mark Scheme Notes

Underlying principle

The mark scheme will clearly indicate the concept that is being rewarded, backed up by examples. It is not a set of model answers.

For example:

(iii)	<u>Horizontal force of hinge on table top</u> 66.3 (N) or 66 (N) and correct indication of direction [no ue] [Some examples of direction: acting from right (to left) / to the left / West / opposite direction to horizontal. May show direction by arrow. Do not accept a minus sign in front of number as direction.]	✓	1
-------	---	---	---

This has a clear statement of the principle for awarding the mark, supported by some examples illustrating acceptable boundaries.

1. Mark scheme format

- 1.1 You will not see 'wtte' (words to that effect). Alternative correct wording should be credited in every answer unless the ms has specified specific words that must be present. Such words will be indicated by underlining e.g. 'resonance'
- 1.2 Bold lower case will be used for emphasis.
- 1.3 Round brackets () indicate words that are not essential e.g. "(hence) distance is increased".
- 1.4 Square brackets [] indicate advice to examiners or examples e.g. [Do not accept gravity] [ecf].

2. Unit error penalties

- 2.1 A separate mark is not usually given for a unit but a missing or incorrect unit will normally mean that the final calculation mark will not be awarded.
- 2.2 Incorrect use of case e.g. 'Watt' or 'w' will **not** be penalised.
- 2.3 There will be no unit penalty applied in 'show that' questions or in any other question where the units to be used have been given, for example in a spreadsheet.
- 2.4 The same missing or incorrect unit will not be penalised more than once within one question (one clip in epen).
- 2.5 Occasionally, it may be decided not to penalise a missing or incorrect unit e.g. the candidate may be calculating the gradient of a graph, resulting in a unit that is not one that should be known and is complex.
- 2.6 The mark scheme will indicate if no unit error penalty is to be applied by means of [no ue].

3. Significant figures

- 3.1 Use of an inappropriate number of significant figures in the theory papers will normally only be penalised in 'show that' questions where use of too few significant figures has resulted in the candidate not demonstrating the validity of the given answer.
- 3.2 The use of $g = 10 \text{ m s}^{-2}$ or 10 N kg^{-1} instead of 9.81 m s^{-2} or 9.81 N kg^{-1} will be penalised by one mark (but not more than once per clip). Accept 9.8 m s^{-2} or 9.8 N kg^{-1}

4. Calculations

- 4.1 Bald (i.e. no working shown) correct answers score full marks unless in a 'show that' question.
- 4.2 If a 'show that' question is worth 2 marks then both marks will be available for a reverse working; if it is worth 3 marks then only 2 will be available.
- 4.3 **use** of the formula means that the candidate demonstrates substitution of physically correct values, although there may be conversion errors e.g. power of 10 error.
- 4.4 **recall** of the correct formula will be awarded when the formula is seen or implied by substitution.
- 4.5 The mark scheme will show a correctly worked answer for illustration only.
- 4.6 Example of mark scheme for a calculation:

<u>'Show that' calculation of weight</u>		
Use of $L \times W \times H$	✓	
Substitution into density equation with a volume and density	✓	
Correct answer [49.4 (N)] to at least 3 sig fig. [No ue]	✓	
[If 5040 g rounded to 5000 g or 5 kg, do not give 3 rd mark; if conversion to kg is omitted and then answer fudged, do not give 3 rd mark]		
[Bald answer scores 0, reverse calculation 2/3]		3
Example of answer:		
$80 \text{ cm} \times 50 \text{ cm} \times 1.8 \text{ cm} = 7200 \text{ cm}^3$		
$7200 \text{ cm}^3 \times 0.70 \text{ g cm}^{-3} = 5040 \text{ g}$		

$5040 \times 10^{-3} \text{ kg} \times 9.81 \text{ N/kg}$		
$= 49.4 \text{ N}$		

5. Quality of Written Communication


- 5.1 Indicated by QoWC in mark scheme. QWC – Work must be clear and organised in a logical manner using technical wording where appropriate.
- 5.2 Usually it is part of a max mark, the final mark not being awarded unless the QoWC condition has been satisfied.

6. Graphs

- 6.1 A mark given for axes requires both axes to be labelled with quantities and units, and drawn the correct way round.
- 6.2 Sometimes a separate mark will be given for units or for each axis if the units are complex. This will be indicated on the mark scheme.
- 6.3 A mark given for choosing a scale requires that the chosen scale allows all points to be plotted, spreads plotted points over more than half of each axis and is not an awkward scale e.g. multiples of 3, 7 etc.
- 6.4 Points should be plotted to within 1 mm.
 - Check the two points furthest from the best line. If both OK award mark.
 - If either is 2 mm out do not award mark.
 - If both are 1 mm out do not award mark.
 - If either is 1 mm out then check another two and award mark if both of these OK, otherwise no mark.
- 6.5 For a line mark there must be a thin continuous line which is the best-fit line for the candidate's results.

Question Number	Answer	Mark
1	C	1
2	B	1
3	B	1
4	D	1
5	A	1
6	C	1
7	C	1
8	C	1
9	B	1
10	A	1

Question Number	Answer	Mark
*11	<p>(QWC – Work must be clear and organised in a logical manner using technical wording where appropriate) Either</p> <p>Initial momentum is zero</p> <p>Nucleus and alpha particle have equal momentum (1)</p> <p>(accept $m_n u_n = m_\alpha u_\alpha$ or $p_n = p_\alpha$) alpha particle and nucleus move in opposite directions Mass of alpha particle < mass of nucleus (therefore $v_n < v_\alpha$) (1)</p> <p>Or</p> <p>The nucleus and alpha particle exert an equal but opposite force on each other. (1)</p> <p>Mass of alpha particle < mass of nucleus (1)</p> <p>Acceleration of nucleus < acceleration of alpha particle</p> <p>Force/acceleration acts for same time so Δv for nucleus is smaller for nucleus (1)</p>	4
Total for question 11		4

Question Number	Answer	Mark
12	<p>Free body force diagram showing 2 forces only</p> <p>Weight/W/mg (1)</p> <p>Tension / T (1)</p> <p>(Each additional forces e.g. horizontal component or resultant force, 1 mark penalty)</p> <p>If θ is angle to the vertical then:</p> <p>(Resolving vertically): $T\cos\theta = mg$ (1)</p> <p>(Resolving horizontally): $T\sin\theta = mv^2/r$ Or $T\sin\theta = mr\omega^2$ (1)</p> <p>Derives $\tan\theta = v^2/rg$ and links to observations</p> <p>Or Derives $\tan\theta = r\omega^2/g$ and links to observations (1)</p> <p>If angle to horizontal is used candidates can score MP3 and 4.[then sin and cos swop over and tan of angle will be reciprocal of above]</p> <p><u>Examples of free body force diagrams</u></p>  <p>(full credit for the last 3 marks can be given to candidates who draw a vector triangle and derive $\tan\theta = T_{\text{horzt}}/mg$ and then $\tan\theta = r\omega^2/g$ and observation)</p>	5
Total for question 12		5

Question Number	Answer	Mark
16(a)	(Electric field strength (at a point in a field) is) the force per unit charge (accept force per coulomb of charge) Acting on a (small) positive charge.	(1) (1) 2
16(b)(i)	Use of $E = kQ/r^2$ Electric field due to $Q_1 = 4.1(1) \times 10^6 \text{ (N C}^{-1}\text{)}$ Use of 11.9 cm to find field due to Q_2 Or Use of $E = kQ/r^2$ Use of $E_1/E_2 = Q_1/r_1^2 / Q_2/r_2^2$ $E_1/E_2 = 1$ <u>Example of calculation</u> Electric field due to Q_1 $= (8.99 \times 10^9 \text{ N m}^2 \text{ C}^{-2}) \times (3 \times 10^{-6} \text{ C}) / (8.1 \times 10^{-2})^2$ $= 4.11 \times 10^6 \text{ N C}^{-1}$ Electric field due to Q_2 $= (8.99 \times 10^9 \text{ N m}^2 \text{ C}^{-2}) \times (6.5 \times 10^{-6} \text{ C}) / (11.9 \times 10^{-2})^2 = 4.13 \times 10^6 \text{ N C}^{-1}$	(1) (1) (1) (1) (1) (1) 3
16(b)(ii)	(Force on charge is) zero/negligible/approx zero (Allow values less than 0.1 N)	(1) 1
16(b)(iii)	At midpoint repulsive force due to $Q_2 >$ repulsive force due to Q_1 Or the <u>resultant</u> field/force is repulsive <u>Work</u> must be done against the repulsive force/field to move the charge to this position.	(1) (1) 2
	Total for question 16	8

7(c)	Max 3 Ultracapacitor used for: overtaking Or going up a hill Or starting (from rest) Or accelerating. (1) Because this requires a large <u>current/power</u> . (1) Batteries used for travelling at constant speed (1) Because this requires a small <u>current/power</u> for a longer time (1)	3
	Total for question 17	15

Question Number	Answer	Mark								
*18(a)	<p>(QWC – Work must be clear and organised in a logical manner using technical wording where appropriate)</p> <p>Electric field Provides a force on the proton/particle (1) Which accelerate the proton/particle Or gives energy to the protons/particles (1)</p> <p>Magnetic field Provides a force on a moving proton Or Provides a force at right angles to the direction of motion (of the protons) (1) Acts as a centripetal force Or produces circular motion (1)</p> <p>Additional detail about either field E field across gap only Or The idea that the E field is reversed /alternates every half cycle Or B field perpendicular to the Dees (1)</p> <p>(this mark may be awarded from a diagram)</p>	5								
18(b)	<p>Division by e (ignore powers of 10 error) (1) multiplication by c^2 (1) Mass = 0.14 (GeV/c^2) (1)</p> <p><u>Example of calculation</u> Mass = $(2.5 \times 10^{-28} \text{ kg} \times 9 \times 10^{16} \text{ m}^2 \text{ s}^{-2}) / 1.6 \times 10^{-19} \text{ C}$ Mass = $0.14 \times 10^9 \text{ eV}/c^2 = 0.14 \text{ GeV}/c^2$</p>	3								
18(c)	<p>2/3 charge of a proton Or 2/3 charge of a positron (1) Or 2/3 <u>positive</u> value of the charge on an electron Or $2/3e^+$</p>	1								
18(d)(i)	<table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Particle</th> <th>Quark combination</th> </tr> </thead> <tbody> <tr> <td>K^-</td> <td>$\bar{s}u$</td> </tr> <tr> <td>K^+</td> <td>$u\bar{s}$</td> </tr> <tr> <td>K^0</td> <td>$\bar{s}d$ or $d\bar{s}$</td> </tr> </tbody> </table>	Particle	Quark combination	K^-	$\bar{s}u$	K^+	$u\bar{s}$	K^0	$\bar{s}d$ or $d\bar{s}$	3
Particle	Quark combination									
K^-	$\bar{s}u$									
K^+	$u\bar{s}$									
K^0	$\bar{s}d$ or $d\bar{s}$									
18(d)(ii)	<p>Mass-energy is conserved Or a comment about $E = mc^2$ (1) Appropriate reference to colliding particles having mass and kinetic energy The extra mass comes from the <u>kinetic</u> energy. (1)</p>	3								
Total for question 18		15								

