

GCE

Physics A

H556/02: Exploring physics

A Level

Mark Scheme for June 2023

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2023

MARKING INSTRUCTIONS

PREPARATION FOR MARKING RM ASSESSOR

- 1. Make sure that you have accessed and completed the relevant training packages for on-screen marking: *RM* Assessor Assessor Online Training; *OCR* Essential Guide to Marking.
- 2. Make sure that you have read and understood the mark scheme and the question paper for this unit. These are posted on the RM Cambridge Assessment Support Portal <u>http://www.rm.com/support/ca</u>
- 3. Log-in to RM Assessor and mark the **required number** of practice responses ("scripts") and the **number of required** standardisation responses.

YOU MUST MARK 10 PRACTICE AND 10 STANDARDISATION RESPONSES BEFORE YOU CAN BE APPROVED TO MARK LIVE SCRIPTS.

MARKING

- 1. Mark strictly to the mark scheme.
- 2. Marks awarded must relate directly to the marking criteria.
- 3. The schedule of dates is very important. It is essential that you meet the RM Assessor 50% and 100% (traditional 40% Batch 1 and 100% Batch 2) deadlines. If you experience problems, you must contact your Team Leader (Supervisor) without delay.
- 4. If you are in any doubt about applying the mark scheme, consult your Team Leader by telephone or the RM Assessor messaging system, or by email.

5. Crossed Out Responses

Where a candidate has crossed out a response and provided a clear alternative then the crossed out response is not marked. Where no alternative response has been provided, examiners may give candidates the benefit of the doubt and mark the crossed out response where legible.

Rubric Error Responses – Optional Questions

Where candidates have a choice of question across a whole paper or a whole section and have provided more answers than required, then all responses are marked and the highest mark allowable within the rubric is given. Enter a mark for each question answered into RM assessor, which will select the

Mark Scheme

highest mark from those awarded. (The underlying assumption is that the candidate has penalised themselves by attempting more questions than necessary in the time allowed.)

Multiple Choice Question Responses

When a multiple choice question has only a single, correct response and a candidate provides two responses (even if one of these responses is correct), then no mark should be awarded (as it is not possible to determine which was the first response selected by the candidate). When a question requires candidates to select more than one option/multiple options, then local marking arrangements need to ensure consistency of approach.

Contradictory Responses

When a candidate provides contradictory responses, then no mark should be awarded, even if one of the answers is correct.

Short Answer Questions (requiring only a list by way of a response, usually worth only one mark per response)

Where candidates are required to provide a set number of short answer responses then only the set number of responses should be marked. The response space should be marked from left to right on each line and then line by line until the required number of responses have been considered. The remaining responses should not then be marked. Examiners will have to apply judgement as to whether a 'second response' on a line is a development of the 'first response', rather than a separate, discrete response. *(The underlying assumption is that the candidate is attempting to hedge their bets and therefore getting undue benefit rather than engaging with the question and giving the most relevant/correct responses.)*

Short Answer Questions (requiring a more developed response, worth two or more marks)

If the candidates are required to provide a description of, say, three items or factors and four items or factors are provided, then mark on a similar basis – that is downwards (as it is unlikely in this situation that a candidate will provide more than one response in each section of the response space.)

Longer Answer Questions (requiring a developed response)

Where candidates have provided two (or more) responses to a medium or high tariff question which only required a single (developed) response and not crossed out the first response, then only the first response should be marked. Examiners will need to apply professional judgement as to whether the second (or a subsequent) response is a 'new start' or simply a poorly expressed continuation of the first response.

- 6. Always check the pages (and additional objects if present) at the end of the response in case any answers have been continued there. If the candidate has continued an answer there, then add a tick to confirm that the work has been seen.
- 7. Award No Response (NR) if:
 - there is nothing written in the answer space

Award Zero '0' if:

• anything is written in the answer space and is not worthy of credit (this includes text and symbols).

Team Leaders must confirm the correct use of the NR button with their markers before live marking commences and should check this when reviewing scripts.

- 8. The RM Assessor comments box is used by your team leader to explain the marking of the practice responses. Please refer to these comments when checking your practice responses. Do not use the comments box for any other reason. If you have any questions or comments for your team leader, use the phone, the RM Assessor messaging system, or e-mail.
- 9. Assistant Examiners will send a brief report on the performance of candidates to their Team Leader (Supervisor) via email by the end of the marking period. The report should contain notes on particular strengths displayed as well as common errors or weaknesses. Constructive criticism of the question paper/mark scheme is also appreciated.
- 10. For answers marked by levels of response:
 - a. To determine the level start at the highest level and work down until you reach the level that matches the answer
 - b. To determine the mark within the level, consider the following

Descriptor	Award mark		
On the borderline of this level and the one below	At bottom of level		
Just enough achievement on balance for this level	Above bottom and either below middle or at middle of level (depending on number of marks available)		
Meets the criteria but with some slight inconsistency	Above middle and either below top of level or at middle of level (depending on number of marks available)		
Consistently meets the criteria for this level	At top of level		

11. Annotations

	Annotation	Meaning
~	Correct response	Used to indicate the point at which a mark has been awarded (one tick per mark awarded).
×	Incorrect response	Used to indicate an incorrect answer or a point where a mark is lost.
AE	Arithmetic error	Do not allow the mark where the error occurs. Then follow through the working/calculation giving full subsequent ECF if there are no further errors.
BOD	Benefit of doubt given	Used to indicate a mark awarded where the candidate provides an answer that is not totally satisfactory, but the examiner feels that sufficient work has been done.
ВР	Blank page	Use BP on additional page(s) to show that there is no additional work provided by the candidates.
CON	Contradiction	No mark can be awarded if the candidate contradicts himself or herself in the same response.
ECF	Error carried forward	Used in <u>numerical answers only</u> , unless specified otherwise in the mark scheme. Answers to later sections of numerical questions may be awarded up to full credit provided they are consistent with earlier incorrect answers. Within a question, ECF can be given for AE, TE and POT errors but not for XP.
L1	Level 1	L1 is used to show 2 marks awarded and L1 [^] is used to show 1 mark awarded.
L2	Level 2	L2 is used to show 4 marks awarded and L2 [^] is used to show 3 marks awarded.
L3	B Level 3 L3 is used to show 6 marks awarded and L3 [^] is used to show 5 marks awarded.	
РОТ	This is usually linked to conversion of SI prefixes. Do not allow the mark where the error occurs. Then follow through the working/calculation giving ECF for subsequent marks if there are no further errors.	

Annotation		Meaning				
SEEN	Seen	To indicate working/text has been seen by the examiner.				
SFError in number of significant figuresWhere more SFs are given than is justified by the question, do not penalise. Fewer significant figures than new will be considered within the mark scheme. Penalised only once in the paper.						
		This error is when there is incorrect transcription of the correct data from the question, graphical read-off, formulae booklet or a previous answer. Do not allow the relevant mark and then follow through the working giving ECF for subsequent marks.				
XP Wrong physics or equation Used in <u>numerical answers only</u> , unless otherwise specified in the mark scheme. Use of an incorrect wrong physics even if it happens to lead to the correct answer.		Used in <u>numerical answers only</u> , unless otherwise specified in the mark scheme. Use of an incorrect equation is wrong physics even if it happens to lead to the correct answer.				
Omission Used t		Used to indicate where more is needed for a mark to be awarded (what is written is not wrong but not enough).				

Annotation	Meaning
1	Alternative and acceptable answers for the same marking point
Reject	Answers which are not worthy of credit
Not	Answers which are not worthy of credit
Ignore	Statements which are irrelevant
Allow	Answers that can be accepted
()	Words which are not essential to gain credit
	Underlined words must be present in answer to score a mark
ECF	Error carried forward
AW	Alternative wording
ORA	Or reverse argument

12. Subject Specific Marking Instructions

CATEGORISATION OF MARKS

The marking schemes categorise marks on the MACB scheme.

- **M** marks These are <u>method</u> marks upon which **A**-marks (accuracy marks) later depend. For an **M**-mark to be scored, the point to which it refers must be seen in the candidate's answers. If a candidate fails to score a particular **M**-mark, then none of the dependent **A**-marks can be scored.
- A marks These are accuracy or <u>answer</u> marks, which either depend on an **M**-mark, or allow a **C**-mark to be scored.
- **C** marks These are <u>compensatory</u> method marks which can be scored even if the points to which they refer are not written down by the candidate, providing subsequent working gives evidence that they must have known it. For example, if an equation carries a **C**-mark and the candidate does not write down the actual equation but does correct working which shows the candidate knew the equation, then the **C**-mark is given.
- **B** marks These are awarded as <u>independent</u> marks, which do not depend on other marks. For a **B**-mark to be scored, the point to which it refers must be seen specifically in the candidate's answers.

SIGNIFICANT FIGURES

If the data given in a question is to 2 sf, then allow an answer to 2 or <u>more</u> significant figures. If an answer is given to fewer than 2 sf, then penalise once only in the <u>entire</u> paper. Any exception to this rule will be mentioned in the Guidance.

June 2023

SECTION A

Question	Answer	Marks	Guidance
1	Α	1	
2	C	1	
3	C	1	
4	Α	1	
5	Α	1	
6	D	1	
7	C	1	
8	Α	1	
9	D	1	
10	C	1	
11	D	1	
12	Α	1	
13	D	1	
14	В	1	
15	C	1	
	Total	15	

SECTION B

General rule: For substitution into an equation, allow any subject - unless stated otherwise in the guidance

	Question	Answer	Mark	Guidance
16	(a)	Interference given in A and D and/or B and C A and D diffraction OR A and D = destructive interference / destructive superposition AND B and C = constructive interference / constructive superposition	B1 B1	IGNORE superposition alone
	(b)	The waves have a constant/same/fixed phase relationship	B1	Constant/same/fixed phase difference IGNORE in phase, frequency, wavelength, diffraction and other correct physics REJECT incorrect references to amplitude and other wave properties
	(c)	Waves arrive in phase / path difference of whole number of wavelengths Phase difference = 4π or 720° / with a path difference of 2λ	C1 A1	Phase difference = 0 or path difference = nλ

Question	Answer	Mark	Guidance
(d)	$\lambda = a x / D = 640 x 10^{-9} = 1.00 x 10^{-5} x x / 4$ (= 0.256 (m))	C1	Correct substitution 0.256m assumes correct substitution
	B to C = $2x = 0.512$ (m)	C1	Recognition that B-C is two fringe separation, and evaluation
	$\theta = \tan^{-1} (0.512 / 4) = 7.3 (\circ)$	A1	Correct to at least 2 significant figures ALLOW $\theta = \sin^{-1} (0.512 / 4) = 7.4 (^{\circ})$
			Note: RAD mode in calculators (0.127) MAX 2 marks
			Note: No doubling of <i>x</i> with correct working leading 3.7° MAX 2 marks
	Alternative method: Use of n $\lambda = d \sin \theta$		
	substitution of λ and <i>d</i> with any integer value of n	(C1)	e.g. n x = 1.00×10^{-5} x sin θ where n is an integer
	recognition that n = 2	(C1)	
	θ = sin ⁻¹ (2 x 640 x 10 ⁻⁹ / 1.00x10 ⁻⁵) = 7.4 (°)	(A1)	Correct to at least 2 significant figures Note: only allow use of incorrect value of $n = 1$, leading to 3.7° MAX 2 marks
			Note: RAD mode in calculators (0.128) MAX 2 marks

Mark Scheme

	Ques	tion	Answer	Mark	Guidance
17	Ques (a) (b*)	tion (i)	AnswerLongitudinal wavesWith a frequency greater than 20 kHzLevel 3 (5–6 marks)Clear explanation of transducer and clear explanation of gelThere is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.Level 2 (3–4 marks)Clear explanation of transducer OR clear explanation of gel	Mark B1 B1 6 x B1	Guidance ALLOW 20 kHz and above Use level of response annotations in RM Assessor Indicative scientific points may include: Transducer • High frequency alternating PD applied to the faces of the piezoelectric crystal. • Causes the piezoelectric crystal to stretch and compress / oscillate / resonate at high frequency. • Ultrasound waves sent in pulses • Between pulses, reflected ultrasound incident on the piezoelectric crystal causes the crystal to change shape / oscillate / resonate
			 or Some explanation of transducer and some explanation of gel There is a line of reasoning presented with some structure. The information presented is in the most part relevant and supported by some evidence. Level 1 (1–2 marks) Limited explanation of transducer or gel There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant. 0 marks No response or no response worthy of credit 		 An alternating PD is induced Gel Acoustic impedance of air is different to body Acoustic impedance of gel is similar to body This allows greater transmission of the ultrasound waves into / out of the patient's body There is less reflection of the ultrasound waves at the body boundary Reference to ^I/_{l_0} = ((Z₂-Z₁)²)/((Z₂+Z₁)²)

Quest	ion	Answer	Mark	Guidance
(c)	(i)	There is no change in (acoustic) impedance		ALLOW there is no change in the density (of the medium) ALLOW there is no boundary / there is nothing to reflect from NOT JUST there is nothing there IGNORE undefined symbols
	(ii)	<u>Acoustic</u> impedance of C is different from B Change in <u>acoustic</u> impedance at B is greater (than at C) (relative to their surrounding medium/A)	C1 A1	IGNORE explanations in terms of density Second point gains both marks ALLOW soft tissue/skin/belly for C ALLOW skull/bone/head for B
(d)		$v = \Delta f c / 2 f \cos \theta$ = $(\frac{10.0000 \times 10^{6} - 9.9987 \times 10^{6}) \times 1600}{2 \times 10.0000 \times 10^{6} \times \cos 50}$ 0.16 (ms ⁻¹)	C1 A1	IGNORE cancelling powers of 10 Correct to at least 2 significant figures

	Question	Answer	Mark	Guidance
18	(a)		B1	Both parts needed – either dotted or solid. Correct curvature needed by eye. Middle node by eye. IGNORE lines outside of tube
	(b)	Frequency <i>f</i> ₀ = 340 / 0.600 = 567 (Hz)	A1	Correct to at least 2 significant figures No ecf from 18(a)
	(c)	Next wavelength for standing wave is $\lambda = 0.300$ m	C1	Ecf from (b) if wavelength used is 0.600m
		Frequency = 340 / 0.300 = 1.13 x 10 ³ (Hz)	A1	Correct to at least 2 significant figures Special case: If wavelength drawn in (a) is 0.300m and f_0 in (b) = 1130 Hz, then allow ecf (C1) for next λ = 0.200m and (A1) frequency as 1700 Hz for full credit.

	Question	Answer	Mark	Guidance
19	(a)	Any four from: In an evacuated tube (Electrons released by) thermionic emission / (low voltage supply causes) emission of electrons from cathode/filament/electron gun Electrons accelerated towards anode / electrons accelerated through (high) pd / electric field Diffracted through a graphite (target) / graphite is thin Electrons detected on a (phosphor) screen / electrons produce visible light on impact with screen	4xB1	ALLOW marking points as labels on the diagram or as the answer lines.
	(b)	 (Provides evidence of) wave nature of electrons Light circles caused by constructive interference / waves arriving in phase Dark circles caused by destructive interference / waves arriving in antiphase 	B1 B1 B1	ALLOW out of phase by 180 / π NOT just out of phase
	(c) (i	Work done = $1.60 \times 10^{-19} \times 5 \times 10^3 = 8.0 \times 10^{-16} (J)$	A1	ALLOW correct answer to 1 significant figure

Question	Answer	Mark	Guidance
(ii)	$W = \frac{1}{2} mv^{2} = 8 \times 10^{-16} = \frac{1}{2} \times 9.11 \times 10^{-31} \times v^{2}$ $= 4.2 \times 10^{7} \text{ ms}^{-1}$ $\lambda = h/mv = 6.63 \times 10^{-34} / 9.11 \times 10^{-31} \times 4.2 \times 10^{7}$	C1	Substitution leading to velocity Ecf from (c)(i)
	= 1.7 x10 ⁻¹¹ (m) OR	A1	ALLOW correct answer to 1 significant figure
	Momentum of electrons = $\sqrt{2 \times m_e \times W}$		
	$= \sqrt{2 \times 9.11 \times 10^{-31} \times 8 \times 10^{-16}} = (3.82 \text{ x}10^{-23} \text{ kgms}^{-1})$ 6.63 x 10 ⁻³⁴ / 3.82 x10 ⁻²³ = 1.74 x10 ⁻¹¹ (m)	(C1) (A1)	Ecf from (c)(i) ALLOW correct answer to 1 significant figure
(iii)	(For diffraction to occur) the gap needs to be approximately the same size as the wavelength so spacing should be $1.74 \times 10^{-11} \text{ m}$	B1	Reason <u>and</u> value required. ALLOW suggested value spacing as (c)(ii) same power of ten

	Question	Answer	Mark	Guidance	
20	(a)	Circuit showing (6V) supply in series with a capacitor and resistor, with a voltmeter in parallel with the capacitor.	B1	No labels required. ALLOW any suitable symbol for d.c. supply	
		Switch/switches allowing discharging of the capacitor through the resistor.	B1	ALLOW this mark if resistor and capacitor in parallel if switch will allow the discharge Examples of correct circuit for both marks	
				e.g.	
	(b)	Charge = 1.0 x 10 ⁻⁶ x 6.0 = 6.0 x 10⁻⁶ (C)	A1	ALLOW correct answer to 1 significant figure	
	(c)	(<i>CR</i> =) 1 x 10 ⁻⁶ x 10 x 10 ³	C1	ALLOW POT error for capacitance if same as in (b)	
		CR / Time constant / τ = 0.010 (s)	A1	ALLOW 1sf ALLOW t for τ	
		Time constant / τ is time taken to fall to 1/e (37%) of initial value	B1	NOT fallen <u>by</u> 37%	
		The time it takes to record the variation of pd / the capacitor to discharge is far less than (human) <u>reaction time</u>	B1		
		OR			

Question	Answer	Mark	Guidance
	$V = V_0 e^{-(\frac{t}{CR})}$ to give $0.6 = 6.0 e^{-(\frac{t}{1.0 \times 10^{-6} \times 10 \times 10^3})}$	(C1)	Substitution into exponential decay equation to fall to a value of <i>V</i> less than $0.9V_0$ (<5.4V). ALLOW calculation in log form e.g ln $0.6 = \ln 6.0 - t/0.01$
	t = 0.023 (s)	(A1)	
	0.6 is the voltage when it has fallen to 10% of the initial voltage	(B1)	Justification/explanation for using a given voltage
	The time it takes to record the variation of pd / for the capacitor to discharge is far less than human <u>reaction time</u>	(B1)	
	OR		
	$V = V_0 e^{-(\frac{t}{CR})}$ to give $V = 6.0 e^{-(\frac{0.02}{1.0 \times 10^{-6} \times 10 \times 10^3})}$	(C1)	Substitution into exponential decay equation with a time less than 0.1s (t < 0.1 s).
	V = 0.81 V	(A1)	ALLOW calculation in log form $\ln V = \ln 6.0 - 0.02/0.01$
	0.02 s is a very short amount of time	(B1)	Justification/explanation for using a given time
	The time it takes to record the variation of pd / for the capacitor to discharge is far less than human <u>reaction time</u>	(B1)	
(d)	Use an oscilloscope / data logger with a voltmeter probe	B1	ALLOW connect a voltmeter to a datalogger
(e) (i)	$V = V_0 e^{-(\frac{t}{CR})}$ to give $4.12 = 6.0 e^{-(\frac{t}{1.0 \times 10^{-6} \times 10 \times 10^3})}$	C1	ALLOW in terms of logs eg $\ln 4.12 = \ln 6.0 - \frac{t}{0.01}$
	= 3.76 x 10 ⁻³ (s)	A1	Correct to at least 2 significant figures IGNORE minus sign in final answer

Question		Answer	Mark	Guidance
(ii)	i)	Change in energy = $\frac{1}{2} CV_1^2 - \frac{1}{2} CV_2^2$	C1	
		= $(\frac{1}{2} \times 1 \times 10^{-6} \times 6^2) - (\frac{1}{2} \times 1 \times 10^{-6} \times 4.12^2) = 9.5 \times 10^{-6} (J)$	C1	ALLOW POT error from (e)(i)
		rate = 9.5 x10 ⁻⁶ / 3.76 x 10 ⁻³		Ecf from (e)(i)
		= 2.53 x 10 ⁻³ (Js ⁻¹)	A1	Correct to at least 2 significant figures

Questio	n	Answer	Mark	Guidance
21 (a)	(i) (ii)	(Nuclear) fuel rods (i) Moderator (ii)	B1 B1	 ALLOW any correctly named fuel e.g. uranium (oxide) for (i) ALLOW any correctly named moderator e.g. (heavy) water for (ii) ALLOW coolant for (ii) ALLOW any correctly named coolant (e.g. CO₂) for (ii)
(b)*		 Level 3 (5–6 marks) Clear description and full correct calculation. There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated. Level 2 (3–4 marks) Clear description or full correct calculation. or Some description and some calculation There is a line of reasoning presented with some structure. The information presented is in the most part relevant and supported by some evidence. 	6 x B1	 Use level of response annotations in RM Assessor Indicative scientific points may include: Description Absorption of neutron by U-235 U-235 nuclei become unstable U236 nuclei U-236 undergoes fission Resulting in smaller nuclei daughter products and additional fast neutrons Moderator slows neutrons so that they can be absorbed by U-235 nuclei Process repeats to form a chain reaction Rate of chain reaction controlled by raising or lowering control rods Which absorb excess neutrons

Question	Answer	Mark	Guidance
	Level 1 (1–2 marks) Limited description or Limited calculation		To allow one neutron per fission to cause further fission
	There is an attempt at a logical structure with a line of reasoning. The information is in the most part		Calculation
	relevant.		Use of ${}^{1}_{0}n + {}^{235}_{92}U \rightarrow {}^{92}_{36}Kr + {}^{141}_{56}Ba + {}^{3}_{0}n$
	0 marks No response or no response worthy of credit.		Mass of U236 = 235.04395 + 1.00867 = 236.05262u
			Total mass of fission products = 140.91440 + 91.92617 + (3 x 1.00867) = 235.86658u
			Difference = 0.18604 u
			Mass defect =0.18604 x 1.661 x 10 ⁻²⁷ = 3.0901244 x 10 ⁻²⁸ kg
			Energy equivalence per fission event = $3.0901244 \times 10^{-28} \times (3 \times 10^{8})^{2} = 2.78111 \times 10^{-11} \text{J}$
			Alternative: Energy equivalence per fission in MeV = 174 MeV
(c)	Marks may be awarded for the following considerations only: technology, reaction, fuel, waste, safety.	4 x B1	ALLOW 1 mark for basic comparison within each consideration if no other marks are awarded.
	 Technology: Fission: technology already established / Fusion: technology not yet established 		

Question	Answer	Mark	Guidance
	 Reaction: Fission: chain reaction self-sustaining once initiated Fusion: very high temperatures/pressures /amounts of energy required to initiate and continue reaction / challenges with storage and containment of plasma 		
	 Fuel: Fission: finite amount of uranium / needs to be enriched Fusion: (unlimited) fuel available from (sea) water 		ALLOW hydrogen is easier to source (than uranium) for 1 mark
	 Waste: Fission: Produces highly / high level / long half- life / long storage time radioactive waste Fusion: Produces (effectively) no radioactive waste (products) / only radioactive material in contained in the reactor 		ALLOW Fission produces radioactive waste and fusion does not for 1 mark. IGNORE nuclear for radioactive
	 Fission: Possibility of serious accident / leakage of radioactive material / "meltdown" AND/OR Fusion: very unlikely to produce serious accident (due to near instant shut down of reaction) / no leakage of radioactive material 		

June 2	023
--------	-----

	Question	Answer	Mark	Guidance	
22	(a)	${}^{18}_{9}\text{F} \rightarrow {}^{18}_{8}\text{O} + {}^{0}_{1}\beta^{+} (+\nu)$	C1 A1	Correct fluorine isotope Correct equation ALLOW numbers written to right of symbol ALLOW e / e ⁺ / \overline{e} / β for positron symbol with correct numbers IGNORE gamma in products ALLOW $_{0}^{0}v$ for neutrino ALLOW v _e for neutrino	
	(b)	$E = mc^{2} = 9.11 \times 10^{-31} \times (3.00 \times 10^{8})^{2}$ $= 8.2 \times 10^{-14} \text{ (J)}$ $E = hc / \lambda = 8.2 \times 10^{-14} = 6.63 \times 10^{-34} \times 3 \times 10^{8} / \lambda$	C1 C1	Working leading to evaluation of E ALLOW factor of 2 leading to 1.6 x 10 ⁻¹³ (J) NOT use of $\lambda = h/mv$ where v is 3x10 ⁸ alone XP ALLOW use of $\lambda = hc/mc^2$ or $\lambda = h/mc$ ALLOW Use of electron mass = 0.511 MeV giving 8.2 x 10 ⁻¹⁴ (J)	
		$\lambda = 3.00 \times 10^8 \div 1.2367 \times 10^{20} = 2.4 \times 10^{-12} $ (m)	A1	Correct to at least 2 significant figures Answer of 4.9 x 10^{-12} or 1.2 x 10^{-12} allow 2/3 due to incorrect accounting for factor of 2	
	(c)	X-rays formed when <u>electrons</u> (in an atom) de-excite	B1	ALLOW X-rays may be produced by acceleration / deceleration of (fast moving) electrons / X-rays are produced when (fast moving) electrons are incident on a metal target / X-rays may be produced when electrons lose kinetic energy	

Question	Answer	Mark	Guidance
	Gamma rays come from the decay of <u>nuclei</u> (in unstable isotopes)	B1	ALLOW gamma rays come from (the decay of) radioactive <u>nuclei</u> / gamma rays come from the <u>nucleus</u> of unstable atoms / gamma rays come from the de- excitation of <u>nuclei</u> / gamma rays come from <u>annihilation</u> of particle-antiparticle (pairs) / IGNORE gamma from fission
(d)	The half-life is short	B1	ALLOW activity is high IGNORE it decays quickly
	Advantage: Exposure of the patient to <u>ionising</u> radiation is kept as low as possible.	B1	
	Disadvantage: (Radiographers have a) short time to scan/diagnose the patient	B1	ALLOW F18 has to be manufactured on site before use / high activity means exposure is high during handling IGNORE short time to treat the patient

	Question		Answer	Mark	Guidance
23	(a)	(i)	4.804 3.839		ALLOW 2dp for either or both values 4.80 and 3.84
		(ii)	Both points plotted correctly	A1	ALLOW plotting of their 23(a)(i) values as long as the values lie within the grid range Plotting to within ½ small square
		(iii)	Appropriate line of best fit	B1	Straight line with negative gradient Allow best fit line from their plotted points Expect even distribution of points across the whole line for their plotted points
	(b)	(i)	$\ln I_x = \ln I_0 - \mu x$	C1	ALLOW log equation in any format
			In $I_x = -\mu x + \ln I_0$ compare with y = mx + c	A1	ALLOW an indication that $y = mx + c$ has been used, with the In equation in the correct order. Negative sign for required μ .
		(ii)	Correct determination of their y intercept	C1	Their intercept recorded to nearest ½ small grid square ALLOW values to more than 1 dp
			Correct e ^{intercept} of their y intercept value	A1	Answer to 2 sf
		(iii)	Evidence of use of graph to determine gradient	B1	Change in x value greater than half of their line AND points correctly read from the line (not the tabulated values unless they lie on the line).

PMT

Question		Answer	Mark	Guidance
				¹ ⁄ ₂ square tolerance on readings of points Ignore sign on gradient.
		Correct answer for their gradient, and given as positive for $\boldsymbol{\mu}$	B1	Correct to at least 2 significant figures NOT if graph shows positive gradient
(c)	(i)	(100 / 5100) x 5 (= 0.098) (V) OR (8000 / 13000) x 5 (= 3.077) (V)	C1	ALLOW rounding of either or both of these values to 2sf
		0.098 to 3.077 OR 2.98 (V)	A1	ALLOW final answer of range of 0.1 to 3 OR 3 (1sf answer) if more than 1sf seen previously
	(ii)	Any two from: Increase the input voltage	2 x B1	ALLOW emf for input voltage but not just potential difference IGNORE current
		Decrease (the resistance of) R_1 / (fixed) resistor		NOT decrease the resistance alone
		Extend range of thickness of jelly used		NOT just make jelly thinner/thicker. Must be clear that values below 1mm or above 5mm are to be used.
	(iii)	As the light intensity increases the resistance of the LDR/ R ₂ decreases the PD measured (across R ₂) decreases OR	B1	ALLOW reverse argument Must be clear that the resistance is that of the LDR ALLOW output voltage/voltmeter reading for PD across R ₂

Mark Scheme

Question		Answer	Mark	Guidance
		As the light intensity increases the total resistance of the circuit decreases the current increases		

Need to get in touch?

If you ever have any questions about OCR qualifications or services (including administration, logistics and teaching) please feel free to get in touch with our customer support centre.

Call us on

01223 553998

Alternatively, you can email us on

support@ocr.org.uk

For more information visit



ocr.org.uk

Twitter/ocrexams

/ocrexams

/company/ocr

/ocrexams



OCR is part of Cambridge University Press & Assessment, a department of the University of Cambridge.

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored. © OCR 2023 Oxford Cambridge and RSA Examinations is a Company Limited by Guarantee. Registered in England. Registered office The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA.

Registered company number 3484466. OCR is an exempt charity.

OCR operates academic and vocational qualifications regulated by Ofqual, Qualifications Wales and CCEA as listed in their qualifications registers including A Levels, GCSEs, Cambridge Technicals and Cambridge Nationals.

OCR provides resources to help you deliver our qualifications. These resources do not represent any particular teaching method we expect you to use. We update our resources regularly and aim to make sure content is accurate but please check the OCR website so that you have the most up-to-date version. OCR cannot be held responsible for any errors or omissions in these resources.

Though we make every effort to check our resources, there may be contradictions between published support and the specification, so it is important that you always use information in the latest specification. We indicate any specification changes within the document itself, change the version number and provide a summary of the changes. If you do notice a discrepancy between the specification and a resource, please <u>contact us</u>.

Whether you already offer OCR qualifications, are new to OCR or are thinking about switching, you can request more information using our Expression of Interest form.

Please get in touch if you want to discuss the accessibility of resources we offer to support you in delivering our qualifications.