



# GCE

## Physics A

Advanced GCE

Unit **G485**: Fields, Particles and Frontiers of Physics

# Mark Scheme for June 2011

---

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications  
PO Box 5050  
Annesley  
NOTTINGHAM  
NG15 0DL

Telephone: 0870 770 6622  
Facsimile: 01223 552610  
E-mail: [publications@ocr.org.uk](mailto:publications@ocr.org.uk)

## CATEGORISATION OF MARKS

The marking schemes categorise marks on the MACB scheme.

- B** marks: These are awarded as independent marks, which do not depend on other marks. For a **B**-mark to be scored, the point to which it refers must be seen specifically in the candidate's answers.
- M** marks: These are method marks upon which **A**-marks (accuracy marks) later depend. For an **M**-mark to be scored, the point to which it refers must be seen in the candidate's answers. If a candidate fails to score a particular **M**-mark, then none of the dependent **A**-marks can be scored.
- C** marks: These are compensatory method marks which can be scored even if the points to which they refer are not written down by the candidate, providing subsequent working gives evidence that they must have known it. For example, if an equation carries a **C**-mark and the candidate does not write down the actual equation but does correct working which shows the candidate knew the equation, then the **C**-mark is given.
- A** marks: These are accuracy or answer marks, which either depend on an **M**-mark, or allow a **C**-mark to be scored.

### Note about significant figures:

Significant figures are rigorously assessed in the practical skills.

If the data given in a question is to 2 sf, then allow answers to 2 or more significant figures.

If an answer is given to fewer than 2 sf, then penalise once only in the entire paper.

Any exception to this rule will be mentioned in the Additional Guidance.

Question		Expected Answers	Marks	Additional guidance
1	(a)	Electromotive force is the energy transferred (from one form of energy) to <u>electrical per</u> unit charge	B1	<b>Allow:</b> 'electrical energy (gained) per unit charge' <b>Not:</b> electrical energy per coulomb
	(b)	Magnetic flux is the product of the (magnetic) flux density and the area (normal to the field)	B1	<b>Allow:</b> $\phi = BA$ , where $B$ = (magnetic) flux density and $A$ = area. If $\phi = BA \cos \theta$ is used, then $\theta$ must be defined as the angle (between the normal to the plane of the area and the magnetic field) <b>Do not allow</b> 'field strength' for 'flux density'
	(c) (i)	A changing (magnetic) flux is produced (in the primary coil / in the iron core)  The iron core links this (magnetic) flux / (magnetic) flux density to the secondary coils  The changing (magnetic) flux / (magnetic) flux density through secondary induces e.m.f. (in secondary coils)	B1  B1  B1	<b>Allow:</b> A changing (magnetic) flux density is produced (in the primary coil) but <b>not</b> ' <i>changing (magnetic) field</i> '  <b>Allow:</b> The rate of change of (magnetic) flux (linkage) induces an e.m.f. (in the secondary coil)
	(ii)	Any <u>one</u> from: More coils / turns on secondary Less coils / turns on primary Laminate the core	B1	<b>Not:</b> Increase frequency of alternating supply
	(d) (i)	$\frac{n_s}{4200} = \frac{12}{230}$ (Any subject) number of turns = 219 or 220	C1  A1	<b>Note:</b> A bald answer 219 or 220 scores 2 marks
	(ii)	current = $(12.0 - 11.8) / 0.35$ current = 0.57 (A) ----- $P = VI$ or $P = I^2R$ or $P = V^2 / R$ $P = 0.2 \times 0.57$ or $P = 0.57^2 \times 0.35$ or $P = 0.2^2 / 0.35$ power = 0.114 (W) or 0.11 (W)	C1 A1  C1  A1	Possible e.c.f. from (ii)1
		<b>Total</b>	<b>12</b>	

Question		Expected Answers	Marks	Additional guidance
2	(a)	capacitance = charge / potential difference	B1	<b>Allow:</b> p.d. and voltage <b>Not:</b> charge per volt or coulombs per p.d
	(b) (i)	$V = Q/C$ and $Q = \text{constant}$ in series circuit $V = \frac{450}{450 + 150} \times 6.0$ potential difference = 4.5 (V)	C1 A1	<b>Allow:</b> 1 mark for an answer of 1.5 (V) <b>Note:</b> Using (b)(ii), alternative marking scheme $V = 6.75 \times 10^{-4} / 150 \times 10^{-6}$ C1 $V = 4.5$ V A1
	(ii)	charge = $150 \times 10^{-6} \times 4.5$ charge = $6.75 \times 10^{-4}$ (C)	B1	Possible e.c.f. <b>Note:</b> Using (b)(iii) ... $Q = 6.0 \times 1.125 \times 10^{-4} = 6.75 \times 10^{-4}$ (C)
	(iii)	$\frac{1}{C} = \frac{1}{150} + \frac{1}{450}$ (working in $\mu\text{F}$ ) capacitance $C_T = 1.125 \times 10^{-4}$ (F) or 113 $\mu\text{F}$	B1	<b>Possible alternative:</b> capacitance = $6.75 \times 10^{-4} / 6.0$ capacitance = $1.125 \times 10^{-4}$ (F) or 113 $\mu\text{F}$ Possible e.c.f. from (ii)
	(c) (i)	time constant = $CR$ time constant = $1.125 \times 10^{-4} \times 45 \times 10^3$ time constant = 5.06 (s)	M1 A0	<b>Note:</b> The mark is for multiplying correct $C$ and $R$ values Possible e.c.f. from (b)(iii)
	(ii)	Graph starting from 6.0 (V)  Correct shaped curve  Approximately correct value of $V$ at $CR$	B1  B1  B1	<b>Note:</b> The (exponential decay) curve must not touch or cut the time axis  <b>Note:</b> $V$ is 2 to 2.5 (V) at $t \approx 5$ s

G485

Mark Scheme

June 2011

Question		Expected Answers	Marks	Additional guidance
	(iii)	$\frac{1}{2} \times 4.5^2 \times 150 \times 10^{-6}$ <u>and</u> $\frac{1}{2} \times 1.5^2 \times 450 \times 10^{-6}$ $\text{ratio} = \frac{0.5 \times 4.5^2 \times 150 \times 10^{-6}}{0.5 \times 1.5^2 \times 450 \times 10^{-6}}$ $\text{ratio} = 3$ <p style="text-align: center;">Or</p> $\frac{1}{2} Q^2 / C_{150} \text{ and } \frac{1}{2} Q^2 / C_{450}$ $\text{ratio} = C_{450} / C_{150}$ $\text{ratio} = 3$	C1  A1  C1  A1	<b>Allow:</b> with or without the $10^{-6}$ Possible e.c.f. from <b>(b)(i)</b> and <b>(b)(ii)</b> <b>Allow:</b> full credit for correct use of either $\frac{1}{2} QV$ or $\frac{1}{2} Q^2 / C$
	(iv)	The ratio remains constant The charge / Q is the same for both capacitors	B1 B1	
<b>Total</b>			<b>13</b>	

Question		Expected Answers	Marks	Additional guidance
3	(a)	(Electric field strength is the) force <u>per</u> (unit positive) charge	B1	<b>Allow:</b> $E = F/Q$ , $F$ is the force on a (positive) charge $Q$
	(b)	Parallel and equally spaced lines at right angles to plates  Correct <u>upward</u> direction of field shown on at least one field line	B1  B1	
	(c) (i)	An arrow vertically downwards at <b>P</b>	B1	
	(ii)	$E = \frac{3400}{0.050} \quad \text{or} \quad E = 6.8 \times 10^4 \text{ (V m}^{-1}\text{)}$ $a = \frac{EQ}{m}$ $a = \frac{6.8 \times 10^4 \times 1.6 \times 10^{-19}}{9.11 \times 10^{-31}} \quad \text{or} \quad a = \frac{1.09 \times 10^{-14}}{9.11 \times 10^{-31}}$ acceleration = $1.19 \times 10^{16} \text{ (m s}^{-2}\text{)}$ or $1.2 \times 10^{16} \text{ (m s}^{-2}\text{)}$	C1  C1  A0	<b>Vital:</b> Candidates using separation of 0.050 cm must be awarded full credit for the analysis shown below $E = \frac{3400}{0.050 \times 10^{-2}} \quad \text{or} \quad E = 6.8 \times 10^6 \text{ (V m}^{-1}\text{)} \quad \text{C1}$ $a = \frac{EQ}{m}$ $a = \frac{6.8 \times 10^6 \times 1.6 \times 10^{-19}}{9.11 \times 10^{-31}}$ acceleration = $1.19 \times 10^{18} \text{ (m s}^{-2}\text{)}$ <span style="float: right;">C1 A0</span>
	(iii)	$t = \frac{0.04}{4.0 \times 10^7}$ time = $1.0 \times 10^{-9} \text{ (s)}$	B1	<b>Allow:</b> $1 \times 10^{-9} \text{ (s)}$ or $10^{-9} \text{ (s)}$
	(iv)	initial vertical velocity = 0, final vertical velocity = $at$  vertical velocity = $1.2 \times 10^{16} \times 1.0 \times 10^{-9}$ <b>(Allow:</b> $1 \times 10^{16} \times 1.0 \times 10^{-9}$ ) vertical velocity = $1.2 \times 10^7 \text{ (m s}^{-1}\text{)}$	M1  A0	<b>Vital:</b> Candidates using separation of 0.050 cm must be awarded full credit for the analysis shown below vertical velocity = $1.2 \times 10^{18} \times 1.0 \times 10^{-9}$ <span style="float: right;">M1</span>  vertical velocity = $1.2 \times 10^9 \text{ (m s}^{-1}\text{)}$ <span style="float: right;">A0</span>

G485

Mark Scheme


June 2011

Question		Expected Answers	Marks	Additional guidance
	(v)	$v^2 = (4.0 \times 10^7)^2 + (1.2 \times 10^7)^2$ velocity = $4.2 \times 10^7$ (m s <sup>-1</sup> ) Or $v^2 = (4.0 \times 10^7)^2 + (1 \times 10^7)^2$ velocity = $4.1 \times 10^7$ (m s <sup>-1</sup> )	C1 A1  C1 A1	Possible ecf from (iv)
	(vi)	$KE = \frac{1}{2} mv^2$ $KE = 0.5 \times 9.11 \times 10^{-31} \times (4.2 \times 10^7)^2$ kinetic energy = $8.04 \times 10^{-16}$ (J) or $8.0 \times 10^{-16}$ (J)	C1 A1	Possible ecf from (v) <b>Allow:</b> 1 sf answer if the answer comes out as $8.0 \times 10^{-16}$ (J)
	(vii)	Graph starts at non-zero value for $E_k$  Between 0 and 0.08 (m) the graph has increasing gradient  Horizontal line after 0.080 (m)	B1  B1  B1	<b>Note:</b> The $E_k$ value for the horizontal line > $E_k$ value at $x = 0$
<b>Total</b>			<b>15</b>	



Question		Expected Answers	Marks	Additional guidance
4	(a)	$E = \frac{Q}{4\pi\epsilon_0 r^2}$ $\frac{(-)4.0 \times 10^{-9}}{4\pi\epsilon_0 \times (1.75 \times 10^{-2})^2} \text{ and } \frac{5.0 \times 10^{-9}}{4\pi\epsilon_0 \times (1.75 \times 10^{-2})^2}$ $E_B = 1.17 \times 10^5 \text{ (N C}^{-1}\text{)} \text{ and } E_A = 1.47 \times 10^5 \text{ (N C}^{-1}\text{)}$ <p>field strength = <math>(1.17 + 1.47) \times 10^5 \text{ (N C}^{-1}\text{)}</math></p> <p>field strength = <math>2.64 \times 10^5 \text{ (N C}^{-1}\text{)}</math> or <math>2.6 \times 10^5 \text{ (N C}^{-1}\text{)}</math></p> <p>direction = to the left / towards B</p>	C1  C1  A1  B1	Ignore signs  <b>Allow:</b> 2 marks for $2.9(4) \times 10^4 \text{ (N C}^{-1}\text{)}$ when the fields are subtracted <b>Allow:</b> 2 marks for $6.6 \times 10^4 \text{ (N C}^{-1}\text{)}$ for using $3.5 \times 10^{-2} \text{ m}$
	(b)	$F = \frac{Qq}{4\pi\epsilon_0 r^2}$ $\text{force} = \frac{4.0 \times 10^{-9} \times 5.0 \times 10^{-9}}{4\pi \times 8.85 \times 10^{-12} \times (3.5 \times 10^{-2})^2}$ $\text{force} = 1.47 \times 10^{-4} \text{ (N)}$	C1  C1 A0	Ignore signs <b>Allow:</b> $\epsilon_0$ in the equation
	(c)	<p>(weight =) <math>4.5 \times 10^{-5} \times 9.81</math> or (weight =) <math>4.4(1) \times 10^{-4} \text{ (N)}</math></p> $\tan \theta = \frac{1.5 \times 10^{-4}}{4.41 \times 10^{-4}}$ <p>angle = <math>18.8^\circ</math> or <math>19^\circ</math></p> <p><b>(Allow:</b> Full credit when angle is determined using a scale diagram)</p>	C1  C1  A1	<b>Allow:</b> weight = $4.5 \times 10^{-5} \times g$  <b>Note:</b> Using force = $1.47 \times 10^{-4} \text{ (N)}$ gives an angle of $18.4^\circ$ ; hence allow $18^\circ$ <b>Allow:</b> 2 marks for $\theta = 71^\circ$ ; this is the complementary angle <b>Allow:</b> 1 mark for ' $\tan \theta = \frac{1.5 \times 10^{-4}}{4.5 \times 10^{-5}}, \theta = 73^\circ$ ', when mass is used instead of weight.
<b>Total</b>			<b>9</b>	

Question		Expected Answers	Marks	Additional guidance
5	(a)	Down(wards)	B1	<b>Note:</b> Can be on Fig. 5.1
	(b)	(Fleming's) left-hand rule	B1	<b>Allow:</b> Thumb in direction of force, first finger in direction of (magnetic) field and second finger in direction of (conventional) current
	(c) (i)	force = $BIL = 0.080 \times 4.0 \times 5.0 \times 10^{-2}$ force = 0.016 (N)	B1	
	(ii)	reading = 2.500 – 0.016 reading = 2.484 (N)  The force on <u>core/magnets</u> is up(wards)  (According to Newton's third law) the forces (on the rod and steel core/magnets) are equal <u>and</u> opposite	B1  B1  B1	<b>Allow:</b> 'up and down' as equivalent to 'opposite'
	(d)	Resistance increases by a factor of 4  Current decreases by a factor of 4  The force decreases by a factor of 4  force = 0.004 (N)	C1  C1  A1	Possible e.c.f. from <b>(c)(i)</b> <b>Note:</b> force = <b>(c)(i)</b> /4 can score full marks <b>Special case:</b> Allow 1 mark for (resistance doubles, current is halved, hence) force = 0.008 (N)
		<b>Total</b>	<b>9</b>	

Question		Expected Answers	Marks	Additional guidance
6	(a)	<p>Any <u>four</u> from 1 to 5:</p> <ol style="list-style-type: none"> <li>1. Most of the alpha particles went straight through (some deviated through small angles)</li> <li>2. Hence most of the atom is empty space</li> <li>3. Some / a very small number of alpha particles were scattered / repelled through large angles / angles more than 90°</li> <li>4. This showed the existence of (a tiny) positive nucleus</li> <li>5. The size of the nucleus is about <math>10^{-14}</math> <u>m</u></li> </ol> <p> QWC: Award a mark for one conclusion correctly linked to an observation</p>	<p>B1×4</p>          <p>B1</p>	<p><b>Must use ticks on Scoris to show where the marks are awarded</b></p>          <p><b>Allow:</b> <math>10^{-15}</math> <u>m</u></p>
	(b)	<p>Any <u>five</u> from:</p> <p>Gravitational (force) This force is attractive <b>AND</b> is long-ranged / obeys ‘<math>1/r^2</math> relationship’</p> <p><u>Strong</u> (nuclear force/interaction) This force is attractive (at larger distances) or repulsive at short distances <b>AND</b> is short-ranged / <math>\sim 10^{-14}</math> m</p> <p>Electrostatic / electrical (force) / coulomb (force) This force is repulsive between protons / zero between neutrons / zero between protons and neutrons <b>AND</b> is long-ranged / obeys ‘<math>1/r^2</math> relationship’</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p><b>Allow:</b> gravity</p> <p><b>Note:</b> Do not allow ‘inverse square law’; allow ‘inverse square law with distance’</p>          <p><b>Allow:</b> Electromagnetic (interaction/force)</p>

Question		Expected Answers	Marks	Additional guidance	
	(c)	(i)	<p>mass = <math>235 \times 1.7 \times 10^{-27}</math> (= <math>3.995 \times 10^{-25}</math> kg)</p> <p>volume = <math>\frac{4}{3} \pi \times (8.8 \times 10^{-15})^3</math> (= <math>2.855 \times 10^{-42}</math> m<sup>3</sup>)</p> <p>density = mass/volume</p> <p>density = <math>1.4 \times 10^{17}</math> (kg m<sup>-3</sup>)</p>	<p>C1</p> <p>C1</p> <p>A1</p>	<p><b>Allow:</b> <math>1.66 \times 10^{-27}</math> kg for mass of nucleon</p> <p><b>Allow:</b> <math>10^{17}</math> (kg m<sup>-3</sup>) for this estimation question  <b>Note:</b> Omitting 235 gives <math>6.0 \times 10^{14}</math> (kg m<sup>-3</sup>), allow 2 mark  <b>Allow:</b> 1 mark if 92 or 143 is used to determine the mass of the nucleus; this gives a density value of <math>5.5 \times 10^{16}</math> (kg m<sup>-3</sup>) and <math>8.5 \times 10^{16}</math> (kg m<sup>-3</sup>) respectively</p>
		(ii)	The nucleons / neutrons and protons are packed together with little or no empty space (AW)	B1	
			<b>Total</b>	14	

Question		Expected Answers	Marks	Additional guidance
7	(a)	The critical density is the density for which the universe will expand towards a (finite) limit or rate of expansion <b>tends</b> to zero / which will result in a <u>flat</u> universe	B1	<b>Not:</b> critical density is given by $\frac{3H_0^2}{8\pi G}$
	(b)	Hubble constant = $\frac{65 \times 10^3}{10^6 \times 3.1 \times 10^{16}}$ Hubble constant = $2.1 \times 10^{-18} \text{ s}^{-1}$ critical density = $\frac{3H_0^2}{8\pi G}$ critical density = $\frac{3 \times (2.1 \times 10^{-18})^2}{8\pi \times 6.67 \times 10^{-11}}$ critical density = $7.9 \times 10^{-27} \text{ (kg m}^{-3}\text{)}$	B1  C1  A1	Possible e.c.f. from value of Hubble constant within this calculation
	(c) (i)	open: (density of universe < critical density hence) the universe will expand forever  closed: (density of universe > critical density hence) the universe will (eventually stop expanding and then) contract / big crunch  flat: (density of universe = critical density hence) the universe will expand towards a (finite) limit / rate of expansion <b>tends</b> to zero	B1  B1  B1	<b>Allow:</b> 'universe continues to expand'   <b>Not:</b> 'The universe stops expanding'  <b>Special case:</b> Award 1 mark for correct sketches if no explanation is given for open, closed and flat
	(ii)	Any <u>one</u> from: Existence of dark matter / black holes / neutrinos / dark energy / $H_0$ is not known accurately	B1	
<b>Total</b>			<b>8</b>	

Question		Expected Answers	Marks	Additional guidance
8	(a)	Less chance of infection	B1	
	(b)	Any <u>two</u> from:  1. Tracer is injected into the body / placed inside the body / circulates the body 2. Tracer is absorbed by organ / shows blockage 3. Beta detector / gamma camera (is used to detect radiation from the body)	B1×2	<b>Note:</b> No marks for ingesting substances (e.g barium)
	(c)	Any <u>five</u> from:  1. A positron / beta-plus emitting tracer / source is used 2. The positron annihilates with an electron (inside the patient) 3. This produces <u>two</u> gamma photons 4. The photons travels in opposite directions 5. The patient is surrounded by a ring of gamma detectors 6. The arrival times of the photons / delay time indicates location (of tumour inside the body) 7. A 3-D image is created (by the computer connected to the detectors)	B1×5	
		<b>Total</b>	<b>8</b>	



Question		Expected Answers	Marks	Additional guidance
10	(a)	A neutron is absorbed by a (massive / uranium) nucleus	B1	
		The nucleus splits into two (smaller/daughter) nuclei and (one or more) neutrons	B1	
	(b)	In a fission reaction there is a decreases in the mass	M1	<b>Allow:</b> The 'BE increases (in the reaction)'
		(According to $\Delta E = \Delta mc^2$ ) mass is converted into energy	A1	
		Or The (total) binding energy of the products / smaller nuclei is greater than the binding energy of the original nucleus	M1	
		The difference in the binding energies is released as energy	A1	
	(c)	Moderator: water / graphite / carbon	B1	<b>Note:</b> If boron is mentioned, then do not award this B1 mark  <b>Allow:</b> They become thermal neutrons
		It slows down the (fast-moving) neutrons / reduces the (kinetic) energy of neutrons	B1	
		Slow-moving neutrons have greater chance of causing fission (than fast-moving neutrons)	B1	
		<b>Total</b>	<b>7</b>	



**OCR (Oxford Cambridge and RSA Examinations)**  
**1 Hills Road**  
**Cambridge**  
**CB1 2EU**

**OCR Customer Contact Centre**

**14 – 19 Qualifications (General)**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

**[www.ocr.org.uk](http://www.ocr.org.uk)**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
**is a Company Limited by Guarantee**  
**Registered in England**  
**Registered Office; 1 Hills Road, Cambridge, CB1 2EU**  
**Registered Company Number: 3484466**  
**OCR is an exempt Charity**

**OCR (Oxford Cambridge and RSA Examinations)**  
**Head office**  
**Telephone: 01223 552552**  
**Facsimile: 01223 552553**

