| Surname | Centre Number | Candidate Number |
|-------------|------------------|---------------------|
| Other Names | | 2 |



GCE AS

B420U20-1

S18-B420U20-1



PHYSICS – AS component 2 Electricity and Light

FRIDAY, 18 MAY 2018 - MORNING

1 hour 30 minutes

| For Examiner's use only | | | |
|-------------------------|-----------------|-----------------|--|
| Question | Maximum Mark | Mark Awarded | |
| 1. | 13 | | |
| 2. | 10 | | |
| 3. | 19 | | |
| 4. | 8 | | |
| 5. | 11 | | |
| 6. | 14 | | |
| Total | 75 | | |

ADDITIONAL MATERIALS

In addition to this paper, you will require a calculator and a Data Booklet.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page. Answer **all** questions.

Write your answers in the spaces provided in this booklet.

INFORMATION FOR CANDIDATES

The total number of marks available for this paper is 75.

The number of marks is given in brackets at the end of each question or part-question.

You are reminded to show all working. Credit is given for correct working even when the final answer is incorrect.

The assessment of the quality of extended response (QER) will take place in **Q5**(*a*).



Examiner only A section of rock undergoes a tensile stress of 900 MPa during an earthquake. Calculate the tensile strain if the Young modulus is 70 GPa for rock. [3] (C) Explain how data obtained by geologists about earthquakes from various monitoring (d) stations can benefit society. [2]

3

B420U201 03

13

2.

| A circ | evit is set up as shown. A $6.0V$ A 6.8Ω C 6.8Ω | Examine |
|--------|---|---------|
| (a) | In the circuit shown, the potential difference between X and Y is 6.0V. Explain what this statement means. [2] (i) Determine the reading on the ammeter if it has an instrument resolution of ± 0.01A. [4] | |
| | (ii) Calculate the potential difference across the 8.2Ω resistor. [2] | |

| (iii) | Calculate the power dissipated in the parallel resistor combination. | [2] | Examiner only |
|-------|--|-----|------------------|
| | | | |
| ••••• | | | |
| ••••• | | | |

10

B420U201 05

BLANK PAGE



[3]

7

(ii) Einstein's photoelectric equation may be written as:

State what is meant by the photoelectric effect.

3.

(a)

(i)

$$E_{k\max} = hf - \phi$$

Explain this equation in terms of energy.

(b) Evaluate why the following circuit is incorrect and cannot be used to measure the maximum kinetic energy of the emitted electrons. [4]



© WJEC CBAC Ltd. (B420U20-1)

(c) Different frequencies of light are used with a **correct** circuit and the following results are only only

| Frequency/10 ¹⁴ Hz | 5.1 | 6.0 | 6.9 | 7.5 |
|-------------------------------|------|------|------|------|
| $E_{kmax}/10^{-19} J$ | 0.36 | 0.93 | 1.50 | 1.95 |

(i) Plot E_{kmax} (*y*-axis) against frequency (*x*-axis) on the grid below and draw a line of best fit through your data. [3]



| Explain whether or not your graph is in agreement with Einstein's photoelectric equation. [3] | Examiner only |
|---|--|
| | |
| Calculate a value for the Planck constant using the gradient of your graph. [3] | |
| An answer can be considered to be accurate if it is within 5% of the accepted value for the Planck constant. Evaluate whether your answer for the Planck constant can be considered to be accurate. | |
| | |
| | Explain whether or not your graph is in agreement with Einstein's photoelectric equation. [3] Calculate a value for the Planck constant using the gradient of your graph. [3] An answer can be considered to be accurate if it is within 5% of the accepted value for the Planck constant. Evaluate whether your answer for the Planck constant can be considered to be accurate. [2] |

Examiner only

[1]

10

Define electric current.

4.

(a)





(i) Describe how the **current** varies from t = 0 to t = 2.5 s giving appropriate values. [4]

| (ii) Calculate the current when $t = 3.0$ s. [3] | Examiner only |
|---|------------------|
| | |
| | |
| | |
| | |

Examiner

only

5.

(a)

Explain how a 3 level laser operates, explaining also why a 2 level laser is not possible. [6 QER]





6. (a) A multimode optical fibre has a core made of glass of refractive index 1.52. The cladding is made of a material with refractive index 1.47.

| ir | A 15° A | Cladding $n = 1.47$ Core $n = 1.52$ Cladding $n = 1.47$ |
|----------|---|---|
| (i) | Calculate the critical angle for the core-cladding bounda | ry. [2] |
| (ii) | A beam of light enters the optical fibre from air at an angle angle A. | of 15° as shown. Calculate [3] |
| (iii) | A technician states that the beam of light entering the fi 15° will not travel down the optical fibre. Evaluate whethe | bre from air at an angle of er the technician is correct. [2] |
| | | |

| 15 | | | |
|-----|---------------|--|--------|
| (b) | Calc lengt | ulate the time taken for the light to travel along the axis of a straight optical fibre c th 15 km. | f |
| | | | |
| (C) | (i) | State how the paths of light in monomode and multimode optical fibres differ. [1 | |
| | (ii) | Explain the advantage of monomode optical fibres over multimode optical fibres fo communicating a rapid sequence of data encoded as light pulses. [3 | r] |
| | | | |
| | ······ | | |
| | | END OF PAPER | |