



GCE Physics - PH4

January 2013 - Markscheme

Question		Marking details	Marks Available
1	(a)	(i) $T = \frac{1}{f} = 1.6$ or $\omega^2 = \frac{k}{m}$ (1) algebra i.e. $m = \frac{T^2 k}{4\pi^2}$ or $\omega = 2\pi f$ (1) $m = \frac{1.6^2 \times 2640}{4\pi^2}$ (1) = [171 kg]	[3]
		(ii) $\frac{1}{2}mv^2 = 2150$ (1) $v = 5.01 \text{ [m s}^{-1}\text{]}$ (1) ecf on m	[2]
		(iii) 2.15 [kJ] (1) conservation of energy stated or implied / all KE transferred to PE (1) (accept energy cannot be created or destroyed)	[2]
		(iv) $v = \omega A$ (1) or suitable alternative $A = 1.28 \text{ [m]}$ (1) ecf	[2]
		(v) $x = \pm A \sin(2\pi ft)$ (1) For 1 st mark ω must be substituted. $a = -\omega^2 x$ used (1) $13.9 \text{ [m s}^{-2}\text{]}$ (1) ecf	[3]
		(b) Resonance / maximum amplitude (1) since natural frequency / $\frac{1}{0.625} = 1.6$ (1)	[2]

Question		Marking details	Marks Available
	(c)	<p>Basic shape (decreasing to 1.4 m with a cos or -cos shape) (1)</p> <p>period = 1.6 s (accept 1.5 – 1.7 s) (1)</p> <p>period constant (1)</p> <div style="text-align: center;">  </div> <p>Question 1 total</p>	<p>[3]</p> <p>[17]</p>
2	(a)	<p>$\frac{1}{2} m \overline{c^2}$ KE of a particle/atom/molecule (1)</p> <p>$\frac{3}{2} nRT$ internal energy (accept total KE) (1)</p>	<p>[1]</p> <p>[1]</p>
	(b)	<p>(i)</p> <p>$N_A \times \frac{1}{2} m \overline{c^2} = \frac{3}{2} \times 1 \times RT$ (1) (or equivalent)</p> <p>e.g. $\frac{1}{2} m \overline{c^2} = \frac{3}{2} kT$</p> <p>$\overline{c^2} = \frac{3RT}{mN_A}$ (1) (i.e. algebra)</p> <p>rms speed = 1350 [m s⁻¹] (1)</p> <p>(ii)</p> <p>$p = \frac{1}{3} \rho \overline{c^2}$ (1)</p> <p>$p = 1.16 \times 10^5$ Pa / Nm⁻² (1) ecf UNIT mark</p> <p>Or suitable alternative method</p> <p>Question 2 total</p>	<p>[3]</p> <p>[2]</p> <p>[7]</p>

Question		Marking details	Marks Available
3	(a)	The [vector] sum of the momenta [of bodies in a system] stays constant [even if forces act between the bodies], (1) provided there is no external [resultant] force. (1)	[2]
	(b)	(i) $1.78 \times 10^{-25} \times u = 5.62 \times 10^5 \times 1.71 \times 10^{-25} \pm 1.36 \times 10^7 \times 6.64 \times 10^{-27}$ (1) i.e. attempt at conservation of momentum $u = \{5.62 \times 10^5 \times 1.71 \times 10^{-25} - 1.36 \times 10^7 \times 6.64 \times 10^{-27}\} / 1.78 \times 10^{-25}$ i.e. correct algebra and sign (1) $u = 32\,600 \text{ [ms}^{-1}\text{]} (1)$	[3]
		(ii) $E = \frac{hc}{\lambda}$ {or $E = hf$ and $c = f\lambda$ } (1) Algebra and $p = \frac{h}{\lambda}$ (1) (Use of both $E = mc^2$ and $p = mc$ award 1 mark only.)	[2]
		(iii) $p = \frac{E}{c}$ attempted (1) $5.62 \times 10^5 \times 1.71 \times 10^{-25}$ used as a denominator (1) $\frac{6.93 \times 10^{-22}}{5.62 \times 10^5 \times 1.71 \times 10^{-25}} \times 100 = 0.72\%$ (1) (accept: $4.5 \times 10^{18}\%$)	[3]
		Question 3 Total	[10]

Question	Marking details	Marks Available
4	<p>(a) horizontal arrow to right at P (1)</p> <p>both other arrows correct direction (1)</p>  <p>(b) $E = \frac{Q}{4\pi\epsilon_0 r^2}$ used (1) e.g. $\frac{6 \times 9 \times 10^9}{3^2}$</p> <p>$E = 6000 \text{ N C}^{-1}$ (1) UNIT mark</p> <p>(c) $E = \frac{Q}{4\pi\epsilon_0 r^2}$ used for negative charge (1) (answer = 1 800)</p> <p>e.g. $\frac{5 \times 9 \times 10^9}{5^2}$ but not $\frac{5 \times 9 \times 10^9}{3^2}$</p> <p>x 2 and x cos θ (1) [= 2 160]</p> <p>resultant = 3 840 [N C⁻¹] [to the right] (1) ecf on arrows</p> <p>(d) (i) correct equation used (1) e.g. $\frac{5 \times 9 \times 10^9}{5}$</p> <p>Attempt at adding 3 potentials (1) e.g. $\frac{(6-5-5) \times 9 \times 10^9}{5}$</p> <p>$\frac{1}{4\pi\epsilon_0} \left\{ \frac{6}{3} - \frac{5}{5} - \frac{5}{5} \right\}$ (1) or equivalent obviously giving zero</p> <p>(ii) (Energy) - final total energy must be zero or final potential is also zero (1) (any implied dissipation of energy loses this mark)</p> <p>Initially (resultant) <u>force / field</u> is to the right (1)</p> <p>Then (resultant) force / field is to the left or deceleration (1)</p> <p>Question 4 Total</p>	<p>[2]</p> <p>[2]</p> <p>[3]</p> <p>[3]</p> <p>[3]</p> <p>[3]</p> <p>[13]</p>

Question		Marking details	Marks Available
5	(a)	$\frac{\Delta\lambda}{\lambda} = \frac{v}{c}$ used (1) $\Delta\lambda = \frac{9.4 \times 10^5}{3 \times 10^8} \times 656 = 2.06 \text{ [nm]} (1)$ $\Delta\lambda = \frac{6.6 \times 10^5}{3 \times 10^8} \times 656 = 1.44 \text{ [nm]} (1)$	[3]
	(b)	$F = \frac{GMm}{r^2}$ used or $g = \frac{GM}{r^2}$ (1) $F = 2.37 \times 10^{-11} \text{ [N]} (1)$	[2]
	(c)	(i) $\frac{mv^2}{r} = \frac{GMm}{r^2}$ (1) convincing algebra (1)	[2]
		(ii) $v = \sqrt{\frac{GM}{r}} = \sqrt{\frac{6.67 \times 10^{-11} \times 8 \times 10^{39}}{1.5 \times 10^{20}}}$ or calculating M using v (1st mark algebra) (1) $v = 60\,000 \text{ [ms}^{-1}\text{]} \text{ or } M = 4.4 \times 10^{40} \text{ or } G = 3.675 \times 10^{-10} (1)$ Comment: (1) allow ecf If v - suggests dark matter since actual v is greater If M - yes If G - yes because larger G or stronger gravity	[3]
		Question 5 Total	[10]

Question		Marking details	Marks Available
6	(a)	<p>period = 44 [days] \pm 2 days (1)</p> <p>correct conversion to seconds (allow ecf) (1) (= 3.83x10⁶ s)</p>	[2]
	(b)	<p>$v = \frac{2\pi r}{T}$ or equivalent e.g. $v = \omega r$ and $\omega = \frac{2\pi}{T}$ (1)</p> <p>$r = \frac{vT}{2\pi} = \frac{18xa}{2\pi}$ (1) (=1.097 x 10⁷) ecf on T</p>	[2]
	(c)	<p>$d^3 = \frac{T^2 G(M_1 + M_2)}{4\pi^2}$ i.e. algebra nearly complete (1)</p> <p>$(M_1 + M_2) \approx M_1$ either written or worded (1)</p> <p>$d = 3.6 \times 10^{10}$ [m] (1) ecf</p>	[3]
	(d)	<p>Values substituted correctly into a correct equation (1)</p> <p>$M_2 = 5.9 \times 10^{26}$ [kg] (1) ecf on d and r</p> <p>i.e. 100 times / [much] larger than the Earth (1) (allow ecf on M)</p> <p>Question 6 Total</p>	[3]
			[10]

Question		Marking details	Marks Available																													
7	(a)	$T = \frac{pV}{nR}$ or implied (1) $T = \frac{84000 \times 2}{49.3 \times 8.31} = 410 \text{ [K]}$ and $T = \frac{104000 \times 1.2}{49.3 \times 8.31} = 305 \text{ [K]}$ (1)	[2]																													
	(b)	(i) $U = 190 \text{ [kJ]}$ allow ecf	[1]																													
		(ii) $U = 250 \text{ [kJ]}$ allow ecf	[1]																													
	(c)	no area under graph or no change in volume	[1]																													
	(d)	temp constant / internal energy only depends on temperature / because they are isotherms	[1]																													
	(e)	(i) A clear valid method (remember show that...) e.g. trapezium (1) (counting squares ok) $DA = \frac{1}{2} (140\,000 + 84\,000) \times 0.8 = 89.6 \text{ [kJ]}$ (1) or better $\frac{1}{2} (140\,000 + 105\,000) \times 0.4$ (no penalty for mysterious -ve sign or +ve sign) $+ \frac{1}{2} (105\,000 + 84\,000) \times 0.4 = \pm 86.8 \text{ [kJ]}$	[2]																													
(ii) $BC = \frac{1}{2} (104\,000 + 64\,000) \times 0.8 = 67.2 \text{ [kJ]}$ (1) or better $\frac{1}{2} (104\,000 + 78\,000) \times 0.4$ (sign penalised here!) $\frac{1}{2} (78\,000 + 64\,000) \times 0.4 = 64.8 \text{ [kJ]}$		[1]																														
(f)	Allow ecf <table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <thead> <tr> <th></th> <th>AB</th> <th>BC</th> <th>CD</th> <th>DA</th> <th>ABCD</th> </tr> </thead> <tbody> <tr> <td>W</td> <td>0</td> <td>67[kJ]</td> <td>0</td> <td>-90 kJ</td> <td>-23[kJ]</td> </tr> <tr> <td>ΔU</td> <td>-60 [kJ]</td> <td>0</td> <td>60 [kJ]</td> <td>0</td> <td>0</td> </tr> <tr> <td>Q</td> <td>-60 [kJ]</td> <td>67[kJ]</td> <td>60 [kJ]</td> <td>-90 kJ</td> <td>-23[kJ]</td> </tr> <tr> <td></td> <td>(1)</td> <td>(1)</td> <td>(1)</td> <td></td> <td>(1)</td> </tr> </tbody> </table>		AB	BC	CD	DA	ABCD	W	0	67[kJ]	0	-90 kJ	-23[kJ]	ΔU	-60 [kJ]	0	60 [kJ]	0	0	Q	-60 [kJ]	67[kJ]	60 [kJ]	-90 kJ	-23[kJ]		(1)	(1)	(1)		(1)	[4]
	AB	BC	CD	DA	ABCD																											
W	0	67[kJ]	0	-90 kJ	-23[kJ]																											
ΔU	-60 [kJ]	0	60 [kJ]	0	0																											
Q	-60 [kJ]	67[kJ]	60 [kJ]	-90 kJ	-23[kJ]																											
	(1)	(1)	(1)		(1)																											
Question 7 Total			[13]																													



WJEC
245 Western Avenue
Cardiff CF5 2YX
Tel No 029 2026 5000
Fax 029 2057 5994
E-mail: exams@wjec.co.uk
website: www.wjec.co.uk